

## CSCI: Regular Exercise Set 2

Prepared by Yufei Tao

**Problem 1.** Prove  $30\sqrt{n} = O(\sqrt{n})$ .

**Solution.** Set  $c_1 = 30$  and  $c_2 = 1$ . The inequality  $30\sqrt{n} \leq c_1\sqrt{n}$  holds for all  $n \geq c_2$ . This completes the proof.

**Problem 2.** Prove  $\sqrt{n} = O(n)$ .

**Solution.** Set  $c_1 = 1$  and  $c_2 = 1$ . The inequality  $\sqrt{n} \leq c_1 n$  holds for all  $n \geq c_2$ . This completes the proof.

**Problem 3.** Let  $f(n)$ ,  $g(n)$ , and  $h(n)$  be functions of integer  $n$ . Prove: if  $f(n) = O(g(n))$  and  $g(n) = O(h(n))$ , then  $f(n) = O(h(n))$ .

**Solution.** Since  $f(n) = O(g(n))$ , there exist constants  $c_1, c_2$  such that for all  $n \geq c_2$ , it holds that

$$f(n) \leq c_1 g(n).$$

Similarly, since  $g(n) = O(h(n))$ , there exist constants  $c'_1, c'_2$  such that for all  $n \geq c'_2$ , it holds that

$$g(n) \leq c'_1 h(n).$$

Set  $c''_1 = c_1 c'_1$  and  $c''_2 = \max\{c_2, c'_2\}$ . From the above, we know that for all  $n \geq c''_2$ , it holds that

$$f(n) \leq c_1 g(n) \leq c_1 c'_1 h(n) = c''_1 h(n).$$

Therefore,  $f(n) = O(h(n))$ .

**Problem 4.** Prove  $(2n + 2)^3 = O(n^3)$ .

**Solution.** Set  $c_1 = 4^3$  and  $c_2 = 1$ . The inequality  $(2n + 2)^3 \leq c_1 n^3$  holds for all  $n \geq c_2$ . This completes the proof.

**Problem 5.** Prove or disprove:  $4^n = O(2^n)$ .

**Solution.** Consider the ratio  $4^n/2^n$ , which equals  $2^n$ . The ratio clearly goes to  $\infty$  when  $n$  tends to  $\infty$ . Therefore, the statement is incorrect.

**Problem 6.** Prove or disprove:  $\frac{1}{n} = O(1)$ .

**Solution.** Set  $c_1 = 1$  and  $c_2 = 1$ . The inequality  $1/n \leq c_1 \cdot 1$  holds for all  $n \geq c_2$ . This completes the proof.

**Problem 7.** Prove that if  $k \log_2 k = \Theta(n)$ , then  $k = \Theta(n/\log n)$ .

**Solution.** Since  $k \log_2 k = O(n)$ , there exist constants  $c_1, c_2$  such that  $k \log_2 k \leq c_1 n$  for all  $n \geq c_2$ . On the other hand,  $k \log_2 k = \Omega(n)$  indicates the existence of constants  $c'_1, c'_2$  such that  $k \log_2 k \geq c'_1 n$  for all  $n \geq c'_2$ . Therefore, for all  $n \geq \max\{c_2, c'_2\}$ , we have:

$$c'_1 n \leq k \log_2 k \leq c_1 n. \quad (1)$$

Set  $c''_2 = \max\{c_1, c_2, c'_2\}$ .

When  $n \geq \max\{c''_2, (1/c'_1)^2\}$ , we derive from (1):

$$\begin{aligned} \log_2(c'_1 n) &\leq \log_2(k \log_2 k) \leq \log_2(c_1 n) \\ \Rightarrow \log_2 c'_1 + \log_2 n &\leq \log_2 k + \log_2 \log_2 k \leq \log_2 c_1 + \log_2 n \\ \Rightarrow \begin{cases} \log_2 k \leq \log_2 c_1 + \log_2 n \leq 2 \log_2 n & (\text{using } n \geq c_1) \\ 2 \log_2 k \geq \log_2 k + \log_2 \log_2 k \geq \log_2 c'_1 + \log_2 n \geq \frac{1}{2} \log_2 n \end{cases} \\ \Rightarrow \frac{\log_2 n}{4} &\leq \log_2 k \leq 2 \log_2 n. \end{aligned} \quad (2)$$

Combining (1) and (2) leads to

$$\begin{cases} k \leq c_1 \frac{n}{\log_2 k} \leq 4c_1 \frac{n}{\log_2 n} \\ k \geq c'_1 \frac{n}{\log_2 k} \geq \frac{c'_1}{2} \frac{n}{\log_2 n} \end{cases}$$

which means  $k = \Theta(n/\log n)$ .