


# Pumping Lemma for Context-Free Languages

CSCI 3130 Formal Languages and Automata Theory

Siu On CHAN

Chinese University of Hong Kong

Fall 2017



$$L_1 = \{a^n b^n \mid n \geq 0\}$$

$$L_2 = \{z \mid z \text{ has the same number of a's and b's}\}$$

$$L_3 = \{a^n b^n c^n \mid n \geq 0\}$$

$$L_4 = \{zz^R \mid z \in \{a, b\}^*\}$$

$$L_5 = \{zz \mid z \in \{a, b\}^*\}$$

These languages are not regular

Are they context-free?

## An attempt

$$L_3 = \{a^n b^n c^n \mid n \geq 0\}$$

Let's try to design a CFG or PDA

$$S \rightarrow aBc \mid \varepsilon$$

$$B \rightarrow ???$$

read a / push x

read b / pop x

???

Suppose we could construct some CFG  $G$  for  $L_3$

e.g.

$$S \rightarrow CC \mid BC \mid a$$

$$B \rightarrow CS \mid b$$

$$C \rightarrow SB \mid c$$

How does a long  
derivation look like?

$$S \Rightarrow CC$$

$$\Rightarrow SBC$$

$$\Rightarrow SCSC$$

$$\Rightarrow SSBSC$$

$$\Rightarrow SSBCC$$

$$\Rightarrow aSBCC$$

$$\Rightarrow aaBBCC$$

$$\Rightarrow aabBCC$$

$$\Rightarrow aabbCC$$

$$\Rightarrow aabbcC$$

$$\Rightarrow aabbcc$$

## Repetition in long derivations

If a derivation is long enough, some variable must appear **twice on the same root-to-leave path** in a parse tree

$S \Rightarrow CC$

$\Rightarrow SBC$

$\Rightarrow SCSC$

$\Rightarrow SSBSC$

$\Rightarrow SSBGCC$

$\Rightarrow aSBGCC$


$\Rightarrow aaSBGCC$

$\Rightarrow aabSBGCC$

$\Rightarrow aabbSBGCC$


$\Rightarrow aabbcbGCC$

$\Rightarrow aabbcc$



## Pumping example

Then we can “cut and paste” part of parse tree




## Pumping example

We can repeat this many times


$$\begin{aligned} \text{a}abcc &\Rightarrow \text{aa}abcbcc \Rightarrow \text{aaa}abcbcbcc \Rightarrow \dots \\ &\Rightarrow a(a)^i b(bc)^i c \end{aligned}$$

Every sufficiently large derivation will have a middle part that can be repeated indefinitely


# Pumping in general




$uvwxy$



$uv^2wx^2y$



$uv^3wx^3y$



$uwy$



## Example

$$L_3 = \{a^n b^n c^n \mid n \geq 0\}$$

If  $L_3$  has a context-free grammar  $G$ , then for any sufficiently long  $s \in L(G)$

$s$  can be split into  $s = uvwxy$  such that  $L(G)$  also contains  $uv^2wx^2y$ ,  
 $uv^3wx^3y, \dots$

What happens if  $s = a^m b^m c^m$


**No matter how** it is split,  $uv^2wx^2y \notin L_3$

## Pumping lemma for context-free languages

For every context-free language  $L$

There exists a number  $m$  such that for every long string  $s$  in  $L$  ( $|s| \geq m$ ),  
we can write  $s = uvwxy$  where

1.  $|vwx| \leq m$
2.  $|vx| \geq 1$
3. For every  $i \geq 0$ , the string  $uv^iwx^iy$  is in  $L$



## Pumping lemma for context-free languages

To prove  $L$  is not context-free, it is enough to show that

For every  $m$  there is a long string  $s \in L$ ,  $|s| \geq m$ , such that **for every way** of writing  $s = uvwxy$  where

1.  $|vwx| \leq m$
2.  $|vx| \geq 1$

there is  $i \geq 0$  such that  $uv^iwx^iy$  is not in  $L$

## Using the pumping lemma

$$L_3 = \{a^n b^n c^n \mid n \geq 0\}$$

1. for every  $m$
2. there is  $s = a^m b^m c^m$  (at least  $m$  symbols)
3. no matter how the pumping lemma splits  $s$  into  $uvwx$   
( $|vwx| \leq m, |vx| \geq 1$ )
4.  $uv^2wx^2y \notin L_3$  (but why?)

## Using the pumping lemma

**Case 1:**  $v$  or  $x$  contains two kinds of symbols

aa aabb bbccccc  
           $\underbrace{\hspace{2em}}$   
           $v$

Then  $uv^2wx^2y \notin L_3$  because the pattern is wrong

**Case 2:**  $v$  and  $x$  both contain one kind of symbol

a aaa b bb bbccccc  
   $\underbrace{\hspace{1em}}$    $\underbrace{\hspace{1em}}$   
   $v$            $x$

Then  $uv^2wx^2y$  does not have the same number of a's, b's and c's

Conclusion:  $uv^2wx^2y \notin L_3$

Which is context-free?

$$L_1 = \{a^n b^n \mid n \geq 0\} \quad \checkmark$$

$$L_2 = \{z \mid z \text{ has the same number of a's and b's}\} \quad \checkmark$$

$$L_3 = \{a^n b^n c^n \mid n \geq 0\} \quad \times$$

$$L_4 = \{zz^R \mid z \in \{a, b\}^*\} \quad \checkmark$$

$$L_5 = \{zz \mid z \in \{a, b\}^*\}$$

## Example

$$L_5 = \{zz \mid z \in \{a, b\}^*\}$$

1. for every  $m$
2. there is  $s = a^m b a^m b$  (at least  $m$  symbols)
3. no matter how the pumping lemma splits  $s$  into  $uvwxy$   
( $|vwx| \leq m, |vx| \geq 1$ )
4. Is  $uv^2wx^2y \notin L_5$ ?

## Example

$$L_5 = \{zz \mid z \in \{a, b\}^*\}$$

1. for every  $m$
2. there is  $s = a^m b a^m b$  (at least  $m$  symbols)
3. no matter how the pumping lemma splits  $s$  into  $uvwxy$   
( $|vwx| \leq m, |vx| \geq 1$ )
4. Is  $uv^2wx^2y \notin L_5$ ?

aaa  $\underbrace{a}_v$  aba  $\underbrace{a}_x$  aaab



## Example

$$L_5 = \{zz \mid z \in \{a, b\}^*\}$$

1. for every  $m$
2. there is  $s = a^m b^m a^m b^m$  (at least  $m$  symbols)
3. no matter how the pumping lemma splits  $s$  into  $uvwxy$   
( $|vwx| \leq m, |vx| \geq 1$ )
4. Is  $uv^iwx^iy \notin L_5$  for some  $i$ ?

Recall that  $|vwx| \leq m$

## Example

### Three cases

- Case 1     $aaa \underbrace{aabb}_{vwx} bbaaaaabbbb$   
 $vwx$  is in the **first half** of  $a^m b^m a^m b^m$
- Case 2     $aaaaabb \underbrace{bbba}_{vwx} aaabbbb$   
 $vwx$  is in the **middle part** of  $a^m b^m a^m b^m$
- Case 3     $aaaaabbbbbaaa \underbrace{aabb}_{vwx} bb$   
 $vwx$  is in the **second half** of  $a^m b^m a^m b^m$

## Example

Apply pumping lemma with  $i = 0$

Case 1     $aaa \underbrace{aabb}_{vwx} bbaaaaabbbb$   
 $uwy$  becomes  $a^j b^k a^m b^m$ , where  $j < m$  or  $k < m$

Case 2     $aaaaabb \underbrace{bbba}_{vwx} aaabbbb$   
 $uwy$  becomes  $a^m b^j a^k b^m$ , where  $j < m$  or  $k < m$

Case 3     $aaaaabbbbbbaaa \underbrace{aabb}_{vwx} bb$   
 $uwy$  becomes  $a^m b^m a^j b^k$ , where  $j < m$  or  $k < m$

Not of the form  $zz$

This covers all cases, so  $L_5$  is not context-free