

Notes 13: Local graph partitioning

1. SMALL SPARSE CUT

Given an undirected graph G with positive edge weights, consider the problem of finding a small sparse cut: a vertex set S with small conductance $\varphi(S)$ and has small size:

$$\operatorname{argmin} \{ \varphi(S) \mid S \subseteq V, |S| \leq \delta n \}$$

This is sometimes motivated by finding a small community in a social network.

The spectral partitioning algorithm of Cheeger–Alon–Milman can find a set of small conductance, but the set may be large (containing up to half of the vertices).

We will study an algorithm with the following guarantee: If a graph G has a subset S with small conductance, then the algorithm will find a subset T with $|T| \leq 16|S|$ and $\varphi(T) \leq O(\sqrt{\varphi(S)} \log |S|)$.

Compared with Cheeger–Alon–Milman, we gain in the guarantee that T is small, but we pay an extra $\sqrt{\log |S|}$ factor in conductance.

2. ANALYTIC SPARSITY

For simplicity we consider only d -regular graphs, and further assume d is normalized to be 1. The proof of Cheeger–Alon–Milman inequality shows that given any $x \in \mathbb{R}^V$, we can find a sparse cut $T \subseteq \operatorname{supp}(x) = \{i \in V \mid x_i \neq 0\}$ and $\varphi(T) \leq \sqrt{2R(x)}$, where $R(x) = x^\top \mathcal{L}x / x^\top x$.

If we can solve the problem of minimizing Rayleigh quotient over vector $x \in \mathbb{R}^V$ of small support,

$$\operatorname{argmin} \{ R(x) \mid x \in \mathbb{R}^V, |\operatorname{supp}(x)| \leq \delta n \},$$

then sweep cut algorithm of Cheeger–Alon–Milman outputs a desired subset T from x . But the combinatorial sparsity condition $|\operatorname{supp}(x)| \leq \delta n$ is difficult to work with.

The idea is to relax the combinatorial sparsity condition to the analytic sparsity condition

$$\|x\|_1^2 \leq \delta n \|x\|_2^2.$$

This condition is satisfied whenever $|\operatorname{supp}(x)| \leq \delta n$ (by Cauchy–Schwarz). Also, if x is the probability vector of a distribution μ , then $\|x\|_1^2 = 1$, and

$$\|x\|_2^2 = \sum_{i \in V} \mu(i)^2 = \mathbb{P}_{i \sim \mu, j \sim \mu} [i = j]$$

is the collision probability of μ (the probability for two independent samples from μ to coincide). In particular, if x is the probability vector of the uniform distribution over a subset $S \subseteq V$, then $\|x\|_2^2 = \sum_{i \in S} 1/|S|^2 = 1/|S|$. Therefore the ratio $\|x\|_1^2 / \|x\|_2^2$ is a robust way to measure the size of the support of a distribution.

Turns out any analytically sparse vector with small Rayleigh quotient can be “rounded” into a combinatorially sparse vector with small Rayleigh quotient.

3. ALGORITHM OUTLINE

At a high level, the algorithm is as follows:

- (1) For every vertex i , run lazy random walk from i for t steps for some t depending on $\varphi(S)$
- (2) Truncate t -step lazy walk probability vector π_t into a vector with small support
- (3) Apply Cheeger–Alon–Milman sweep cut to this vector and output a small sparse cut

Why do we expect this algorithm to work? If the random walk starts at a vertex $i \in S$, since $\varphi(S)$ is small, most of the probability mass of $\pi_t^\top = \mathbb{1}_i^\top W^t$ will stay inside S . Here W is the transition matrix of the lazy random walk, and $\mathbb{1}_i$ is the indicator vector for vertex i (the probability vector for the initial distribution of starting the random walk at i). After some time t , the lazy random walk should have become close to the “stationary distribution” in S . Therefore Cheeger–Alon–Milman thresholding should reveal S .

To analyze step (1), we will show that π_t has small Rayleigh quotient, provided the collision probability $\|\pi_t\|_2^2$ is not too small (due to having substantial mass in S).

To analyze step (2), we will show that if there is a small sparse cut S , then π_t will be analytically sparse for some starting vertex $i \in S$. Further, an analytically sparse vector can be truncated to a combinatorially sparse vector with similar Rayleigh quotient.

To analyze step (3), we apply a lemma in proving Cheeger–Alon–Milman inequality.

4. COLLISION PROBABILITY AND RAYLEIGH QUOTIENT

We keep track of how the collision probability $\|\pi_t\|_2^2$ changes over time.

- Initially, $\|\pi_0\|_2^2 = \|\mathbb{1}_i\|_2^2 = 1$.
- $\|\pi_{t+1}\|_2^2 = \|W\pi_t\|_2^2 \leq \|\pi_t\|_2^2$, as W has all eigenvalues bounded by 1 in magnitude. So collision probability $\|\pi_t\|_2^2$ can only decrease over time.
- $\|\pi_t\|_2^2 \rightarrow \|\mathbb{1}/n\|_2^2 = 1/n$ as t grows.

In fact, the ratio $\|\pi_{t+1}\|_2^2/\|\pi_t\|_2^2$ is nondecreasing in t , so $\|\pi_t\|_2^2$ converges to $\|\mathbb{1}/n\|_2^2$ more and more slowly over time. This is proved in the following claim.

Claim 4.1. $\frac{\|\pi_{t+1}\|_2^2}{\|\pi_t\|_2^2} \leq \frac{\|\pi_{t+2}\|_2^2}{\|\pi_{t+1}\|_2^2}$.

Proof. Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of W and v_1, \dots, v_n be its orthonormal eigenvectors. Using the eigen-expansion $\pi_t = \sum_{1 \leq \ell \leq n} c_\ell \lambda_\ell^t v_\ell$ of π_t , we have $\|\pi_t\|_2^2 = \sum_{1 \leq \ell \leq n} c_\ell^2 \lambda_\ell^{2t}$. The desired inequality is $\|\pi_{t+1}\|_2^4 \leq \|\pi_{t+2}\|_2^2 \|\pi_t\|_2^2$, and it becomes

$$\left(\sum_{1 \leq \ell \leq n} c_\ell^2 \lambda_\ell^{2t+2} \right)^2 \leq \left(\sum_{1 \leq \ell \leq n} c_\ell^2 \lambda_\ell^{2t+4} \right) \left(\sum_{1 \leq \ell \leq n} c_\ell^2 \lambda_\ell^{2t} \right),$$

which follows by Cauchy–Schwarz. \square

What happens when $\|\pi_t\|_2^2$ decreases slowly? $\|\pi_{t+1}\|_2^2/\|\pi_t\|_2^2$ will be close to 1, or equivalently $1 - (\|\pi_{t+1}\|_2^2/\|\pi_t\|_2^2)$ is close to 0. We can express

$$1 - \frac{\|\pi_{t+1}\|_2^2}{\|\pi_t\|_2^2} = 1 - \frac{\|W\pi_t\|_2^2}{\|\pi_t\|_2^2} = \frac{\pi_t^\top (I - W^\top W) \pi_t}{\pi_t^\top \pi_t} = \frac{\pi_t^\top \mathcal{L}' \pi_t}{\pi_t^\top \pi_t}$$

as the Rayleigh quotient $R_{\mathcal{L}'}(\pi_t)$ for the matrix $\mathcal{L}' = I - W^2$. Turns out \mathcal{L}' is the normalized Laplacian of some graph H ! This graph H is the two-step lazy random walk, where every step in H corresponds to two consecutive steps in W . More precisely, H also has vertex set V , and every edge (i, k) in H corresponds to a length-2 path $(i, j), (j, k)$ in the lazy random walk W . The weight w_{ik} of (i, k) in H is $w_{ij}w_{jk}$, the product of weights of the two edges in the path in W . H has normalized adjacency matrix W^2 . We won't prove these claims about H since our proof does not depend on them, and will leave them as easy exercises.

This means when $\|\pi_t\|_2^2$ decreases slowly at time t , the probability vector π_t corresponds two small Rayleigh quotient (hence a sparse cut, by Cheeger–Alon–Milman) in the two-step lazy random walk graph H .

We can translate small Rayleigh quotient $R_{\mathcal{L}'}(\pi_t)$ (for the two-step walk) into small Rayleigh quotient $R(\pi_t)$ (for the original lazy walk) using the following claim:

Claim 4.2. For any $x \in \mathbb{R}^V$ and lazy random walk transition W , $x^\top W^2 x \leq x^\top W x$. Therefore

$$R_{\mathcal{L}'}(x) = \frac{x^\top (I - W^2) x}{x^\top x} \geq \frac{x^\top (I - W) x}{x^\top x} = R(x).$$

Proof. $W = I^{-1}W$ coincides with the normalized adjacency matrix of G , since G is assumed to be 1-regular, so the degree matrix is I .

Since W is lazy, $W = \frac{1}{2}I + \frac{1}{2}W'$, where W' is the transition/normalized adjacency matrix of the non-lazy random walk on G . Then $W - W^2 = \frac{1}{4}I - \frac{1}{4}(W')^2 = \frac{1}{4}\mathcal{L}_{W'} \succeq 0$. \square

The above claim is the only place we require the random walk to be lazy.

We get the following upperbound on Rayleigh quotient $R(\pi_{t-1})$ if we can lower bound the collision probability $\|\pi_t\|_2^2$.

Proposition 4.3. $R(\pi_{t-1}) \leq 1 - \|\pi_t\|_2^{2/t}$.

Proof. Since $\|\pi_0\|_2^2 = 1$,

$$\|\pi_t\|_2^2 = \frac{\|\pi_t\|_2^2}{\|\pi_0\|_2^2} = \frac{\|\pi_t\|_2^2}{\|\pi_{t-1}\|_2^2} \frac{\|\pi_{t-1}\|_2^2}{\|\pi_{t-2}\|_2^2} \dots \frac{\|\pi_1\|_2^2}{\|\pi_0\|_2^2} \leq \left(\frac{\|\pi_t\|_2^2}{\|\pi_{t-1}\|_2^2} \right)^t,$$

where the inequality is [Claim 4.1](#). This inequality and [Claim 4.2](#) implies

$$R(\pi_{t-1}) \leq R_{\mathcal{L}'}(\pi_{t-1}) = 1 - \frac{\pi_{t-1}^\top W^\top W \pi_{t-1}}{\pi_{t-1}^\top \pi_{t-1}} = 1 - \frac{\|\pi_t\|_2^2}{\|\pi_{t-1}\|_2^2} \leq 1 - \|\pi_t\|_2^{2/t}. \quad \square$$

5. TRUNCATING ANALYTICALLY SPARSE VECTOR

Lemma 5.1. *Suppose $x \in \mathbb{R}_{\geq 0}^V$ satisfies $\|x\|_1^2 \leq s\|x\|_2^2$. Then it can be truncated into a vector $y \in \mathbb{R}_{\geq 0}^V$ with $|\text{supp}(y)| \leq 4s$ and $R(y) \leq 2R(x)$.*

Proof. By scaling, assume $\|x\|_2^2 = s$ and $\|x\|_1 \leq s$.

Let $y \in \mathbb{R}_{\geq 0}^V$ be the vector $y_i = \max\{x_i - 1/4, 0\}$.

Then $s \geq \|x\|_1 \geq \sum_{i \in \text{supp}(y)} x_i \geq |\text{supp}(y)| \frac{1}{4}$, because every $i \in \text{supp}(y)$ contributes $x_i \geq 1/4$ to $\|x\|_1$. Hence $|\text{supp}(y)| \leq 4s$.

We will compare $R(y)$ and $R(x)$, where $R(x) = \frac{x^\top \mathcal{L}x}{x^\top x} = \frac{\sum_{(i,j) \in E} (x_i - x_j)^2}{d \sum_{i \in V} x_i^2}$.

For the numerator, $(y_i - y_j)^2 \leq (x_i - x_j)^2$ because truncation can only reduce the difference. Hence $y^\top \mathcal{L}y \leq x^\top \mathcal{L}x$.

For the denominator, we have $y_i^2 \geq x_i^2 - \frac{1}{2}x_i$, so

$$\sum_{i \in V} y_i^2 \geq \sum_{i \in V} x_i^2 - \frac{1}{2} \sum_{i \in V} x_i \geq s - \frac{1}{2}s = \frac{s}{2} = \frac{1}{2} \sum_{i \in V} x_i^2.$$

Hence $y^\top y \geq x^\top x/2$.

Therefore $R(y) = y^\top \mathcal{L}y / y^\top y \leq x^\top \mathcal{L}x / (x^\top x/2) = 2R(x)$. \square

6. ANALYTICALLY SPARSE VECTOR FROM SMALL SPARSE CUT

Given a probability π over V , we write $\pi(S) = \sum_{i \in S} \pi(i)$ to denote its total probability in $S \subseteq V$.

Claim 6.1. *If initial distribution $\mu_0 = \mathbb{1}_S/|S|$ is uniform over subset S , and $\mu_t = W^t \mu_0$, then $\mu_t(S) \geq 1 - t\varphi(S)$.*

Proof. We lowerbound $\mu_t(S)$ by the probability the random walk stays inside S for all t steps. We will upperbound the probability it leaves S in any of the t steps.

Every vertex i in the initial distribution μ_0 carries $\mu_0(i) = 1/|S|$ probability. Since the graph is d -regular, an edge going out of S carries $\frac{w_{ij}}{d|S|}$ probability out of S . Total probability escaping out of S in the first step is $\sum_{i \in S, j \in \bar{S}} \frac{w_{ij}}{d|S|} = \varphi(S)$.

We can finish the proof if the escape probability for every step is at most $\varphi(S)$. This is true by repeating the above calculations (changing “=” to “ \leq ”), and observing every vertex i at any time t carries probability $\mu_t(i)$ at most $1/|S|$.

Why is $\mu_i(t) \leq 1/|S|$ for any i and any t ? This is true for initially $t = 0$ for all vertices i . For future time steps, $\mu_i(t+1)$ is a weighted average of $\mu_j(t)$ over neighbors j of i , so it remains true for time $t+1$. \square

Corollary 6.2. *There is a starting point $i \in S$ such that if $\pi_0^{(i)} = \mathbb{1}_i$ and $\pi_t^{(i)} = W^t \pi_0^{(i)}$, then $\pi_t^{(i)}(S) \geq 1 - t\varphi(S)$.*

Proof. The uniform distribution μ_0 over S is the average, over a uniformly random $i \in S$, of initial distributions $\mathbb{1}_i$ starting from a single vertex i in S , because $\mu_0 = \frac{\mathbb{1}_S}{|S|} = \mathbb{E}_{i \sim \mu_0} [\mathbb{1}_i]$.

Now $\mu_t(S)$ is the same averaging of $\pi_t^{(i)}(S)$, because

$$\mu_t(S) = (W^t \mu_0)(S) = \left(W^t \mathbb{E}_{i \sim \mu_0} [\mathbb{1}_i] \right) (S) = \mathbb{E}_{i \sim \mu_0} [W^t \mathbb{1}_i](S) = \mathbb{E}_{i \sim \mu_0} [\pi_t^{(i)}(S)].$$

The key observation here is that the t -step lazy random walk W^t is a linear operator, so taking average first and then t -step walk is the same as taking t -step walk first

Some vertex i in S must achieve staying probability $\pi_t^{(i)}(S)$ at least the average $\mu_t(S)$. \square

Lemma 6.3. *For any distribution π , its collision probability $\|\pi\|_2^2 \geq \pi(S)^2/|S|$.*

Proof. Expand $\|\pi\|_2^2$ and apply Cauchy–Schwarz,

$$\|\pi\|_2^2 \geq \sum_{j \in S} \pi(j)^2 \geq \frac{1}{|S|} \left(\sum_{j \in S} \pi(j) \right)^2 = \frac{1}{|S|} \pi(S)^2.$$

This Cauchy–Schwarz inequality implies that the distribution over S with the smallest collision probability is the uniform distribution, and has collision probability $1/|S|$. \square

7. ALGORITHM

We know the graph contains a small subset S with conductance $\varphi(S)$. **Corollary 6.2** implies that if we are lucky to choose $i \in S$ as the starting point of our random walk, then even after $t + 1 = 1/2\varphi(S)$ steps, there is still $\pi_{t+1}(S) \geq 1/2$ probability mass of staying in S .

Lemma 6.3 then implies the collision probability $\|\pi_{t+1}\|_2^2 \geq 1/4|S|$.

Proposition 4.3 gives the following upperbound on Rayleigh quotient:

$$R(\pi_t) \leq 1 - \|\pi_{t+1}\|_2^{2/(t+1)} \leq 1 - \frac{1}{(4|S|)^{2\varphi(S)}} = 1 - \exp(-2\varphi(S) \ln(4|S|)) = O(\varphi(S) \ln |S|),$$

where the last equality is due to $1 - e^{-x} = O(x)$ for small x near 0.

π_t is analytically sparse and has sparsity ratio $\|\pi_t\|_1^2/\|\pi_t\|_2^2 = 1/\|\pi_t\|_2^2 \leq 1/\|\pi_{t+1}\|_2^2 \leq 4|S|$.

Lemma 5.1 truncates π_t to some nonnegative vector y with $|\text{supp}(y)| \leq 16|S|$ and $R(y) = O(\varphi(S) \ln |S|)$.

Cheeger–Alon–Milman outputs a super-level set $T = \{i \in V \mid y_i > r\}$ of y with $|T| \leq 16|S|$ and $\varphi(T) \leq \sqrt{2R(y)} = O(\sqrt{\varphi(S) \ln |S|})$.

8. SMALL-SET EXPANSION

The above conductance guarantee has an extra $\sqrt{\log |S|}$ factor. Is there an efficient approximation algorithm whose approximation factor is independent of the size of S ?

Such an algorithm, if exists, will solve the Small-Set-Expansion problem, defined as follows:

Small-Set-Expansion

Parameters: conductance bound ε and size bound δ
Input: regular undirected graph G
Goal: decide between the following two cases:
 (Yes) Some $S \subseteq V$ with $|S| \leq \delta n$ satisfies $\varphi(S) \leq \varepsilon$
 (No) All $S \subseteq V$ with $|S| \leq 16\delta n$ satisfies $\varphi(S) \geq 1 - \varepsilon$

You may think of the problem as asking if a graph has a hidden small “community” (subset with small conductance). And it only asks for deciding between two extreme cases of conductance: either some small subset has conductance very close to 0, or all small subsets have conductance very close to 1.

A conjecture known as Small-Set-Expansion Hypothesis says that Small-Set-Expansion is hard to solve.

Conjecture 8.1 (Raghavendra and Steurer 2010). *For every $\varepsilon > 0$, there is $\delta > 0$ such that Small-Set-Expansion with parameters ε and δ is NP-hard.*

In particular, if Small-Set-Expansion Hypothesis holds and $P \neq NP$, then no efficient algorithm can avoid the dependence on $|S|$.

Small-Set-Expansion Hypothesis also implies the Unique-Games Conjecture, a central open problem in approximation algorithms that we will not define here. The latter conjecture says that certain constraint satisfaction problem called Unique-Games is NP-hard to approximate.

If Unique-Games Conjecture holds and $P \neq NP$, then a simple SDP algorithm will be the best approximation algorithm for many problems. A consequence is that Goemans–Williamson rounding algorithm for MaxCut (with approximation factor 0.878...) will be optimal.