

NOTE ON MATH2060B: ELEMENTARY ANALYSIS II (2020-21)

CHI-WAI LEUNG

1. DIFFERENTIATION

Throughout this section, let I be an open interval (not necessarily bounded) and let f be a real-valued function defined on I .

Definition 1.1. Let $c \in I$. We say that f is differentiable at c if the following limit exists:

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

In this case, we write $f'(c)$ for the above limit and we call it the derivative of f at c . We say that if f is differentiable on I if $f'(x)$ exists for every point x in I .

Proposition 1.2. Let $c \in I$. Then $f'(c)$ exists if and only if there is a function φ defined on I such that the function φ is continuous at c and

$$f(x) - f(c) = \varphi(x)(x - c)$$

for all $x \in I$.

In this case, $\varphi(c) = f'(c)$.

Proof. Assume that $f'(c)$ exists. Define a function $\varphi : I \rightarrow \mathbb{R}$ by

$$\varphi(x) = \begin{cases} \frac{f(x) - f(c)}{x - c} & \text{if } x \neq c; \\ f'(c) & \text{if } x = c. \end{cases}$$

Clearly, we have $f(x) - f(c) = \varphi(x)(x - c)$ for all $x \in I$. We want to show that the function φ is continuous at c . In fact, let $\varepsilon > 0$, by the definition of the limit of a function, there is $\delta > 0$ such that

$$\left| f'(c) - \frac{f(x) - f(c)}{x - c} \right| < \varepsilon$$

whenever $x \in I$ with $0 < |x - c| < \delta$. Therefore, we have $|f'(c) - \varphi(x)| < \varepsilon$ as $x \in I$ with $0 < |x - c| < \delta$. Since $\varphi(c) = f'(c)$, we have $|f'(c) - \varphi(x)| < \varepsilon$ as $x \in I$ with $|x - c| < \delta$, hence the function φ is continuous at c as desired.

The converse is clear since $\varphi(x) = \frac{f(x) - f(c)}{x - c}$ if $x \neq c$. The proof is complete. \square

Proposition 1.3. Using the notation as above, if f is differentiable at c , then f is continuous at c .

Proof. By using Proposition 1.2, if $f'(c)$ exists, then there is a function φ defined on I such that the function φ is continuous at c and we have $f(x) - f(c) = \varphi(x)(x - c)$ for all $x \in I$. This implies that $\lim_{x \rightarrow c} f(x) = f(c)$, so f is continuous at c as desired. \square

Remark 1.4. In general, the converse of Proposition 1.3 does not hold, for example, the function $f(x) := |x|$ is a continuous function on \mathbb{R} but $f'(0)$ does not exist.

Proposition 1.5. *Let f and g be the functions defined on I . Assume that f and g both are differentiable at $c \in I$. We have the following assertions.*

- (i) $(f + g)'(c)$ exists and $(f + g)'(c) = f'(c) + g'(c)$.
- (ii) The product $(f \cdot g)'(c)$ exists and $(f \cdot g)'(c) = f'(c)g(c) + f(c)g'(c)$.
- (iii) If $g(c) \neq 0$, then we have $(\frac{f}{g})'(c)$ exists and $(\frac{f}{g})'(c) = \frac{f'(c)g(c) - f(c)g'(c)}{g(c)^2}$.

Proof. Part (i) clearly follows from the definition of the limit of a function.

For showing Part (ii), note that we have

$$\frac{f(x)g(x) - f(c)g(c)}{x - c} = \frac{f(x) - f(c)}{x - c}g(x) + f(c)\frac{g(x) - g(c)}{x - c}$$

for all $x \in I$ with $x \neq c$. From this, together with Proposition 1.3, Part (ii) follows.

For Part (iii), by using Part (ii), it suffices to show that $(\frac{1}{g})'(c) = -\frac{g'(c)}{g(c)^2}$. In fact, $g'(c)$ exists, so g is continuous at c . Since $g(c) \neq 0$, there is $\delta_1 > 0$ so that $g(x) \neq 0$ for all $x \in I$ with $|x - c| < \delta_1$. Then we have

$$\frac{1}{x - c} \left(\frac{1}{g(x)} - \frac{1}{g(c)} \right) = \frac{1}{x - c} \left(\frac{g(c) - g(x)}{g(x)g(c)} \right)$$

for all $x \in I$ with $0 < |x - c| < \delta_1$. By taking $x \rightarrow c$, we see that $(\frac{1}{g})'(c)$ exists and $(\frac{1}{g})'(c) = \frac{-g'(c)}{g(c)^2}$. The proof is complete. \square

Proposition 1.6. (Chain Rule): *Let f, g be functions defined on \mathbb{R} . Let $d = f(c)$ for some $c \in \mathbb{R}$. Suppose that $f'(c)$ and $g'(d)$ exist. Then the derivative of composition $(g \circ f)'(c)$ exists and $(g \circ f)'(c) = g'(d)f'(c)$.*

Proof. By using Proposition 1.2, we want to find a function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$g \circ f(x) - g \circ f(c) = \varphi(x)(x - c)$$

for all $x \in \mathbb{R}$ and the function $\varphi(x)$ is continuous at c , and so $(g \circ f)'(c) = \varphi(c)$.

Let $y = f(x)$. By using Proposition 1.2 again, there is a function $\beta(y)$ so that $g(y) - g(d) = \beta(y)(y - d)$ for all $y \in \mathbb{R}$ and $\beta(y)$ is continuous at d . Similarly, there is a function $\alpha(x)$ we have $f(x) - f(c) = \alpha(x)(x - c)$ for all $x \in \mathbb{R}$ and $\alpha(x)$ is continuous at c . These two equations imply that

$$g \circ f(x) - g \circ f(c) = \beta(f(x))(f(x) - f(c)) = \beta(f(x))\alpha(x)(x - c)$$

for all $x \in \mathbb{R}$. Let $\varphi(x) := \beta(f(x)) \cdot \alpha(x)$ for $x \in \mathbb{R}$. Since $\beta(d) = g'(d)$ and $\alpha(c) = f'(c)$, we see that $\varphi(c) = \beta(f(c))\alpha(c) = g'(d)f'(c)$. It remains to show that the function φ is continuous at c . In fact, $f'(c)$ exists, so f is continuous at c , and hence the composition $\beta \circ f(x)$ is continuous at c . In addition, the function α is continuous at c . Therefore, the function $\varphi := (\beta \circ f) \cdot \alpha$ is continuous at c , and so $(g \circ f)'(c)$ exists with $(g \circ f)'(c) = \varphi(c) = g'(d)f'(c)$. The proof is complete. \square

Proposition 1.7. *Let I and J be open intervals. Let f be a strictly increasing function from I onto J . Let $d = f(c)$ for $c \in I$. Assume that $f'(c)$ exists and the inverse of f , write $g := f^{-1}$, is continuous at d . If $f'(c) \neq 0$, then $g'(d)$ exists and $g'(d) = \frac{1}{f'(c)}$.*

Proof. Let $y = f(x)$. Note that by using Proposition 1.2, there is a function F on I such that $f(x) - f(c) = F(x)(x - c)$ for all $x \in I$ and F is continuous at c with $F(c) = f'(c) \neq 0$. F is continuous at c , so there are open intervals I_1 and J_1 such that $c \in I_1 \subseteq I$ and $d \in f(I_1) = J_1$, moreover, $F(x) \neq 0$ for all $x \in I_1$. Note that since $f(x) - f(c) = F(x)(x - c)$, we have $y - d = f(g(y)) - f(g(c)) = F(g(y))(g(y) - g(d))$ for all $y \in J_1$. Since $F(x) \neq 0$ for all $x \in I_1$, we have $g(y) - g(d) = F(g(y))^{-1}(y - d)$ for all $y \in J_1$. Note that the function $F(g(y))^{-1}$ is continuous at d . Thus, $g'(d)$ exists and $g'(d) = F(g(d))^{-1} = \frac{1}{f'(c)}$ as desired. \square

Definition 1.8. Let D be a non-empty subset of \mathbb{R} and let g be a real-valued function defined on D .

(i) We say that g has an absolute maximum (resp. absolute minimum) at a point $c \in D$ if $g(c) \geq g(x)$ (resp. $g(c) \leq g(x)$) for all $x \in D$.

In this case, c is called an absolute extreme point of g .

(ii) We say that g has a local maximum (resp. local minimum) at a point $c \in D$ if there is $r > 0$ such that $(c - r, c + r) \subseteq D$ and $g(c) \geq g(x)$ (resp. $g(c) \leq g(x)$) for all $x \in (c - r, c + r)$.

In this case, c is called a local extreme point of g .

Remark 1.9. Note that an absolute extreme point of a function g need not be a local extreme point, for example if $g(x) := x$ for $x \in [0, 1]$, then g has an absolute maximum point at $x = 1$ of g but 1 is not a local maximum point of g .

Proposition 1.10. Let I be an open interval and let f be a function on I . Assume that f has a local extreme point at $c \in I$ and $f'(c)$ exists. Then $f'(c) = 0$.

Proof. Without lost the generality, we may assume that f has local minimum at c . Then there is $r > 0$ such that $f(x) \geq f(c)$ for $x \in (c - r, c + r) \subseteq I$. Since $f'(c)$ exists, by using Proposition 1.2, there is a function φ defined on I such that $f(x) - f(c) = \varphi(x)(x - c)$ for all $x \in I$ and φ is continuous at c with $\varphi(c) = f'(c)$. Thus, we have $\varphi(x)(x - c) \geq 0$ for all $x \in (c - r, c + r)$. From this we see that $\varphi(x) \geq 0$ as $x \in (c, c + r)$, similarly, $\varphi(x) \leq 0$ as $x \in (c - r, c)$. The function φ is continuous at c , so $\varphi(c) = 0$ and hence $f'(c) = \varphi(c) = 0$ as desired. \square

Proposition 1.11. Rolle's Theorem: Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function. Assume that $f'(x)$ exists for all $x \in (a, b)$ and $f(a) = f(b)$. Then there is a point $c \in (a, b)$ such that $f'(c) = 0$.

Proof. Recall a fact that every continuous function defined a compact attains absolute points, that is, there are c_1 and c_2 such that $f(c_1) = \min_{x \in [a, b]} f(x)$ and $f(c_2) = \max_{x \in [a, b]} f(x)$, hence, $f(c_1) \leq f(x) \leq f(c_2)$ for all $x \in [a, b]$. If $f(c_1) = f(c_2)$, then $f(x) \equiv f(c_1) = f(c_2)$ for all $x \in [a, b]$, so $f'(x) \equiv 0$ for all $x \in (a, b)$.

Otherwise, suppose that $f(c_1) < f(c_2)$. Since $f(a) = f(b)$, we have $c_1 \in (a, b)$ or $c_2 \in (a, b)$. We may assume that $c_1 \in (a, b)$. Then $x = c_1$ is a local minimum point of f . Therefore, $f'(c_1) = 0$ by using Proposition 1.10. \square

Theorem 1.12. Main Value Theorem: If $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function and is differentiable on (a, b) , then there is a point $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Proof. Define a function $\varphi : [a, b] \rightarrow \mathbb{R}$ by

$$\varphi(x) = f(x) - f(a) - \frac{f(b) - f(a)}{b - a}(x - a)$$

for $x \in [a, b]$. Note that the function φ is continuous on $[a, b]$ with $\varphi(a) = \varphi(b) = 0$, in addition, $\varphi'(x)$ exists for all $x \in (a, b)$. The Rolle's Theorem implies that there is a point $c \in (a, b)$ such that

$$0 = \varphi'(c) = f'(c) - \frac{f(b) - f(a)}{b - a}.$$

The proof is complete. \square

Corollary 1.13. Assume that $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function and is differentiable on (a, b) . If $f' \equiv 0$ on (a, b) , then f is a constant function.

Proof. Fix any point $z \in (a, b)$. Let $x \in (z, b]$. By using the Mean Value Theorem, there is a point $c \in (z, x)$ such that $f(x) - f(z) = f'(c)(x - z)$. If $f' \equiv 0$ on (a, b) , so $f(x) = f(z)$ for all $x \in [z, b]$. Similarly, we have $f(x) = f(z)$ for all $x \in [a, z]$. The proof is complete. \square

Definition 1.14. We call a function f is a C^1 -function on I if $f'(x)$ exists and continuous on I . In addition, we define the n -derivatives of f by $f^{(n)}(x) := f^{(n-1)}(x)$ for $n \geq 2$, provided it exists. In this case, we say that f is a C^n -function on I . In particular, we call f a C^∞ -function (or smooth function) if f is a C^n -function for all $n = 1, 2, \dots$

For example, the exponential function $\exp x$ is a very important example of smooth function on \mathbb{R} .

Corollary 1.15. Inverse Mapping Theorem: Let f be a C^1 -function on an open interval I and let $c \in I$. Assume that $f'(c) \neq 0$. Then there is $r > 0$ such that the function f is a strictly monotone function on $(c - r, c + r) \subseteq I$. If we let $J := f(c - r, c + r)$, then the inverse function $g := f^{-1} : J \rightarrow (c - r, c + r)$ is also a C^1 -function.

Proof. We may assume that $f'(c) > 0$. $f'(x)$ is continuous on I , so there is $r > 0$ such that $f'(x) > 0$ for all $x \in (c - r, c + r) \subseteq I$. For any x_1 and x_2 in $(c - r, c + r)$ with $x_1 < x_2$, by using the Mean Value Theorem, we have $f(x_2) - f(x_1) = f'(v)(x_2 - x_1)$ for some $v \in (x_1, x_2)$, and hence $f(x_2) > f(x_1)$. Therefore the restriction of f on $(c - r, c + r)$ is a strictly increasing function, thus, it is an injection. Let $J := f((c - r, c + r))$. Then J is an interval by the Intermediate Value Theorem. Moreover, J is an open interval because f is strictly increasing. Also, if we let $g = f^{-1}$ on J , then g is continuous on J due to the fact that every continuous bijection on a compact set is a homeomorphism. Therefore, by Proposition 1.7, we see that $g'(y)$ exists on J and $g'(y) = \frac{1}{f'(x)}$ for $y = f(x)$ and $x \in (c - r, c + r)$. Therefore, g is a C^1 function on J . The proof is complete. \square

Proposition 1.16. Cauchy Mean Value Theorem: Let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous functions with $g(a) \neq g(b)$. Assume that f, g are differentiable functions on (a, b) and $g'(x) \neq 0$ for all $x \in (a, b)$. Then there is a point $c \in (a, b)$ such that $\frac{f(b)-f(a)}{g(b)-g(a)} = \frac{f'(c)}{g'(c)}$.

Proof. Define a function ψ on $[a, b]$ by $\psi(x) = f(x) - f(a) - \frac{f(b)-f(a)}{g(b)-g(a)}(g(x) - g(a))$ for $x \in [a, b]$. Then by using the similar argument as in the Mean Value Theorem, the result follows. \square

Theorem 1.17. Lagrange Remainder Theorem: Let f be a $C^{(n+1)}$ function defined on (a, b) . Let $x_0 \in (a, b)$. Then for each $x \in (a, b)$, there is a point c between x_0 and x such that

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{f^{(n+1)}(c)}{(n+1)!} (x - x_0)^{n+1}.$$

Proof. We may assume that $x_0 < x < b$. **Case:** We first assume that $f^{(k)}(x_0) = 0$ for all $k = 0, 1, \dots, n$. Put $g(t) = (t - x_0)^{n+1}$ for $t \in [x_0, x]$. Then $g'(t) = (n+1)(t - x_0)^n$ and $g(x_0) = 0$. Then by the Cauchy Mean Value Theorem, there is $x_1 \in (x_0, x)$ such that $\frac{f(x)}{g(x)} = \frac{f(x)-f(x_0)}{g(x)-g(x_0)} = \frac{f'(x_1)}{g'(x_1)}$. Using the same step for f' and g' on $[x_0, x_1]$, there is $x_2 \in (x_0, x_1)$ such that $\frac{f'(x_1)}{g'(x_1)} = \frac{f'(x_1)-f'(x_0)}{g'(x_1)-g'(x_0)} = \frac{f^{(2)}(x_2)}{g^{(2)}(x_2)}$. To repeat the same step, there are x_1, x_2, \dots, x_{n+1} in (a, b) such that $x_k \in (x_0, x_{k-1})$ for $k = 1, 2, \dots, n+1$ and

$$\frac{f(x)}{g(x)} = \frac{f'(x_1)}{g'(x_1)} = \dots = \frac{f^{(n+1)}(x_{n+1})}{g^{(n+1)}(x_{n+1})}.$$

In addition, note that $g^{n+1}(x_{n+1}) = (n+1)!$. Therefore, we have $\frac{f(x)}{g(x)} = \frac{f^{(n+1)}(x_{n+1})}{(n+1)!}$, and hence $f(x) = \frac{f^{(n+1)}(x_{n+1})}{(n+1)!} (x - x_0)^{n+1}$. Note $x_{n+1} \in (x_0, x)$ and thus, the result holds for this case.

For the general case, put $G(x) = f(x) - \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$ for $x \in (a, b)$. Note that we have $G(x_0) = G'(x_0) = \dots = G^{(n)}(x_0) = 0$. Then by the Claim above, there is a point $c \in (x_0, x)$ such that $G(x) = \frac{G^{(n+1)}(c)}{(n+1)!}$. Since $G^{(n+1)}(c) = f^{(n+1)}(c)$, $f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{f^{(n+1)}(c)}{(n+1)!}$. The proof is complete. \square

Example 1.18. Recall that the exponential function e^x is defined by

$$e^x := \sum_{k=0}^{\infty} \frac{x^k}{k!} := \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{x^k}{k!}$$

for $x \in \mathbb{R}$. Note that the above limit always exists for all $x \in \mathbb{R}$ (shown in the last chapter).

Show that the natural base e is an irrational number.

Put $f(x) := e^x$ for $x \in \mathbb{R}$. It is a known fact f is a C^∞ function and $f^{(n)}(x) = e^x$ for all $x \in \mathbb{R}$. Fix any $x > 0$. Then by the Lagrange Theorem, for each positive integer n , there is $c_n \in (0, x)$ such that

$$f(x) = \sum_{k=0}^n \frac{x^k}{k!} + \frac{e^{c_n}}{(n+1)!} x^{n+1}.$$

In particular, taking $x = 1$, we have

$$0 < \frac{e^{c_n}}{(n+1)!} = e - \sum_{k=0}^n \frac{1}{k!} < \frac{3}{(n+1)!}$$

for all positive integer n . Now if $e = p/q$ for some positive integers p and q , and thus, we have

$$0 < \frac{p}{q} - \sum_{k=0}^n \frac{1}{k!} < \frac{3}{(n+1)!}$$

for all $n = 1, 2, \dots$. Now we can choose n large enough such that $(n!)^2 \in \mathbb{N}$. It leads to a contradiction because we have

$$0 < (n!)^{\frac{p}{q}} - (n!) \sum_{k=0}^n \frac{1}{k!} < \frac{3(n!)}{(n+1)!} = \frac{3}{n+1} < 1.$$

Therefore, e is irrational.

Proposition 1.19. Let f be a C^2 function on an open interval I and $x_0 \in I$. Assume that $f'(x_0) = 0$. Then f has local maximum (resp. local minimum) at x_0 if $f^{(2)}(x_0) < 0$ (resp. $f^{(2)}(x_0) > 0$).

Proof. We assume that $f^{(2)}(x_0) > 0$. We want to show that x_0 is a local minimum point of f . The proof of another case is similar. Note that for any $x \in I \setminus \{x_0\}$. Then by the Lagrange Theorem, there is a point c between x_0 and x such that

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{1}{2} f^{(2)}(x_0)(x - x_0)^2 = f(x_0) + \frac{1}{2} f^{(2)}(x_0)(x - x_0)^2.$$

$f^{(2)}$ is continuous at x_0 and $f^{(2)}(x_0) > 0$, and so there is $r > 0$ such that $f^{(2)}(x) > 0$ for all $x \in (x_0 - r, x_0 + r) \subseteq I$. Therefore, we have

$$f(x) = f(x_0) + \frac{1}{2} f^{(2)}(x)(x - x_0)^2 \geq f(x_0)$$

for all $x \in (x_0 - r, x_0 + r)$ and thus, x_0 is a local minimum point of f as desired. \square

Proposition 1.20. L'Hospital's Rule: Let f and g be the differentiable functions on (a, b) and let $c \in (a, b)$. Assume that $f(c) = g(c) = 0$, in addition, $g'(x) \neq 0$ and $g(x) \neq 0$ for all $x \in (a, b) \setminus \{c\}$. If the limit $L := \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$ exists, then so does $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$, moreover, we have $L = \lim_{x \rightarrow c} \frac{f(x)}{g(x)}$.

Proof. Fix $c < x < b$. Then by the Cauchy Mean Value Theorem, there is a point $x_1 \in (c, x)$ such that

$$\frac{f(x)}{g(x)} = \frac{f(x) - f(c)}{g(x) - g(c)} = \frac{f'(x_1)}{g'(x_1)}$$

$x_1 \in (c, x)$, so if $L := \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$ exists, then $\lim_{x \rightarrow c^+} \frac{f(x)}{g(x)}$ exists and is equal to L .

Similarly, we also have $\lim_{x \rightarrow c^-} \frac{f(x)}{g(x)} = L$. The proof is finished. \square

Proposition 1.21. Let f be a function on (a, b) and let $c \in (a, b)$.

(i) If $f'(c)$ exists, then the following limit exists (also called the symmetric derivatives of f at c):

$$f'(c) = \lim_{t \rightarrow 0} \frac{f(c+t) - f(c-t)}{2t}.$$

(ii) If $f^{(2)}(c)$ exists, then

$$f^{(2)}(c) = \lim_{t \rightarrow 0} \frac{f(c+t) - 2f(c) + f(c-t)}{t^2}.$$

Proof. For showing (i), note that we have

$$f'(c) = \lim_{t \rightarrow 0^+} \frac{f(c+t) - f(c)}{t} = \lim_{t \rightarrow 0^-} \frac{f(c+t) - f(c)}{t}.$$

Putting $t = -s$ into the second equality above, we see that

$$f'(c) = \lim_{s \rightarrow 0^+} \frac{f(c-s) - f(c)}{-s}.$$

To sum up the two equations above, we have

$$f'(c) = \lim_{t \rightarrow 0^+} \frac{f(c+t) - f(c-t)}{2t}.$$

Similarly, we have $f'(c) = \lim_{t \rightarrow 0^-} \frac{f(c+t) - f(c-t)}{2t}$. Part (i) follows.

For showing Part (ii), let $h(t) := f(c+t) - 2f(c) + f(c-t)$ for $t \in \mathbb{R}$. Then $h(0) = 0$ and $h'(t) = f'(c+t) - f'(c-t)$. By using the L'Hospital's Rule and Part (i), we have

$$\lim_{t \rightarrow 0} \frac{f(c+t) - 2f(c) + f(c-t)}{t^2} = \lim_{t \rightarrow 0} \frac{h'(t)}{(t^2)'} = \lim_{t \rightarrow 0} \frac{f'(c+t) - f'(c-t)}{2t} = f^{(2)}(c).$$

The proof is complete. \square

Definition 1.22. A function f defined on (a, b) is said to be convex if for any pair $a < x_1 < x_2 < b$, we have

$$f((1-t)x_1 + tx_2) \leq (1-t)f(x_1) + tf(x_2)$$

for all $t \in [0, 1]$.

Proposition 1.23. Let f be a C^2 function on (a, b) . Then f is a convex function if and only if $f^{(2)}(x) \geq 0$ for all $x \in (a, b)$.

Proof. For showing (\Rightarrow): assume that f is a convex function. Fix a point $c \in (a, b)$. f is convex, so we have $f(c) = f(\frac{1}{2}(c+t) + \frac{1}{2}(c-t)) \leq \frac{1}{2}f(c+t) + \frac{1}{2}f(c-t)$ for all $t \in \mathbb{R}$ with $c \pm t \in (a, b)$. By Proposition 1.21, we have

$$f^{(2)}(c) = \lim_{t \rightarrow 0} \frac{f(c+t) - 2f(c) + f(c-t)}{t^2}.$$

Therefore, we have $f^{(2)}(c) \geq 0$.

For (\Leftarrow), assume that $f^{(2)}(x) \geq 0$ for all $x \in (a, b)$. Fix $a < x_1 < x_2 < b$ and $t \in [0, 1]$. Let $c := (1-t)x_1 + tx_2$. Then by the Lagrange Remainder Theorem, there are points $z_1 \in (x_1, c)$ and $z_2 \in (c, x_2)$ such that

$$f(x_2) = f(c) + f'(c)(x_2 - c) + \frac{1}{2}f^{(2)}(z_2)(x_2 - c)^2$$

and

$$f(x_1) = f(c) + f'(c)(x_1 - c) + \frac{1}{2}f^{(2)}(z_1)(x_1 - c)^2.$$

These two equations implies that

$$(1-t)f(x_1) + tf(x_2) = f(c) + (1-t)\frac{1}{2}f^{(2)}(z_1)(x_1 - c)^2 + t\frac{1}{2}f^{(2)}(z_2)(x_2 - c)^2 \geq f(c).$$

since $f^{(2)}(z_1)$ and $f^{(2)}(z_2)$ both are non-negative. Thus, f is convex. \square

Corollary 1.24. *Let $p > 0$. The function $f(x) := x^p$ is convex on $(0, \infty)$ if and only if $p \geq 1$.*

Proof. Note that $f^{(2)}(x) = p(p-1)x^{p-2}$ for all $x > 0$. Then the result follows immediately from Proposition 1.23. \square

Proposition 1.25. Netwon's Method: *Let f be a continuous real-valued function defined on $[a, b]$ with $f(a) < 0 < f(b)$ and $f(z) = 0$ for some $z \in (a, b)$. Assume that f is a C^2 function on (a, b) and $f'(x) \neq 0$ for all $x \in (a, b)$. Then there is $\delta > 0$ with $J := [z - \delta, z + \delta] \subseteq [a, b]$ which have the following property:*

if we fix any $x_1 \in J$ and let

$$(1.1) \quad x_{n+1} := x_n - \frac{f(x_n)}{f'(x_n)}$$

for $n = 1, 2, \dots$, then we have $z = \lim x_n$.

Proof. We first choose $r > 0$ such that $[z - r, z + r] \subseteq (a, b)$. We fix any point $x_1 \in (z - r, z + r)$ with $x_1 \neq z$. Then by the Lagrange Remainder Theorem, there is a point ξ between z and x_1 such that

$$0 = f(z) = f(x_1) + f'(x_1)(z - x_1) + \frac{1}{2}f^{(2)}(\xi)(z - x_1)^2.$$

This, together with Eq 1.1 above, we have

$$x_2 - x_1 = -\frac{f(x_1)}{f'(x_1)} = z - x_1 + \frac{f^{(2)}(\xi)}{2f'(x_1)}(z - x_1)^2.$$

Therefore, we have

$$(1.2) \quad x_2 - z = \frac{f^{(2)}(\xi)}{2f'(x_1)}(z - x_1)^2.$$

Note that the functions $f'(x)$ and $f^{(2)}(x)$ are continuous on $[z - r, z + r]$ and $f'(x) \neq 0$, hence, there is $M > 0$ such that $|\frac{f^{(2)}(u)}{2f'(v)}| \leq M$ for all $u, v \in [z - r, z + r]$. Then the Eq 1.2 implies that

$$(1.3) \quad |x_2 - z| = \left| \frac{f^{(2)}(\xi)}{2f'(x_1)} (z - x_1)^2 \right| \leq M(z - x_1)^2.$$

Choose $\delta > 0$ such that $M\delta < 1$ and $J := [z - \delta, z + \delta] \subseteq (z - r, z + r)$. Note that Now we take any $x_1 \in J$. Eq 1.3 implies that $|x_2 - z| \leq M \cdot |z - x_1|^2 \leq (M\delta) \cdot |x_1 - z| < \delta$. By using Eq 1.1 inductively, we have a sequence (x_n) in J such that

$$|x_{n+1} - z| \leq M \cdot |z - x_n|^2 \leq (M\delta) \cdot |x_n - z|$$

for all $n = 1, 2, \dots$. Therefore, we have

$$|x_{n+1} - z| \leq (M\delta)^n \cdot |x_1 - z|$$

for all $n = 1, 2, \dots$, thus, $\lim x_n = z$. The proof is complete. \square

2. RIEMANN INTEGRABLE FUNCTIONS

We will use the following notation throughout this chapter.

- (i): All functions f, g, h, \dots are bounded real valued functions defined on $[a, b]$ and $m \leq f \leq M$ on $[a, b]$.
- (ii): Let $P : a = x_0 < x_1 < \dots < x_n = b$ denote a partition on $[a, b]$; Put $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max \Delta x_i$.
- (iii): $M_i(f, P) := \sup\{f(x) : x \in [x_{i-1}, x_i]\}$; $m_i(f, P) := \inf\{f(x) : x \in [x_{i-1}, x_i]\}$.
Set $\omega_i(f, P) = M_i(f, P) - m_i(f, P)$.
- (iv): (the *upper sum* of f): $U(f, P) := \sum M_i(f, P)\Delta x_i$
(the *lower sum* of f): $L(f, P) := \sum m_i(f, P)\Delta x_i$.

Remark 2.1. *It is clear that for any partition on $[a, b]$, we always have*

- (i) $m(b - a) \leq L(f, P) \leq U(f, P) \leq M(b - a)$.
- (ii) $L(-f, P) = -U(f, P)$ and $U(-f, P) = -L(f, P)$.

The following lemma is the critical step in this section.

Lemma 2.2. *Let P and Q be the partitions on $[a, b]$. We have the following assertions.*

- (i) *If $P \subseteq Q$, then $L(f, P) \leq L(f, Q) \leq U(f, Q) \leq U(f, P)$.*
- (ii) *We always have $L(f, P) \leq U(f, Q)$.*

Proof. For Part (i), we first claim that $L(f, P) \leq L(f, Q)$ if $P \subseteq Q$. By using the induction on $l := \#Q - \#P$, it suffices to show that $L(f, P) \leq L(f, Q)$ as $l = 1$. Let $P : a = x_0 < x_1 < \dots < x_n = b$ and $Q = P \cup \{c\}$. Then $c \in (x_{s-1}, x_s)$ for some s . Notice that we have

$$m_s(f, P) \leq \min\{m_s(f, Q), m_{s+1}(f, Q)\}.$$

So, we have

$$m_s(f, P)(x_s - x_{s-1}) \leq m_s(f, Q)(c - x_{s-1}) + m_{s+1}(f, Q)(x_s - c).$$

This gives the following inequality as desired.

$$(2.1) \quad L(f, Q) - L(f, P) = m_s(f, Q)(c - x_{s-1}) + m_{s+1}(f, Q)(x_s - c) - m_s(f, P)(x_s - x_{s-1}) \geq 0.$$

Now by considering $-f$ in the Inequality 2.1 above, we see that $U(f, Q) \leq U(f, P)$.

For Part (ii), let P and Q be any pair of partitions on $[a, b]$. Notice that $P \cup Q$ is also a partition on $[a, b]$ with $P \subseteq P \cup Q$ and $Q \subseteq P \cup Q$. So, Part (i) implies that

$$L(f, P) \leq L(f, P \cup Q) \leq U(f, P \cup Q) \leq U(f, Q).$$

The proof is complete. □

The following notion plays an important role in this chapter.

Definition 2.3. *Let f be a bounded function on $[a, b]$. The upper integral (resp. lower integral) of f over $[a, b]$, write $\overline{\int_a^b} f$ (resp. $\underline{\int_a^b} f$), is defined by*

$$\overline{\int_a^b} f = \inf\{U(f, P) : P \text{ is a partation on } [a, b]\}.$$

(resp.

$$\int_a^b f = \sup\{L(f, P) : P \text{ is a partition on } [a, b]\}.)$$

Notice that the upper integral and lower integral of f must exist by Remark 2.1.

Remark 2.4. Appendix: We call a partially set (I, \leq) a *directed set* if for each pair of elements i_1 and i_2 in I , there is $i_3 \in I$ such that $i_1 \leq i_3$ and $i_2 \leq i_3$.

A *net* in \mathbb{R} is a real-valued function f defined on a directed set I , write $f = (x_i)_{i \in I}$, where $x_i := f(i)$ for $i \in I$.

We say that a net (x_i) converges to a point $L \in \mathbb{R}$ (call a limit of (x_i)) if for any $\varepsilon > 0$, there is $i_0 \in I$ such that $|x_i - L| < \varepsilon$ for all $i \geq i_0$.

Using the similar argument as in the sequence case, a limit of (x_i) is unique if it exists and we write $\lim_i x_i$ for its limits.

Example 2.5. Appendix: Using the notation given as before, let

$$I := \{P : P \text{ is a partition on } [a, b]\}.$$

We say that $P_1 \leq P_2$ for $P_1, P_2 \in I$ if $P_1 \subseteq P_2$. Clearly, I is a directed set with this order. If we put $u_P := U(f, P)$, then we have

$$\lim_P u_P = \int_a^b f.$$

In fact, let $\varepsilon > 0$. Then by the definition of an upper integral, there is $P_0 \in I$ such that

$$\int_a^b f \leq U(f, P_0) \leq \int_a^b f + \varepsilon.$$

Lemma 2.2 tells us that whenever $P \in I$ with $P \geq P_0$, we have $U(f, P) \leq U(f, P_0)$. Thus we have $|u_P - \int_a^b f| < \varepsilon$ whenever $P \geq P_0$ as desired.

Proposition 2.6. *Let f and g both are bounded functions on $[a, b]$. With the notation as above, we always have*

(i)

$$\int_a^b f \leq \int_a^b f.$$

(ii) $\int_a^b (-f) = -\int_a^b f.$

(iii)

$$\int_a^b f + \int_a^b g \leq \int_a^b (f + g) \leq \int_a^b (f + g) \leq \int_a^b f + \int_a^b g.$$

Proof. Part (i) follows from Lemma 2.2 at once.

Part (ii) is clearly obtained by $L(-f, P) = -U(f, P)$.

For proving the inequality $\int_a^b f + \int_a^b g \leq \int_a^b (f + g) \leq$ first. It is clear that we have $L(f, P) + L(g, P) \leq L(f + g, P)$ for all partitions P on $[a, b]$. Now let P_1 and P_2 be any partition on $[a, b]$. Then by Lemma 2.2, we have

$$L(f, P_1) + L(g, P_2) \leq L(f, P_1 \cup P_2) + L(g, P_1 \cup P_2) \leq L(f + g, P_1 \cup P_2) \leq \int_a^b (f + g).$$

So, we have

$$(2.2) \quad \int_a^b f + \int_a^b g \leq \int_a^b (f + g).$$

As before, we consider $-f$ and $-g$ in the Inequality 2.2, we get $\overline{\int_a^b (f + g)} \leq \overline{\int_a^b f} + \overline{\int_a^b g}$ as desired. \square

The following example shows the strict inequality in Proposition 2.6 (iii) may hold in general.

Example 2.7. Define a function $f, g : [0, 1] \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \cap \mathbb{Q}; \\ -1 & \text{otherwise.} \end{cases}$$

and

$$g(x) = \begin{cases} -1 & \text{if } x \in [0, 1] \cap \mathbb{Q}; \\ 1 & \text{otherwise.} \end{cases}$$

Then it is easy to see that $f + g \equiv 0$ and

$$\int_0^1 f = \int_0^1 g = 1 \quad \text{and} \quad \int_0^1 f = \int_0^1 g = -1.$$

So, we have

$$-2 = \int_a^b f + \int_a^b g < \int_a^b (f + g) = 0 = \overline{\int_a^b (f + g)} < \overline{\int_a^b f} + \overline{\int_a^b g} = 2.$$

We can now reaching the main definition in this chapter.

Definition 2.8. Let f be a bounded function on $[a, b]$. We say that f is Riemann integrable over $[a, b]$ if $\overline{\int_a^b f} = \underline{\int_a^b f}$. In this case, we write $\int_a^b f$ for this common value and it is called the Riemann integral of f over $[a, b]$.

Also, write $R[a, b]$ for the class of Riemann integrable functions on $[a, b]$.

Proposition 2.9. With the notation as above, $R[a, b]$ is a vector space over \mathbb{R} and the integral

$$\int_a^b : f \in R[a, b] \mapsto \int_a^b f \in \mathbb{R}$$

defines a linear functional, that is, $\alpha f + \beta g \in R[a, b]$ and $\int_a^b (\alpha f + \beta g) = \alpha \int_a^b f + \beta \int_a^b g$ for all $f, g \in R[a, b]$ and $\alpha, \beta \in \mathbb{R}$.

Proof. Let $f, g \in R[a, b]$ and $\alpha, \beta \in \mathbb{R}$. Notice that if $\alpha \geq 0$, it is clear that $\overline{\int_a^b \alpha f} = \alpha \overline{\int_a^b f} = \alpha \int_a^b f = \alpha \underline{\int_a^b f} = \underline{\int_a^b \alpha f}$. Also, if $\alpha < 0$, we have $\overline{\int_a^b \alpha f} = \alpha \underline{\int_a^b f} = \alpha \int_a^b f = \alpha \overline{\int_a^b f} = \underline{\int_a^b \alpha f}$. Therefore, we have $\int_a^b \alpha f = \alpha \int_a^b f$ for all $\alpha \in \mathbb{R}$. For showing $f + g \in R[a, b]$ and $\int_a^b (f + g) = \int_a^b f + \int_a^b g$, these will follow from Proposition 2.6 (iii) at once. The proof is finished. \square

The following result is the important characterization of a Riemann integrable function. Before showing this, we will use the following notation in the rest of this chapter.

For a partition $P : a = x_0 < x_1 < \cdots < x_n = b$ and $1 \leq i \leq n$, put

$$\omega_i(f, P) := \sup\{|f(x) - f(x')| : x, x' \in [x_{i-1}, x_i]\}.$$

It is easy to see that $U(f, P) - L(f, P) = \sum_{i=1}^n \omega_i(f, P) \Delta x_i$.

Theorem 2.10. *Let f be a bounded function on $[a, b]$. Then $f \in R[a, b]$ if and only if for all $\varepsilon > 0$, there is a partition $P : a = x_0 < \cdots < x_n = b$ on $[a, b]$ such that*

$$(2.3) \quad 0 \leq U(f, P) - L(f, P) = \sum_{i=1}^n \omega_i(f, P) \Delta x_i < \varepsilon.$$

Proof. Suppose that $f \in R[a, b]$. Let $\varepsilon > 0$. Then by the definition of the upper integral and lower integral of f , we can find the partitions P and Q such that $U(f, P) < \overline{\int_a^b} f + \varepsilon$ and $\underline{\int_a^b} f - \varepsilon < L(f, Q)$. By considering the partition $P \cup Q$, we see that

$$\underline{\int_a^b} f - \varepsilon < L(f, Q) \leq L(f, P \cup Q) \leq U(f, P \cup Q) \leq U(f, P) < \overline{\int_a^b} f + \varepsilon.$$

Since $\int_a^b f = \overline{\int_a^b} f = \underline{\int_a^b} f$, we have $0 \leq U(f, P \cup Q) - L(f, P \cup Q) < 2\varepsilon$. So, the partition $P \cup Q$ is as desired.

Conversely, let $\varepsilon > 0$, assume that the Inequality 2.3 above holds for some partition P . Notice that we have

$$L(f, P) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq U(f, P).$$

So, we have $0 \leq \overline{\int_a^b} f - \underline{\int_a^b} f < \varepsilon$ for all $\varepsilon > 0$. The proof is finished. \square

Remark 2.11. *Theorem 8.3 tells us that a bounded function f is Riemann integrable over $[a, b]$ if and only if the “size” of the discontinuous set of f is arbitrary small.*

Example 2.12. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be the function defined by*

$$f(x) = \begin{cases} \frac{1}{p} & \text{if } x = \frac{q}{p}, \text{ where } p, q \text{ are relatively prime positive integers;} \\ 0 & \text{otherwise.} \end{cases}$$

Then $f \in R[0, 1]$.

(Notice that the set of all discontinuous points of f , say D , is just the set of all $(0, 1] \cap \mathbb{Q}$. Since the set $(0, 1] \cap \mathbb{Q}$ is countable, we can write $(0, 1] \cap \mathbb{Q} = \{z_1, z_2, \dots\}$. So, if we let $m(D)$ be the “size” of the set D , then $m(D) = m(\bigcup_{i=1}^{\infty} \{z_i\}) = \sum_{i=1}^{\infty} m(\{z_i\}) = 0$, in here, you may think that the size of each set $\{z_i\}$ is 0.)

Proof. Let $\varepsilon > 0$. By Theorem 8.3, it aims to find a partition P on $[0, 1]$ such that

$$U(f, P) - L(f, P) < \varepsilon.$$

Notice that for $x \in [0, 1]$ such that $f(x) \geq \varepsilon$ if and only if $x = q/p$ for a pair of relatively prime positive integers p, q with $\frac{1}{p} \geq \varepsilon$. Since $1 \leq q \leq p$, there are only finitely many pairs of relatively prime positive integers p and q such that $f(\frac{q}{p}) \geq \varepsilon$. So, if we let $S := \{x \in [0, 1] : f(x) \geq \varepsilon\}$, then S is a finite subset

of $[0, 1]$. Let L be the number of the elements in S . Then, for any partition $P : a = x_0 < \cdots < x_n = 1$, we have

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i = \left(\sum_{i: [x_{i-1}, x_i] \cap S = \emptyset} + \sum_{i: [x_{i-1}, x_i] \cap S \neq \emptyset} \right) \omega_i(f, P) \Delta x_i.$$

Notice that if $[x_{i-1}, x_i] \cap S = \emptyset$, then we have $\omega_i(f, P) \leq \varepsilon$ and thus,

$$\sum_{i: [x_{i-1}, x_i] \cap S = \emptyset} \omega_i(f, P) \Delta x_i \leq \varepsilon \sum_{i: [x_{i-1}, x_i] \cap S = \emptyset} \Delta x_i \leq \varepsilon(1 - 0).$$

On the other hand, since there are at most $2L$ sub-intervals $[x_{i-1}, x_i]$ such that $[x_{i-1}, x_i] \cap S \neq \emptyset$ and $\omega_i(f, P) \leq 1$ for all $i = 1, \dots, n$, so, we have

$$\sum_{i: [x_{i-1}, x_i] \cap S \neq \emptyset} \omega_i(f, P) \Delta x_i \leq 1 \cdot \sum_{i: [x_{i-1}, x_i] \cap S \neq \emptyset} \Delta x_i \leq 2L \|P\|.$$

We can now conclude that for any partition P , we have

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i \leq \varepsilon + 2L \|P\|.$$

So, if we take a partition P with $\|P\| < \varepsilon/(2L)$, then we have $\sum_{i=1}^n \omega_i(f, P) \Delta x_i \leq 2\varepsilon$. The proof is finished. \square

Proposition 2.13. *Let f be a function defined on $[a, b]$. If f is either monotone or continuous on $[a, b]$, then $f \in R[a, b]$.*

Proof. We first show the case of f being monotone. We may assume that f is monotone increasing. Notice that for any partition $P : a = x_0 < \cdots < x_n = b$, we have $\omega_i(f, P) = f(x_i) - f(x_{i-1})$. So, if $\|P\| < \varepsilon$, we have

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i = \sum_{i=1}^n (f(x_i) - f(x_{i-1})) \Delta x_i < \|P\| \sum_{i=1}^n (f(x_i) - f(x_{i-1})) = \|P\| (f(b) - f(a)) < \varepsilon (f(b) - f(a)).$$

Therefore, $f \in R[a, b]$ if f is monotone.

Suppose that f is continuous on $[a, b]$. Then f is uniform continuous on $[a, b]$. Then for any $\varepsilon > 0$, there is $\delta > 0$ such that $|f(x) - f(x')| < \varepsilon$ as $x, x' \in [a, b]$ with $|x - x'| < \delta$. So, if we choose a partition P with $\|P\| < \delta$, then $\omega_i(f, P) < \varepsilon$ for all i . This implies that

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i \leq \varepsilon \sum_{i=1}^n \Delta x_i = \varepsilon(b - a).$$

The proof is complete. \square

Proposition 2.14. *We have the following assertions.*

(i) *If $f, g \in R[a, b]$ with $f \leq g$, then $\int_a^b f \leq \int_a^b g$.*

(ii) *If $f \in R[a, b]$, then the absolute valued function $|f| \in R[a, b]$. In this case, we have $|\int_a^b f| \leq \int_a^b |f|$.*

Proof. For Part (i), it is clear that we have the inequality $U(f, P) \leq U(g, P)$ for any partition P . So, we have $\int_a^b f = \overline{\int_a^b f} \leq \overline{\int_a^b g} = \int_a^b g$.

For Part (ii), the integrability of $|f|$ follows immediately from Theorem 8.3 and the simple inequality $||f|(x') - |f|(x'')| \leq |f(x') - f(x'')|$ for all $x', x'' \in [a, b]$. Thus, we have $U(|f|, P) - L(|f|, P) \leq$

$U(f, P) - L(f, P)$ for any partition P on $[a, b]$.

Finally, since we have $-f \leq |f| \leq f$, by Part (i), we have $|\int_a^b f| \leq \int_a^b |f|$ at once. \square

Proposition 2.15. *Let $a < c < b$. We have $f \in R[a, b]$ if and only if the restrictions $f|_{[a, c]} \in R[a, c]$ and $f|_{[c, b]} \in R[c, b]$. In this case we have*

$$(2.4) \quad \int_a^b f = \int_a^c f + \int_c^b f.$$

Proof. Let $f_1 := f|_{[a, c]}$ and $f_2 := f|_{[c, b]}$.

It is clear that we always have

$$U(f_1, P_1) - L(f_1, P_1) + U(f_2, P_2) - L(f_2, P_2) = U(P, f) - L(f, P)$$

for any partition P_1 on $[a, c]$ and P_2 on $[c, b]$ with $P = P_1 \cup P_2$.

From this, we can show the sufficient condition at once.

For showing the necessary condition, since $f \in R[a, b]$, for any $\varepsilon > 0$, there is a partition Q on $[a, b]$ such that $U(f, Q) - L(f, Q) < \varepsilon$ by Theorem 8.3. Notice that there are partitions P_1 and P_2 on $[a, c]$ and $[c, b]$ respectively such that $P := Q \cup \{c\} = P_1 \cup P_2$. Thus, we have

$$U(f_1, P_1) - L(f_1, P_1) + U(f_2, P_2) - L(f_2, P_2) = U(f, P) - L(f, P) \leq U(f, Q) - L(f, Q) < \varepsilon.$$

So, we have $f_1 \in R[a, c]$ and $f_2 \in R[c, b]$.

It remains to show the Equation 2.4 above. Notice that for any partition P_1 on $[a, c]$ and P_2 on $[c, b]$, we have

$$L(f_1, P_1) + L(f_2, P_2) = L(f, P_1 \cup P_2) \leq \int_a^b f = \int_a^b f.$$

So, we have $\int_a^c f + \int_c^b f \leq \int_a^b f$. Then the inverse inequality can be obtained at once by considering the function $-f$. Then the result is obtained by using Theorem 8.3. \square

Proposition 2.16. *Let f and g be Riemann integrable functions defined on $[a, b]$. Then the pointwise product function $f \cdot g \in R[a, b]$.*

Proof. We first show that the square function f^2 is Riemann integrable. In fact, if we let $M = \sup\{|f(x)| : x \in [a, b]\}$, then we have $\omega_k(f^2, P) \leq 2M\omega_k(f, P)$ for any partition $P : a = x_0 < \dots < x_n = b$ because we always have $|f^2(x) - f^2(x')| \leq 2M|f(x) - f(x')|$ for all $x, x' \in [a, b]$. Then by Theorem 8.3, the square function $f^2 \in R[a, b]$.

This, together with the identity $f \cdot g = \frac{1}{2}((f+g)^2 - f^2 - g^2)$. The result follows. \square

Remark 2.17. *In the proof of Proposition 2.16, we have shown that if $f \in R[a, b]$, then so is its square function f^2 . However, the converse does not hold. For example, if we consider $f(x) = 1$ for $x \in \mathbb{Q} \cap [0, 1]$ and $f(x) = -1$ for $x \in \mathbb{Q}^c \cap [0, 1]$, then $f \notin R[0, 1]$ but $f^2 \equiv 1$ on $[0, 1]$.*

Proposition 2.18. (Mean Value Theorem for Integrals)

Let f and g be the functions defined on $[a, b]$. Assume that f is continuous and g is a non-negative Riemann integrable function. Then, there is a point $\xi \in (a, b)$ such that

$$(2.5) \quad \int_a^b f(x)g(x)dx = f(\xi) \int_a^b g(x)dx.$$

Proof. By the continuity of f on $[a, b]$, there exist two points x_1 and x_2 in $[a, b]$ such that

$$f(x_1) = m := \min f(x); \text{ and } f(x_2) = M := \max f(x).$$

We may assume that $a \leq x_1 < x_2 \leq b$. From this, since $g \leq 0$, we have

$$mg(x) \leq f(x)g(x) \leq Mg(x)$$

for all $x \in [a, b]$. From this and Proposition 2.16 above, we have

$$m \int_a^b g \leq \int_a^b fg \leq M \int_a^b g.$$

So, if $\int_a^b g = 0$, then the result follows at once.

We may now suppose that $\int_a^b g > 0$. The above inequality shows that

$$m = f(x_1) \leq \frac{\int_a^b fg}{\int_a^b g} \leq f(x_2) = M.$$

Therefore, there is a point $\xi \in [x_1, x_2] \subseteq [a, b]$ so that the Equation 2.5 holds by using the Intermediate Value Theorem for the function f . Thus, it remains to show that such element ξ can be chosen in (a, b) .

Let $a \leq x_1 < x_2 \leq b$ be as above.

If x_1 and x_2 can be found so that $a < x_1 < x_2 < b$, then the result is proved immediately since $\xi \in [x_1, x_2] \subset (a, b)$ in this case.

Now suppose that x_1 or x_2 does not exist in (a, b) , i.e., $m = f(a) < f(x)$ for all $x \in (a, b]$ or $f(x) < f(b) = M$ for all $x \in [a, b)$.

Claim 1: If $f(a) < f(x)$ for all $x \in (a, b]$, then $\int_a^b fg > f(a) \int_a^b g$ and hence, $\xi \in (a, x_2] \subseteq (a, b]$.

For showing **Claim 1**, put $h(x) := f(x) - f(a)$ for $x \in [a, b]$. Then h is continuous on $[a, b]$ and $h > 0$ on $(a, b]$. This implies that $\int_c^d h > 0$ for any subinterval $[c, d] \subseteq [a, b]$. (**Why?**)

On the other hand, since $\int_a^b g = \int_a^b g > 0$, there is a partition $P : a = x_0 < \dots < x_n = b$ so that $L(g, P) > 0$. This implies that $m_k(g, P) > 0$ for some sub-interval $[x_{k-1}, x_k]$. Therefore, we have

$$\int_a^b hg \geq \int_{x_{k-1}}^{x_k} hg \geq m_k(g, P) \int_{x_{k-1}}^{x_k} h > 0.$$

Hence, we have $\int_a^b fg > f(a) \int_a^b g$. **Claim 1** follows.

Similarly, one can show that if $f(x) < f(b) = M$ for all $x \in [a, b)$, then we have $\int_a^b fg < f(b) \int_a^b g$.

This, together with **Claim 1** give us that such ξ can be found in (a, b) . The proof is finished. \square

Now if $f \in R[a, b]$, then by Proposition 2.15, we can define a function $F : [a, b] \rightarrow \mathbb{R}$ by

$$(2.6) \quad F(c) = \begin{cases} 0 & \text{if } c = a \\ \int_a^c f & \text{if } a < c \leq b. \end{cases}$$

Theorem 2.19. Fundamental Theorem of Calculus: *With the notation as above, assume that $f \in R[a, b]$, we have the following assertion.*

- (i) *If there is a continuous function F on $[a, b]$ which is differentiable on (a, b) with $F' = f$, then $\int_a^b f = F(b) - F(a)$. In this case, F is called an indefinite integral of f . (**note:** if F_1 and F_2 both are the indefinite integrals of f , then by the Mean Value Theorem, we have $F_2 = F_1 + \text{constant}$).*
- (ii) *The function F defined as in Eq. 2.6 above is continuous on $[a, b]$. Furthermore, if f is continuous on $[a, b]$, then F' exists on (a, b) and $F' = f$ on (a, b) .*

Proof. For Part (i), notice that for any partition $P : a = x_0 < \cdots < x_n = b$, then by the Mean Value Theorem, for each $[x_{i-1}, x_i]$, there is $\xi_i \in (x_{i-1}, x_i)$ such that $F(x_i) - F(x_{i-1}) = F'(\xi_i)\Delta x_i = f(\xi_i)\Delta x_i$. So, we have

$$L(f, P) \leq \sum f(\xi_i)\Delta x_i = \sum F(x_i) - F(x_{i-1}) = F(b) - F(a) \leq U(f, P)$$

for all partitions P on $[a, b]$. This gives

$$\int_a^b f = \int_a^b f \leq F(b) - F(a) \leq \overline{\int_a^b f} = \int_a^b f$$

as desired.

For showing the continuity of F in Part (ii), let $a < c < x < b$. If $|f| \leq M$ on $[a, b]$, then we have $|F(x) - F(c)| = |\int_c^x f| \leq M(x - c)$. So, $\lim_{x \rightarrow c^+} F(x) = F(c)$. Similarly, we also have $\lim_{x \rightarrow c^-} F(x) = F(c)$. Thus F is continuous on $[a, b]$.

Now assume that f is continuous on $[a, b]$. Notice that for any $t > 0$ with $a < c < c + t < b$, we have

$$\inf_{x \in [c, c+t]} f(x) \leq \frac{1}{t}(F(c+t) - F(c)) = \frac{1}{t} \int_c^{c+t} f \leq \sup_{x \in [c, c+t]} f(x).$$

Since f is continuous at c , we see that $\lim_{t \rightarrow 0^+} \frac{1}{t}(F(c+t) - F(c)) = f(c)$. Similarly, we have $\lim_{t \rightarrow 0^-} \frac{1}{t}(F(c+t) - F(c)) = f(c)$. So, we have $F'(c) = f(c)$ as desired. The proof is finished. \square

Definition 2.20. For each function f on $[a, b]$ and a partition $P : a = x_0 < \cdots < x_n = b$, we call $R(f, P, \{\xi_i\}) := \sum_{i=1}^n f(\xi_i)\Delta x_i$, where $\xi_i \in [x_{i-1}, x_i]$, the Riemann sum of f over $[a, b]$.

We say that the Riemann sum $R(f, P, \{\xi_i\})$ converges to a number A as $\|P\| \rightarrow 0$, write $A = \lim_{\|P\| \rightarrow 0} R(f, P, \{\xi_i\})$, if for any $\varepsilon > 0$, there is $\delta > 0$ such that

$$|A - R(f, P, \{\xi_i\})| < \varepsilon$$

whenever $\|P\| < \delta$ and for any $\xi_i \in [x_{i-1}, x_i]$.

Proposition 2.21. Let f be a function defined on $[a, b]$. If the limit $\lim_{\|P\| \rightarrow 0} R(f, P, \{\xi_i\}) = A$ exists, then f is automatically bounded.

Proof. Suppose that f is unbounded. Then by the assumption, there exists a partition $P : a = x_0 < \cdots < x_n = b$ such that $|\sum_{k=1}^n f(\xi_k)\Delta x_k| < 1 + |A|$ for any $\xi_k \in [x_{k-1}, x_k]$. Since f is unbounded, we may assume that f is unbounded on $[a, x_1]$. In particular, we choose $\xi_k = x_k$ for $k = 2, \dots, n$. Also, we can choose $\xi_1 \in [a, x_1]$ such that

$$|f(\xi_1)|\Delta x_1 > 1 + |A| + \left| \sum_{k=2}^n f(x_k)\Delta x_k \right|.$$

It leads to a contradiction because we have $1 + |A| > |f(\xi_1)|\Delta x_1 - \left| \sum_{k=2}^n f(x_k)\Delta x_k \right|$. The proof is finished. \square

Lemma 2.22. $f \in R[a, b]$ if and only if for any $\varepsilon > 0$, there is $\delta > 0$ such that $U(f, P) - L(f, P) < \varepsilon$ whenever $\|P\| < \delta$.

Proof. The converse follows from Theorem 8.3.

Assume that f is integrable over $[a, b]$. Let $\varepsilon > 0$. Then there is a partition $Q : a = y_0 < \dots < y_l = b$ on

$[a, b]$ such that $U(f, Q) - L(f, Q) < \varepsilon$. Now take $0 < \delta < \varepsilon/l$. Suppose that $P : a = x_0 < \dots < x_n = b$ with $\|P\| < \delta$. Then we have

$$U(f, P) - L(f, P) = I + II$$

where

$$I = \sum_{i: Q \cap [x_{i-1}, x_i] = \emptyset} \omega_i(f, P) \Delta x_i;$$

and

$$II = \sum_{i: Q \cap [x_{i-1}, x_i] \neq \emptyset} \omega_i(f, P) \Delta x_i$$

Notice that we have

$$I \leq U(f, Q) - L(f, Q) < \varepsilon$$

and

$$II \leq (M - m) \sum_{i: Q \cap [x_{i-1}, x_i] \neq \emptyset} \Delta x_i \leq (M - m) \cdot 2l \cdot \frac{\varepsilon}{l} = 2(M - m)\varepsilon.$$

The proof is finished. □

Theorem 2.23. $f \in R[a, b]$ if and only if the Riemann sum $R(f, P, \{\xi_i\})$ is convergent. In this case, $R(f, P, \{\xi_i\})$ converges to $\int_a^b f(x)dx$ as $\|P\| \rightarrow 0$.

Proof. For the proof (\Rightarrow): we first note that we always have

$$L(f, P) \leq R(f, P, \{\xi_i\}) \leq U(f, P)$$

and

$$L(f, P) \leq \int_a^b f(x)dx \leq U(f, P)$$

for any partition P and $\xi_i \in [x_{i-1}, x_i]$.

Now let $\varepsilon > 0$. Lemma 2.22 gives $\delta > 0$ such that $U(f, P) - L(f, P) < \varepsilon$ as $\|P\| < \delta$. Then we have

$$\left| \int_a^b f(x)dx - R(f, P, \{\xi_i\}) \right| < \varepsilon$$

as $\|P\| < \delta$ and $\xi_i \in [x_{i-1}, x_i]$. The necessary part is proved and $R(f, P, \{\xi_i\})$ converges to $\int_a^b f(x)dx$.

For (\Leftarrow): assume that there is a number A such that for any $\varepsilon > 0$, there is $\delta > 0$, we have

$$A - \varepsilon < R(f, P, \{\xi_i\}) < A + \varepsilon$$

for any partition P with $\|P\| < \delta$ and $\xi_i \in [x_{i-1}, x_i]$.

Note that f is automatically bounded in this case by Proposition 2.21.

Now fix a partition P with $\|P\| < \delta$. Then for each $[x_{i-1}, x_i]$, choose $\xi_i \in [x_{i-1}, x_i]$ such that $M_i(f, P) - \varepsilon \leq f(\xi_i)$. This implies that we have

$$U(f, P) - \varepsilon(b - a) \leq R(f, P, \{\xi_i\}) < A + \varepsilon.$$

Thus, we have shown that for any $\varepsilon > 0$, there is a partition \mathcal{P} such that

$$(2.7) \quad \int_a^b f(x)dx \leq U(f, P) \leq A + \varepsilon(1 + b - a).$$

By considering $-f$, note that the Riemann sum of $-f$ will converge to $-A$. The inequality 8.1 will imply that for any $\varepsilon > 0$, there is a partition P such that

$$A - \varepsilon(1 + b - a) \leq \int_a^b f(x)dx \leq \overline{\int_a^b f(x)dx} \leq A + \varepsilon(1 + b - a).$$

The proof is complete. \square

Theorem 2.24. *Let $f \in R[c, d]$ and let $\phi : [a, b] \rightarrow [c, d]$ be a strictly increasing C^1 function with $f(a) = c$ and $f(b) = d$.*

Then $f \circ \phi \in R[a, b]$, moreover, we have

$$\int_c^d f(x)dx = \int_a^b f(\phi(t))\phi'(t)dt.$$

Proof. Let $A = \int_c^d f(x)dx$. By using Theorem 2.23, we need to show that for all $\varepsilon > 0$, there is $\delta > 0$ such that

$$|A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| < \varepsilon$$

for all $\xi_k \in [t_{k-1}, t_k]$ whenever $Q : a = t_0 < \dots < t_m = b$ with $\|Q\| < \delta$.

Now let $\varepsilon > 0$. Then by Lemma 2.22 and Theorem 2.23, there is $\delta_1 > 0$ such that

$$(2.8) \quad |A - \sum f(\eta_k)\Delta x_k| < \varepsilon$$

and

$$(2.9) \quad \sum \omega_k(f, P)\Delta x_k < \varepsilon$$

for all $\eta_k \in [x_{k-1}, x_k]$ whenever $P : c = x_0 < \dots < x_m = d$ with $\|P\| < \delta_1$.

Now put $x = \phi(t)$ for $t \in [a, b]$.

Now since ϕ and ϕ' are continuous on $[a, b]$, there is $\delta > 0$ such that $|\phi(t) - \phi(t')| < \delta_1$ and $|\phi'(t) - \phi'(t')| < \varepsilon$ for all t, t' in $[a, b]$ with $|t - t'| < \delta$.

Now let $Q : a = t_0 < \dots < t_m = b$ with $\|Q\| < \delta$. If we put $x_k = \phi(t_k)$, then $P : c = x_0 < \dots < x_m = d$ is a partition on $[c, d]$ with $\|P\| < \delta_1$ because ϕ is strictly increasing.

Note that the Mean Value Theorem implies that for each $[t_{k-1}, t_k]$, there is $\xi_k^* \in (t_{k-1}, t_k)$ such that

$$\Delta x_k = \phi(t_k) - \phi(t_{k-1}) = \phi'(\xi_k^*)\Delta t_k.$$

This yields that

$$(2.10) \quad |\Delta x_k - \phi'(\xi_k)\Delta t_k| < \varepsilon\Delta t_k$$

for any $\xi_k \in [t_{k-1}, t_k]$ for all $k = 1, \dots, m$ because of the choice of δ .

Now for any $\xi_k \in [t_{k-1}, t_k]$, we have

$$(2.11) \quad \begin{aligned} |A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| &\leq |A - \sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k| \\ &+ |\sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k - \sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k| \\ &+ |\sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| \end{aligned}$$

Notice that inequality 8.2 implies that

$$|A - \sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k| = |A - \sum f(\phi(\xi_k^*))\Delta x_k| < \varepsilon.$$

Moreover, since we have $|\phi'(\xi_k^*) - \phi'(\xi_k)| < \varepsilon$ for all $k = 1, \dots, m$, we have

$$|\sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k - \sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k| \leq M(b - a)\varepsilon$$

where $|f(x)| \leq M$ for all $x \in [c, d]$.

On the other hand, by using inequality 8.4 we have

$$|\phi'(\xi_k)\Delta t_k| \leq \Delta x_k + \varepsilon\Delta t_k$$

for all k . This, together with inequality 8.3 imply that

$$\begin{aligned} & \left| \sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k \right| \\ & \leq \sum \omega_k(f, P)|\phi'(\xi_k)\Delta t_k| \quad (\because \phi(\xi_k^*), \phi(\xi_k) \in [x_{k-1}, x_k]) \\ & \leq \sum \omega_k(f, P)(\Delta x_k + \varepsilon\Delta t_k) \\ & \leq \varepsilon + 2M(b-a)\varepsilon. \end{aligned}$$

Finally by inequality 8.5, we have

$$|A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| \leq \varepsilon + M(b-a)\varepsilon + \varepsilon + 2M(b-a)\varepsilon.$$

The proof is complete. \square

3. IMPROPER RIEMANN INTEGRALS

Definition 3.1. Let $-\infty < a < b < \infty$.

(i) Let f be a function defined on $[a, \infty)$. Assume that the restriction $f|_{[a, T]}$ is integrable over

$[a, T]$ for all $T > a$. Put $\int_a^\infty f := \lim_{T \rightarrow \infty} \int_a^T f$ if this limit exists.

Similarly, we can define $\int_{-\infty}^b f$ if f is defined on $(-\infty, b]$.

(ii) If f is defined on $(a, b]$ and $f|_{[c, b]} \in R[c, b]$ for all $a < c < b$. Put $\int_a^b f := \lim_{c \rightarrow a^+} \int_c^b f$ if it exists.

Similarly, we can define $\int_a^b f$ if f is defined on $[a, b)$.

(iii) As f is defined on \mathbb{R} , if $\int_0^\infty f$ and $\int_{-\infty}^0 f$ both exist, then we put $\int_{-\infty}^\infty f = \int_{-\infty}^0 f + \int_0^\infty f$.

In the cases above, we call the resulting limits the improper Riemann integrals of f and say that the integrals are convergent.

Example 3.2. Define (formally) an improper integral $\Gamma(s)$ (called the Γ -function) as follows:

$$\Gamma(s) := \int_0^\infty x^{s-1}e^{-x}dx$$

for $s \in \mathbb{R}$. Then $\Gamma(s)$ is convergent if and only if $s > 0$.

Proof. Put $I(s) := \int_0^1 x^{s-1}e^{-x}dx$ and $II(s) := \int_1^\infty x^{s-1}e^{-x}dx$. We first claim that the integral $II(s)$ is convergent for all $s \in \mathbb{R}$.

In fact, if we fix $s \in \mathbb{R}$, then we have

$$\lim_{x \rightarrow \infty} \frac{x^{s-1}}{e^{x/2}} = 0.$$

So there is $M > 1$ such that $\frac{x^{s-1}}{e^{x/2}} \leq 1$ for all $x \geq M$. Thus we have

$$0 \leq \int_M^\infty x^{s-1}e^{-x}dx \leq \int_M^\infty e^{-x/2}dx < \infty.$$

Therefore we need to show that the integral $I(s)$ is convergent if and only if $s > 0$. Note that for $0 < \eta < 1$, we have

$$0 \leq \int_{\eta}^1 x^{s-1} e^{-x} dx \leq \int_{\eta}^1 x^{s-1} dx = \begin{cases} \frac{1}{s}(1 - \eta^s) & \text{if } s - 1 \neq -1; \\ -\ln \eta & \text{otherwise.} \end{cases}$$

Thus the integral $I(s) = \lim_{\eta \rightarrow 0^+} \int_{\eta}^1 x^{s-1} e^{-x} dx$ is convergent if $s > 0$.

Conversely, we also have

$$\int_{\eta}^1 x^{s-1} e^{-x} dx \geq e^{-1} \int_{\eta}^1 x^{s-1} dx = \begin{cases} \frac{e^{-1}}{s}(1 - \eta^s) & \text{if } s - 1 \neq -1; \\ -e^{-1} \ln \eta & \text{otherwise.} \end{cases}$$

So if $s \leq 0$, then $\int_{\eta}^1 x^{s-1} e^{-x} dx$ is divergent as $\eta \rightarrow 0^+$. The result follows. \square

4. SOME RESULTS OF SEQUENCES OF FUNCTIONS

Proposition 4.1. *Let $f_n : (a, b) \rightarrow \mathbb{R}$ be a sequence of functions. Assume that it satisfies the following conditions:*

- (i) : $f_n(x)$ point-wise converges to a function $f(x)$ on (a, b) ;
- (ii) : each f_n is a C^1 function on (a, b) ;
- (iii) : $f'_n \rightarrow g$ uniformly on (a, b) .

Then f is a C^1 -function on (a, b) with $f' = g$.

Proof. Fix $c \in (a, b)$. Then for each x with $c < x < b$ (similarly, we can prove it in the same way as $a < x < c$), the Fundamental Theorem of Calculus implies that

$$f_n(x) = \int_c^x f'_n(t) dt + f_n(c).$$

Since $f'_n \rightarrow g$ uniformly on (a, b) , we see that

$$\int_c^x f'_n(t) dt \rightarrow \int_c^x g(t) dt.$$

This gives

$$(4.1) \quad f(x) = \int_c^x g(t) dt + f(c).$$

for all $x \in (c, b)$. Similarly, we have $f(x) = \int_c^x g(t) dt + f(c)$ for all $x \in (a, b)$.

On the other hand, g is continuous on (a, b) since each f'_n is continuous and $f'_n \rightarrow g$ uniformly on (a, b) . Equation 9.1 will tell us that f' exists and $f' = g$ on (a, b) . The proof is finished. \square

Proposition 4.2. *Let (f_n) be a sequence of differentiable functions defined on (a, b) . Assume that*

- (i): there is a point $c \in (a, b)$ such that $\lim f_n(c)$ exists;
- (ii): f'_n converges uniformly to a function g on (a, b) .

Then

- (a): f_n converges uniformly to a function f on (a, b) ;
- (b): f is differentiable on (a, b) and $f' = g$.

Proof. For Part (a), we will make use the Cauchy theorem.

Let $\varepsilon > 0$. Then by the assumptions (i) and (ii), there is a positive integer N such that

$$|f_m(c) - f_n(c)| < \varepsilon \quad \text{and} \quad |f'_m(x) - f'_n(x)| < \varepsilon$$

for all $m, n \geq N$ and for all $x \in (a, b)$. Now fix $c < x < b$ and $m, n \geq N$. To apply the Mean Value Theorem for $f_m - f_n$ on (c, x) , then there is a point ξ between c and x such that

$$(4.2) \quad f_m(x) - f_n(x) = f_m(c) - f_n(c) + (f'_m(\xi) - f'_n(\xi))(x - c).$$

This implies that

$$|f_m(x) - f_n(x)| \leq |f_m(c) - f_n(c)| + |f'_m(\xi) - f'_n(\xi)||x - c| < \varepsilon + (b - a)\varepsilon$$

for all $m, n \geq N$ and for all $x \in (c, b)$. Similarly, when $x \in (a, c)$, we also have

$$|f_m(x) - f_n(x)| < \varepsilon + (b - a)\varepsilon.$$

So Part (a) follows.

Let f be the uniform limit of (f_n) on (a, b)

For Part (b), we fix $u \in (a, b)$. We are going to show

$$\lim_{x \rightarrow u} \frac{f(x) - f(u)}{x - u} = g(u).$$

Let $\varepsilon > 0$. Since (f'_n) is uniformly convergent on (a, b) , there is $N \in \mathbb{N}$ such that

$$(4.3) \quad |f'_m(x) - f'_n(x)| < \varepsilon$$

for all $m, n \geq N$ and for all $x \in (a, b)$

Note that for all $m \geq N$ and $x \in (a, b) \setminus \{u\}$, applying the Mean value Theorem for $f_m - f_N$ as before, we have

$$\frac{f_m(x) - f_N(x)}{x - u} = \frac{f_m(u) - f_N(u)}{x - u} + (f'_m(\xi) - f'_N(\xi))$$

for some ξ between u and x .

So Eq.9.3 implies that

$$(4.4) \quad \left| \frac{f_m(x) - f_m(u)}{x - u} - \frac{f_N(x) - f_N(u)}{x - u} \right| \leq \varepsilon$$

for all $m \geq N$ and for all $x \in (a, b)$ with $x \neq u$.

Taking $m \rightarrow \infty$ in Eq.9.4, we have

$$\left| \frac{f(x) - f(u)}{x - u} - \frac{f_N(x) - f_N(u)}{x - u} \right| \leq \varepsilon.$$

Hence we have

$$\begin{aligned} \left| \frac{f(x) - f(u)}{x - u} - f'_N(u) \right| &\leq \left| \frac{f(x) - f(u)}{x - u} - \frac{f_N(x) - f_N(u)}{x - u} \right| + \left| \frac{f_N(x) - f_N(u)}{x - u} - f'_N(u) \right| \\ &\leq \varepsilon + \left| \frac{f_N(x) - f_N(u)}{x - u} - f'_N(u) \right|. \end{aligned}$$

So if we can take $0 < \delta$ such that $\left| \frac{f_N(x) - f_N(u)}{x - u} - f'_N(u) \right| < \varepsilon$ for $0 < |x - u| < \delta$, then we have

$$(4.5) \quad \left| \frac{f(x) - f(u)}{x - u} - f'_N(u) \right| \leq 2\varepsilon$$

for $0 < |x - u| < \delta$. On the other hand, by the choice of N , we have $|f'_m(y) - f'_N(y)| < \varepsilon$ for all $y \in (a, b)$ and $m \geq N$. So we have $|g(u) - f'_N(u)| \leq \varepsilon$. This together with Eq.9.5 give

$$\left| \frac{f(x) - f(u)}{x - u} - g(u) \right| \leq 3\varepsilon$$

as $0 < |x - u| < \delta$, that is we have

$$\lim_{x \rightarrow u} \frac{f(x) - f(u)}{x - u} = g(u).$$

The proof is finished. □

Remark 4.3. *The uniform convergence assumption of (f'_n) in the Propositions above is essential.*

Example 4.4. *Let $f_n(x) := \frac{x}{1+n^2x^2}$ for $x \in (-1, 1)$. Then we have*

$$g(x) := \lim_n f'_n(x) := \lim_n \frac{1 - n^2x^2}{(1 + n^2x^2)^2} = \begin{cases} 0 & \text{if } x \neq 0; \\ 1 & \text{if } x = 0. \end{cases}$$

On the other hand, $f_n \rightarrow 0$ uniformly on $(-1, 1)$. In fact, if $f'_n(1/n) = 0$ for all $n = 1, 2, \dots$, then f_n attains the maximal value $f_n(1/n) = \frac{1}{2n}$ at $x = 1/n$ for each $n = 1, \dots$ and hence, $f_n \rightarrow 0$ uniformly on $(-1, 1)$.

So Propositions 9.1 and 9.2 does not hold. Note that (f'_n) does not converge uniformly to g on $(-1, 1)$.

Proposition 4.5. (Dini's Theorem): *Let A be a compact subset of \mathbb{R} and $f_n : A \rightarrow \mathbb{R}$ be a sequence of continuous functions defined on A . Suppose that*

- (i) *for each $x \in A$, we have $f_n(x) \leq f_{n+1}(x)$ for all $n = 1, 2, \dots$;*
- (ii) *the pointwise limit $f(x) := \lim_n f_n(x)$ exists for all $x \in A$;*
- (iii) *f is continuous on A .*

Then f_n converges to f uniformly on A .

Proof. Let $g_n := f - f_n$ defined on A . Then each g_n is continuous and $g_n(x) \downarrow 0$ pointwise on A . It suffices to show that g_n converges to 0 uniformly on A .

Method I: Suppose not. Then there is $\varepsilon > 0$ such that for all positive integer N , we have

$$(4.6) \quad g_n(x_n) \geq \varepsilon.$$

for some $n \geq N$ and some $x_n \in A$. From this, by passing to a subsequence we may assume that $g_n(x_n) \geq \varepsilon$ for all $n = 1, 2, \dots$. Then by using the compactness of A , there is a convergent subsequence (x_{n_k}) of (x_n) in A . Let $z := \lim_k x_{n_k} \in A$. Since $g_{n_k}(z) \downarrow 0$ as $k \rightarrow \infty$. So, there is a positive integer K such that $0 \leq g_{n_K}(z) < \varepsilon/2$. Since g_{n_K} is continuous at z and $\lim_i x_{n_i} = z$, we have $\lim_i g_{n_K}(x_{n_i}) = g_{n_K}(z)$. So, we can choose i large enough such that $i > K$

$$g_{n_i}(x_{n_i}) \leq g_{n_K}(x_{n_i}) < \varepsilon/2$$

because $g_m(x_{n_i}) \downarrow 0$ as $m \rightarrow \infty$. This contradicts to the Inequality 4.6.

Method II: Let $\varepsilon > 0$. Fix $x \in A$. Since $g_n(x) \downarrow 0$, there is $N(x) \in \mathbb{N}$ such that $0 \leq g_n(x) < \varepsilon$ for all $n \geq N(x)$. Since $g_{N(x)}$ is continuous, there is $\delta(x) > 0$ such that $g_{N(x)}(y) < \varepsilon$ for all $y \in A$ with $|x - y| < \delta(x)$. If we put $J_x := (x - \delta(x), x + \delta(x))$, then $A \subseteq \bigcup_{x \in A} J_x$. Then by the compactness of A , there are finitely many x_1, \dots, x_m in A such that $A \subseteq J_{x_1} \cup \dots \cup J_{x_m}$. Put $N := \max(N(x_1), \dots, N(x_m))$. Now if $y \in A$, then $y \in J(x_i)$ for some $1 \leq i \leq m$. This implies that

$$g_n(y) \leq g_{N(x_i)}(y) < \varepsilon$$

for all $n \geq N \geq N(x_i)$. □

REFERENCES

- [1] R.G. Bartle and D.R. Sherbert, Introduction to real analysis, Fourth edition, Wiley, (2011).