

THE CHINESE UNIVERSITY OF HONG KONG  
Department of Mathematics  
**MATH2050C Mathematical Analysis I**  
**Tutorial 9 (March 25)**

**Definition.** Let  $A \subseteq \mathbb{R}$ , let  $f : A \rightarrow \mathbb{R}$  and let  $c \in A$ .

- We say that  $f$  is **continuous at**  $c$  if, given any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for all  $x \in A$  satisfying  $|x - c| < \delta$ , then  $|f(x) - f(c)| < \varepsilon$ .
- Let  $B \subseteq A$ . We say that  $f$  is **continuous on**  $B$  if  $f$  is continuous at every point of  $B$ .

*Remarks.* (1) We do not assume that  $c$  is a cluster point of  $A$ .

Case 1: If  $c \in A$  is a cluster point of  $A$ , then  $f$  is continuous at  $c \iff \lim_{x \rightarrow c} f = f(c)$ .

Case 2: If  $c \in A$  is not a cluster point of  $A$ , then  $f$  is automatically continuous at  $c$ .

(2) “ $f$  is continuous on  $B$ ” and “ $f|_B$  is continuous” are different.

**Example 1.** (a) The function  $g(x) := \sin(1/x)$  for  $x \neq 0$  does not have a limit at  $x = 0$ . Thus there is no value that we can assign at  $x = 0$  to obtain a continuous extension of  $g$  at  $x = 0$ .

(b) Let  $f(x) := x \sin(1/x)$  for  $x \neq 0$ . If we define  $F : \mathbb{R} \rightarrow \mathbb{R}$  by

$$F(x) := \begin{cases} 0 & \text{for } x = 0, \\ x \sin(1/x) & \text{for } x \neq 0, \end{cases}$$

then  $F$  is continuous at  $x = 0$ .

**Example 2.** Show that the sine function is continuous on  $\mathbb{R}$ .

Suppose  $A \subseteq \mathbb{R}$ ,  $f : A \rightarrow \mathbb{R}$  and  $c \in A$ .

**Sequential Criterion for Continuity.**  $f$  is continuous at  $c$  if and only if for every sequence  $(x_n)$  in  $A$  that converges to  $c$ , the sequence  $(f(x_n))$  converges to  $f(c)$ .

**Discontinuity Criterion.**  $f$  is discontinuous at  $c$  if and only if there is a sequence  $(x_n)$  in  $A$  that converges to  $c$  but the sequence  $(f(x_n))$  does not converge to  $f(c)$ .

**Example 3.** Determine the points of continuity of the function  $f(x) := [1/x]$ ,  $x \neq 0$ . Here  $[\cdot]$  is the greatest integer function defined by

$$[x] := \sup\{n \in \mathbb{Z} : n \leq x\}.$$

**Solution.** First we show that  $f$  is continuous at each  $1/m$ ,  $m \in \mathbb{Z} \setminus \{0\}$ . Let  $x_n = (m - \frac{1}{2n})^{-1}$  for  $n \geq 1$ . Then  $\lim(x_n) = 1/m$ . However,  $f(x_n) = [m - \frac{1}{2n}] = m - 1$ , so that

$$\lim f(x_n) = m - 1 \neq m = f(1/m).$$

By discontinuity criterion,  $f$  is discontinuous at  $1/m$ .

Next we show that  $f$  is continuous at each  $c \in \mathbb{R} \setminus (\{0\} \cup \{1/m : m \in \mathbb{Z} \setminus \{0\}\})$ . Observe that,  $\delta := \min\{1/c - [1/c], [1/c] + 1 - 1/c\}/2$  satisfies

$$\left| \frac{1}{x} - \frac{1}{c} \right| < \delta \implies \left[ \frac{1}{x} \right] = \left[ \frac{1}{c} \right].$$

Let  $\varepsilon > 0$  be given. Take  $\delta' := \min\{|c|/2, \delta|c|^2/2\}$ . If  $x \in \mathbb{R} \setminus \{0\}$  and  $|x - c| < \delta'$ , then  $|x| > |c|/2$ , and

$$\left| \frac{1}{x} - \frac{1}{c} \right| = \frac{|x - c|}{|x||c|} < \frac{2}{|c|^2}|x - c| < \frac{2\delta'}{|c|^2} \leq \delta,$$

so that

$$|f(x) - f(c)| = \left| \left[ \frac{1}{x} \right] - \left[ \frac{1}{c} \right] \right| = 0 < \varepsilon.$$

Hence  $f$  is continuous at  $c$ . ◀

## Classwork

1. Determine the points of continuity of the function  $g(x) := x[x]$ .
2. Give an example for each of the following:
  - (a)  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous everywhere except at one point.
  - (b)  $f : \mathbb{R} \rightarrow \mathbb{R}$  is discontinuous everywhere.
  - (c)  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous exactly at one point.
  - (d)  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous on  $\mathbb{R} \setminus \mathbb{Q}$  but discontinuous on  $\mathbb{Q}$ .