

Fourier Analysis

Note Title

2/16/2020

Review: Good kernels and the Convergence Theorem

Def. A sequence $(K_n)_{n=1}^{\infty} = \mathcal{R}[-\pi, \pi]$ is said to be a good kernel if

$$(1) \frac{1}{2\pi} \int_{-\pi}^{\pi} K_n(x) dx = 1 \text{ for all } n \in \mathbb{N}$$

$$(2) \int_{-\pi}^{\pi} |K_n(x)| dx \leq M \text{ for all } n \in \mathbb{N},$$

where M is a constant

$$(3) \forall \delta > 0,$$

$$\int_{\delta < |x| \leq \pi} |K_n(x)| dx \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Similarly, we can define "good kernel" for a family of integrable functions

$$(K_t)_{t \in (a, b)}$$

on the circle as $t \rightarrow t_0$

Thm (Convergence Thm)

Let $(K_t(x))_{t \in (a,b)}$ be a good kernel on the circle as $t \rightarrow t_0$. Let $f \in \mathcal{R}[-\pi, \pi]$.

Then

① $f * K_t(x) \rightarrow f(x)$ as $t \rightarrow t_0$
if f is continuous at x .

② If f is continuous on the circle,
then

$f * K_t(x) \Rightarrow f(x)$ as $t \rightarrow t_0$
on the circle.

Examples of good kernels :

Fejér kernel and Poisson kernel

1. Fejér kernel

Let $D_N(x) = \sum_{|n| \leq N} e^{inx}$ denote the N -th Dirichlet kernel.

The N -th Fejér kernel :

$$\begin{aligned} F_N(x) &= \frac{D_0(x) + \dots + D_{N-1}(x)}{N} \\ &= \sum_{n=-N}^N \left(1 - \frac{|n|}{N}\right) e^{inx} \\ &= \frac{\sin^2\left(\frac{N}{2}x\right)}{N \sin^2\left(\frac{x}{2}\right)}. \end{aligned}$$

As was proved, $(F_N)_{N=1}^{\infty}$ is a good kernel as $N \rightarrow \infty$.

Corollary (Fejér's Thm)

Let $f \in \mathcal{R}[-\pi, \pi]$. Then

(1) $f * F_N(x) \rightarrow f(x)$ as $N \rightarrow \infty$
if f is cts at x .

(2) If f is cts on the circle, then
 $f * F_N(x) \Rightarrow f(x)$ on the circle.

Remark: Suppose $f \sim \sum_{n=-\infty}^{\infty} a_n e^{inx}$,

then $f * K_N(x) = \sum_{n=-N}^N \left(1 - \frac{|n|}{N}\right) a_n e^{inx}$.

Write also

$$\sigma_N(f)(x) = f * K_N(x)$$

and call it

"the N -th Cesaro mean of the
Fourier series of f ."

Corollary. Continuous functions on the circle can be unif. approximated by trigonometric polynomials.

That is, if f is cts on $[-\pi, \pi]$ and $f(\pi) = f(-\pi)$, then $\forall \varepsilon > 0$, \exists a trigonometric polynomial P such that

$$|f(x) - P(x)| < \varepsilon \quad \text{for all } x \in [-\pi, \pi]$$

Pf. Notice that $\sigma_N(f)(x) \Rightarrow f(x)$ and $\sigma_N(f)$ are trigonometric polynomials. \square

Corollary. Let f be a continuous function on the circle so that

$\hat{f}(n) = 0$ for all $n \in \mathbb{Z}$.
Then $f \equiv 0$.

Pf. Since $\hat{f}(n) \equiv 0$,

$$\sigma_N(f) \equiv 0.$$

However,

$\sigma_N(f) \Rightarrow f$ as $N \rightarrow \infty$
So $f \equiv 0$.

2. Poisson kernel on the circle:

$(P_r(x))_{0 \leq r < 1}$ where

$$P_r(x) = \frac{1-r^2}{1-2r \cos x + r^2}$$
$$= \sum_{n=-\infty}^{\infty} r^{|n|} e^{inx}.$$

As was proved, $(P_r)_{0 \leq r < 1}$ is a good kernel
as $r \rightarrow 1$.

Corollary. Let $f \in \mathcal{R}[-\pi, \pi]$. Then

$$\lim_{r \rightarrow 1} f * P_r(x) = f(x) \text{ if } f \text{ is cts at } x.$$

Moreover

$f * P_r(x) \Rightarrow f(x)$ on the circle
if f is cts on the circle.

Remark: If $f(x) \sim \sum_{n=-\infty}^{\infty} a_n e^{inx}$,

then

$$f * P_r(x) = \sum_{n=-\infty}^{\infty} r^{|n|} a_n e^{inx}.$$

We also write

$$A_r(f)(x) = f * P_r(x)$$

and call it the Abel mean of the
Fourier series of f .