

HW 7

- Using the Structure Theorem for Open Sets (and check by epsilon-delta terminology) show that each continuous function f on a closed set F in \mathbb{R} can be continuously extended to be on the whole of \mathbb{R} (this is known as the Tietze extension Theorem).
- Let f be a measurable real-valued function on a set E of finite measure. Show that there exists a sequence of continuous functions convergent to f almost everywhere on E . Hence, for any $r > 0$, there exist a closed set F contained in E with $m(E \setminus F) < r$ such that the above convergence is uniform on F and the restriction of f to F is continuous.
- Let f be a measurable real-valued function on a measurable set E of possibly infinite measure, and let $r > 0$. Apply Q2 to get a corresponding closed set F_n contained in the intersection of E with $(n, n+1]$ for each integer n . Show that the union F of F_n is closed and that the restriction of f to F is continuous. Moreover we can arrange in such a way that $m(E \setminus F) < r$.
- Let f be a non-negative extended real function on a measurable set E . Show that the sequence (f_n) of simple functions monotonically increases and converges point-wisely to f , where

$$f_n := \sum_{k=1}^n \frac{k-1}{2^n} \chi_{B_{n,k}} + n \chi_{A_n}$$

$$A_n = \{x \in E \cap [-n, n] : n \leq f(x)\}$$

$$B_{n,k} = \{x \in E \cap [-n, n] : \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n}\}, \quad k=1, \dots, n \cdot 2^n$$

(This result is also known as Littlewood's 2nd principle = one of the versions)

- Let $m(E) < +\infty$, and $\mathcal{L}(E)$ consist of all measurable functions f on E such that $\int_E |f| < +\infty$. Let

$f_n, g_n, f, g \in \mathcal{L}(E)$ be such that $f_n \rightarrow f, g_n \rightarrow g$

and $f_n \leq g_n$ $\forall n$ (all are pointwise or a.e. on E).

Suppose further that $\lim_n \int_E g_n = \int_E g$. Show that

$$\lim_n \int_E |f_n - f| = 0.$$

- Let $f_n, f \in \mathcal{L}(E)$ and $f_n \rightarrow f$ a.e. on E . Suppose

$$\lim_n \int_E |f_n| = \int_E |f|. \quad \text{Show that } \lim_n \int_E f_n = \int_E f.$$