

NOTE OF ELEMENTARY ANALYSIS II

CHI-WAI LEUNG

1. RIEMANN INTEGRALS

Notation 1.1. .

- (i) : All functions f, g, h, \dots are bounded real valued functions defined on $[a, b]$. And $m \leq f \leq M$.
- (ii) : $\mathcal{P} : a = x_0 < x_1 < \dots < x_n = b$ denotes a partition on $[a, b]$; $\Delta x_i = x_i - x_{i-1}$ and $\|\mathcal{P}\| = \max \Delta x_i$.
- (iii) : $M_i(f, \mathcal{P}) := \sup\{f(x) : x \in [x_{i-1}, x_i]\}$; $m_i(f, \mathcal{P}) := \inf\{f(x) : x \in [x_{i-1}, x_i]\}$. And $\omega_i(f, \mathcal{P}) = M_i(f, \mathcal{P}) - m_i(f, \mathcal{P})$.
- (iv) : $U(f, \mathcal{P}) := \sum M_i(f, \mathcal{P})\Delta x_i$; $L(f, \mathcal{P}) := \sum m_i(f, \mathcal{P})\Delta x_i$.
- (v) : $\mathcal{R}(f, \mathcal{P}, \{\xi_i\}) := \sum f(\xi_i)\Delta x_i$, where $\xi_i \in [x_{i-1}, x_i]$.
- (vi) : $\mathcal{R}[a, b]$ is the class of all Riemann integral functions on $[a, b]$.

Definition 1.2. We say that the Riemann sum $\mathcal{R}(f, \mathcal{P}, \{\xi_i\})$ converges to a number A as $\|\mathcal{P}\| \rightarrow 0$ if for any $\varepsilon > 0$, there is $\delta > 0$ such that

$$|A - \mathcal{R}(f, \mathcal{P}, \{\xi_i\})| < \varepsilon$$

for any $\xi_i \in [x_{i-1}, x_i]$ whenever $\|\mathcal{P}\| < \delta$.

Theorem 1.3. $f \in \mathcal{R}[a, b]$ if and only if for any $\varepsilon > 0$, there is a partition \mathcal{P} such that $U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$.

Lemma 1.4. $f \in \mathcal{R}[a, b]$ if and only if for any $\varepsilon > 0$, there is $\delta > 0$ such that $U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$ whenever $\|\mathcal{P}\| < \delta$.

Proof. The converse follows from Theorem 1.3.

Assume that f is integrable over $[a, b]$. Let $\varepsilon > 0$. Then there is a partition $\mathcal{Q} : a = y_0 < \dots < y_l = b$ on $[a, b]$ such that $U(f, \mathcal{Q}) - L(f, \mathcal{Q}) < \varepsilon$. Now take $0 < \delta < \varepsilon/l$. Suppose that $\mathcal{P} : a = x_0 < \dots < x_n = b$ with $\|\mathcal{P}\| < \delta$. Then we have

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) = I + II$$

where

$$I = \sum_{i: \mathcal{Q} \cap (x_{i-1}, x_i) = \emptyset} \omega_i(f, \mathcal{P})\Delta x_i;$$

and

$$II = \sum_{i: \mathcal{Q} \cap (x_{i-1}, x_i) \neq \emptyset} \omega_i(f, \mathcal{P})\Delta x_i$$

Notice that we have

$$I \leq U(f, \mathcal{Q}) - L(f, \mathcal{Q}) < \varepsilon$$

Date: March 2, 2016.

and

$$II \leq (M - m) \sum_{i: Q \cap (x_{i-1}, x_i) \neq \emptyset} \Delta x_i \leq (M - m) \cdot l \cdot \frac{\varepsilon}{l} = (M - m)\varepsilon.$$

The proof is finished. \square

Theorem 1.5. $f \in \mathcal{R}[a, b]$ if and only if the Riemann sum $\mathcal{R}(f, \mathcal{P}, \{\xi_i\})$ is convergent. In this case, $\mathcal{R}(f, \mathcal{P}, \{\xi_i\})$ converges to $\int_a^b f(x)dx$ as $\|\mathcal{P}\| \rightarrow 0$.

Proof. For the proof (\Rightarrow): we first note that we always have

$$L(f, \mathcal{P}) \leq \mathcal{R}(f, \mathcal{P}, \{\xi_i\}) \leq U(f, \mathcal{P})$$

and

$$L(f, \mathcal{P}) \leq \int_a^b f(x)dx \leq U(f, \mathcal{P})$$

for any $\xi_i \in [x_{i-1}, x_i]$ and for all partition \mathcal{P} .

Now let $\varepsilon > 0$. Lemma 1.4 gives $\delta > 0$ such that $U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$ as $\|\mathcal{P}\| < \delta$. Then we have

$$\left| \int_a^b f(x)dx - \mathcal{R}(f, \mathcal{P}, \{\xi_i\}) \right| < \varepsilon$$

as $\|\mathcal{P}\| < \delta$. The necessary part is proved and $\mathcal{R}(f, \mathcal{P}, \{\xi_i\})$ converges to $\int_a^b f(x)dx$.

For (\Leftarrow): there exists a number A such that for any $\varepsilon > 0$, there is $\delta > 0$, we have

$$A - \varepsilon < \mathcal{R}(f, \mathcal{P}, \{\xi_i\}) < A + \varepsilon$$

for any partition \mathcal{P} with $\|\mathcal{P}\| < \delta$ and $\xi_i \in [x_{i-1}, x_i]$.

Now fix a partition \mathcal{P} with $\|\mathcal{P}\| < \delta$. Then for each $[x_{i-1}, x_i]$, choose $\xi_i \in [x_{i-1}, x_i]$ such that $M_i(f, \mathcal{P}) - \varepsilon \leq f(\xi_i)$. This implies that we have

$$U(f, \mathcal{P}) - \varepsilon(b - a) \leq \mathcal{R}(f, \mathcal{P}, \{\xi_i\}) < A + \varepsilon.$$

So we have shown that for any $\varepsilon > 0$, there is a partition \mathcal{P} such that

$$(1.1) \quad \int_a^b f(x)dx \leq U(f, \mathcal{P}) \leq A + \varepsilon(1 + b - a).$$

By considering $-f$, note that the Riemann sum of $-f$ will converge to $-A$. The inequality 1.1 will imply that for any $\varepsilon > 0$, there is a partition \mathcal{P} such that

$$A - \varepsilon(1 + b - a) \leq \int_a^b f(x)dx \leq \int_a^b f(x)dx \leq A + \varepsilon(1 + b - a).$$

The proof is finished. \square

Theorem 1.6. Let $f \in \mathcal{R}[c, d]$ and let $\phi : [a, b] \rightarrow [c, d]$ be a strictly increasing C^1 function with $f(a) = c$ and $f(b) = d$.

Then $f \circ \phi \in \mathcal{R}[a, b]$, moreover, we have

$$\int_c^d f(x)dx = \int_a^b f(\phi(t))\phi'(t)dt.$$

Proof. Let $A = \int_c^d f(x)dx$. By Theorem 1.5, we need to show that for all $\varepsilon > 0$, there is $\delta > 0$ such that

$$|A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| < \varepsilon$$

for all $\xi_k \in [t_{k-1}, t_k]$ whenever $\mathcal{Q} : a = t_0 < \dots < t_m = b$ with $\|\mathcal{Q}\| < \delta$.

Now let $\varepsilon > 0$. Then by Lemma 1.4 and Theorem 1.5, there is $\delta_1 > 0$ such that

$$(1.2) \quad |A - \sum f(\eta_k)\Delta x_k| < \varepsilon$$

and

$$(1.3) \quad \sum \omega_k(f, \mathcal{P})\Delta x_k < \varepsilon$$

for all $\eta_k \in [x_{k-1}, x_k]$ whenever $\mathcal{P} : c = x_0 < \dots < x_m = d$ with $\|\mathcal{P}\| < \delta_1$.

Now put $x = \phi(t)$ for $t \in [a, b]$.

Now since ϕ and ϕ' are continuous on $[a, b]$, there is $\delta > 0$ such that $|\phi(t) - \phi(t')| < \delta_1$ and $|\phi'(t) - \phi'(t')| < \varepsilon$ for all t, t' in $[a, b]$ with $|t - t'| < \delta$.

Now let $\mathcal{Q} : a = t_0 < \dots < t_m = b$ with $\|\mathcal{Q}\| < \delta$. If we put $x_k = \phi(t_k)$, then $\mathcal{P} : c = x_0 < \dots < x_m = d$ is a partition on $[c, d]$ with $\|\mathcal{P}\| < \delta_1$ because ϕ is strictly increasing.

Note that the Mean Value Theorem implies that for each $[t_{k-1}, t_k]$, there is $\xi_k^* \in (t_{k-1}, t_k)$ such that

$$\Delta x_k = \phi(t_k) - \phi(t_{k-1}) = \phi'(\xi_k^*)\Delta t_k.$$

This yields that

$$(1.4) \quad |\Delta x_k - \phi'(\xi_k)\Delta t_k| < \varepsilon\Delta t_k$$

for any $\xi_k \in [t_{k-1}, t_k]$ for all $k = 1, \dots, m$ because of the choice of δ .

Now for any $\xi_k \in [t_{k-1}, t_k]$, we have

$$(1.5) \quad \begin{aligned} |A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| &\leq |A - \sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k| \\ &+ |\sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k - \sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k| \\ &+ |\sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| \end{aligned}$$

Notice that inequality 1.2 implies that

$$|A - \sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k| = |A - \sum f(\phi(\xi_k^*))\Delta x_k| < \varepsilon.$$

Also, since we have $|\phi'(\xi_k^*) - \phi'(\xi_k)| < \varepsilon$ for all $k = 1, \dots, m$, we have

$$|\sum f(\phi(\xi_k^*))\phi'(\xi_k^*)\Delta t_k - \sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k| \leq M(b-a)\varepsilon$$

where $|f(x)| \leq M$ for all $x \in [c, d]$.

On the other hand, by using inequality 1.4 we have

$$|\phi'(\xi_k)\Delta t_k| \leq \Delta x_k + \varepsilon\Delta t_k$$

for all k . This, together with inequality 1.3 imply that

$$\begin{aligned}
& \left| \sum f(\phi(\xi_k^*))\phi'(\xi_k)\Delta t_k - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k \right| \\
& \leq \sum \omega_k(f, \mathcal{P})|\phi'(\xi_k)\Delta t_k| \quad (\because \phi(\xi_k^*), \phi(\xi_k) \in [x_{k-1}, x_k]) \\
& \leq \sum \omega_k(f, \mathcal{P})(\Delta x_k + \varepsilon\Delta t_k) \\
& \leq \varepsilon + 2M(b-a)\varepsilon.
\end{aligned}$$

Finally by inequality 1.5, we have

$$|A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k| \leq \varepsilon + M(b-a)\varepsilon + \varepsilon + 2M(b-a)\varepsilon.$$

The proof is finished. □

REFERENCES

(Chi-Wai Leung) DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

E-mail address: cwleung@math.cuhk.edu.hk