

MATH 4030 Differential Geometry
Homework 2

due 20/9/2015 (Tue) at 5PM

Problems

(to be handed in)

Unless otherwise stated, we use I, J to denote connected open intervals in \mathbb{R} .

1. Consider the *logarithmic spiral* $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$ given by

$$\alpha(t) = (ae^{bt} \cos t, ae^{bt} \sin t)$$

with $a > 0, b < 0$. Compute the arc length function $S : \mathbb{R} \rightarrow \mathbb{R}$ from $t_0 \in \mathbb{R}$. Reparametrize this curve by arc length and study its trace.

2. Let $\alpha : I \rightarrow \mathbb{R}^2$ be a regular curve (not necessarily parametrized by arc length). Prove that

$$k_\alpha(t) = \frac{1}{|\alpha'(t)|^3} \det(\alpha'(t), \alpha''(t)).$$

3. Consider the *catenary* given by $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$ where $\alpha(t) = (t, \cosh t)$, compute the curvature $k_\alpha(t)$.

4. Let $\alpha : T \rightarrow \mathbb{R}^2$ be a curve p.b.a.l. and $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a rigid motion. Show that $\beta = \varphi \circ \alpha : I \rightarrow \mathbb{R}^2$ is a curve p.b.a.l. and that

$$k_\beta(s) = \begin{cases} k_\alpha(s) & \text{if } \varphi \text{ is orientation-preserving} \\ -k_\alpha(s) & \text{if } \varphi \text{ is orientation-reversing} \end{cases}$$

5. Let $\alpha : I = (-a, a) \rightarrow \mathbb{R}^2$ be a curve p.b.a.l. for some $a > 0$. Suppose that $k_\alpha(s) = k_\alpha(-s)$ for each $s \in (-a, a)$. Prove that the trace of α is symmetric relative to the normal line of α at $s = 0$.

Suggested Exercises

(no need to hand in)

1. Let $\alpha : I \rightarrow \mathbb{R}^2$ be a plane curve with $[a, b] \subset I$. Suppose $P = \{a = t_0 < t_1 < t_2 < \dots < t_n = b\}$ is a partition of $[a, b]$. Define

$$L_a^b(\alpha, P) = \sum_{i=1}^n |\alpha(t_i) - \alpha(t_{i-1})|$$

to be the length of the polygonal line. Prove that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$\left| L_a^b(\alpha, P) - \int_a^b |\alpha'(t)| dt \right| < \epsilon$$


whenever $|P| := \max_{1 \leq i \leq n} |t_i - t_{i-1}| < \delta$. *Hint: use mean value theorems.* Moreover, show that

$$L_a^b(\alpha) = \sup\{L_a^b(\alpha, P) \mid P \text{ is a partition of } [a, b]\}.$$


2. Let $\alpha : I \rightarrow \mathbb{R}^3$ be a regular curve. Show that $|\alpha'(t)|$ is a nonzero constant if and only if $\langle \alpha(t), \alpha'(t) \rangle = 0$ for all $t \in I$.
3. Show that the tangent lines to the regular curve $\alpha(t) = (3t, 3t^2, 2t^3)$ make a constant angle with the line $\ell = \{y = 0, z = x\}$.
4. A circular disk of radius 1 in the xy -plane rolls without slipping along the x -axis. The figure described by a point of the circumference of the disk is called a *cycloid* (see Figure 1(a)). Find a curve $\alpha : I \rightarrow \mathbb{R}^2$ whose trace is the cycloid. Determine the value of $t \in I$ at which α is not regular. Moreover, compute the arc length of the cycloid corresponding to a complete rotation of the disk.
5. Let $\alpha : (0, \pi) \rightarrow \mathbb{R}^2$ be given by

$$\alpha(t) = (\sin t, \cos t + \log \tan \frac{t}{2}),$$

where t is the angle the y -axis make with the vector $\alpha'(t)$. The trace of α is called the *tractrix* (see Figure 1(b)). Show that α is regular except at $t = \pi/2$. Moreover, prove that the length of the segment of the tangent of the tractrix between the point of tangency and the y -axis is constantly equal to 1.



(a) The cycloid



(b) The tractrix

Figure 1

6. Let $\alpha : I \rightarrow \mathbb{R}^2$ be a regular curve such that all the normal lines pass through a fixed point $p \in \mathbb{R}^2$. Prove that the trace $\alpha(I)$ is contained in a circle of some radius $r > 0$ centered at p .
7. Let $\alpha : I \rightarrow \mathbb{R}^2$ be a regular curve such that all the tangent lines pass through a fixed point $p \in \mathbb{R}^2$. Prove that the trace $\alpha(I)$ is contained in a straight line passing through p . What if α is not regular?