

## Math 2230A, Complex Variables with Applications

Use residues to derive the integration formulas in Question 1 and Question 2.

1.


$$\int_0^{\infty} \frac{dx}{(x^2 + 1)^2} = \frac{\pi}{4}$$

2.

$$\int_0^{\infty} \frac{x^2 dx}{x^6 + 1} = \frac{\pi}{6}$$

3. Use a residue and a contour shown in Fig.95, where  $R > 1$ , to establish the integration formula

$$\int_0^{\infty} \frac{dx}{x^3 + 1} = \frac{2\pi}{3\sqrt{3}}$$



**FIGURE 95**

4. Use residues to derive the integration formula

$$\int_0^{\infty} \frac{\cos ax}{(x^2 + b^2)^2} dx = \frac{\pi}{4b^3} (1 + ab)e^{-ab} \quad (a > 0, b > 0).$$

5. Use residues to derive the integration formula

$$\int_{-\infty}^{\infty} \frac{x^3 \sin ax}{x^4 + 4} dx = \pi e^{-a} \cos a \quad (a > 0).$$

6. Use residues to find the Cauchy principal values of the improper integrals

$$\int_{-\infty}^{\infty} \frac{\sin x dx}{x^2 + 4x + 5}$$

7. Follow the steps below to evaluate the *Fresnel integrals*, which are important in diffraction theory:

$$\int_0^\infty \cos(x^2) dx = \int_0^\infty \sin(x^2) dx = \frac{1}{2} \sqrt{\frac{\pi}{2}}$$

- (a) By integrating the function  $\exp(iz^2)$  around the positively oriented boundary of the sector  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \frac{\pi}{4}$  (Fig. 99) and appealing to the Cauchy-Goursat theorem, show that

$$\int_0^R \cos(x^2) dx = \frac{1}{\sqrt{2}} \int_0^R e^{-r^2} dr - \operatorname{Re} \int_{C_R} e^{iz^2} dz$$

and

$$\int_0^R \sin(x^2) dx = \frac{1}{\sqrt{2}} \int_0^R e^{-r^2} dr - \operatorname{Im} \int_{C_R} e^{iz^2} dz,$$




FIGURE 99

where  $C_R$  is the arc  $z = Re^{i\theta}$  ( $0 \leq \theta \leq \frac{\pi}{4}$ ).

- (b) Show that the value of the integral along the arc  $C_R$  in part (a) tends to zero as  $R$  tends to infinity by obtaining the inequality

$$\left| \int_{C_R} e^{iz^2} dz \right| \leq \frac{R}{2} \int_0^{\pi/2} e^{-R^2 \sin \phi} d\phi$$

and then referring to the form (2), Sec. 81, of Jordan's inequality.

- (c) Use the results in part (a) and (b), together with the known formula

$$\int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2},$$

to complete the exercise.

8. Use the function  $f(z) = (e^{iaz} - e^{ibz})/z^2$  and the indented contour in Fig.108(Sec.89) to derive the integration formula

$$\int_0^\infty \frac{\cos(ax) - \cos(bx)}{x^2} dx = \frac{\pi}{2}(b - a) \quad (a \geq 0, b \geq 0)$$

Then with the aid of the trigonometric identity  $1 - \cos(2x) = 2 \sin^2 x$ , point out how it follows that

$$\int_0^\infty \frac{\sin^2 x}{x^2} dx = \frac{\pi}{2}.$$

9. Derive the integration formula

$$\int_0^\infty \frac{dx}{\sqrt{x}(x^2 + 1)} = \frac{\pi}{\sqrt{2}}$$

by integrating the function

$$f(z) = \frac{z^{-1/2}}{z^2 + 1} = \frac{e^{(-1/2)\log z}}{z^2 + 1} \quad (|z| > 0, -\frac{\pi}{2} < \arg z < \frac{3\pi}{2})$$

over the indented contour appearing in Fig. 109 (Sec. 90).

10. Use residues to establish the following integration formula:

(a)

$$\int_0^{2\pi} \frac{d\theta}{5 + 4 \sin \theta} = \frac{2\pi}{3}.$$

(b)

$$\int_{-\pi}^{\pi} \frac{d\theta}{1 + \sin^2 \theta} = \sqrt{2}\pi.$$

(c)

$$\int_0^{2\pi} \frac{\cos^2 3\theta d\theta}{5 - 4 \cos 2\theta} = \frac{3\pi}{8}.$$

(d)

$$\int_0^{2\pi} \frac{d\theta}{1 + a \cos \theta} = \frac{2\pi}{\sqrt{1 - a^2}} \quad (-1 < a < 1).$$