

THE CHINESE UNIVERSITY OF HONG KONG  
Department of Mathematics  
MATH2230 Tutorial 2  
(Prepared by Tai Ho Man)

## 0.1 Polynomial and Rational function

The domain of definition (or simply the domain) of a function is the set of input for which the function value is defined.

**Definition 1.** We call the function in the form of

$$P_n(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n \text{ for } n = 0, 1, 2, \dots \text{ and } a_n \neq 0$$

to be the polynomial of degree  $n$ .

Remark: The domain of definition of polynomial is clearly  $\mathbb{C}$ .

Remark: The condition of  $a_n \neq 0$  is meaningful. Otherwise, the degree of the polynomial is no longer  $n$ , it could be smaller than  $n$ .

**Definition 2.** Given two polynomials  $P_n(z)$  and  $Q_m(z)$ , the function  $R(z) = \frac{P_n(z)}{Q_m(z)}$  is called the rational function.

Remark: The domain of definition of rational function is clearly  $\mathbb{C} \setminus \{z_1, z_2, \dots, z_m\}$  where  $z_1, z_2, \dots, z_m$  are the roots of  $Q_m(z) = 0$ .

## 0.2 Trigonometric function

$$\begin{aligned} \sin z &= \frac{e^{iz} - e^{-iz}}{2i} & \cos z &= \frac{e^{iz} + e^{-iz}}{2} \\ \sinh z &= -i \sin(iz) = \frac{e^z - e^{-z}}{2} & \cosh z &= \cos(iz) = \frac{e^z + e^{-z}}{2} \end{aligned}$$

Remark: The domain of definition of these trigonometric functions are clearly  $\mathbb{C}$  since that of exponential function is also  $\mathbb{C}$ .

## 0.3 Logarithmic function

If the logarithmic is the "inverse" of exponential function, we let  $z_0 = re^{i\theta} \neq 0$  for  $-\pi < \theta \leq \pi$  ( $\theta = \text{Arg}(z)$ ) and we expect to have

$$\log(z_0) = \log(re^{i\theta}) = \log(r) + i\theta$$

However,  $z_0 = re^{i\theta} = re^{i\theta+2k\pi i}$  for any integers  $k$ , hence we have

$$\log(z_0) = \log(re^{i\theta+2k\pi i}) = \log(r) + i\theta + 2k\pi i$$

It shows that the logarithmic defined in this way is not a function. We see that  $\log(z_0)$  represent a set

$$\log(z_0) := \{ \log(r) + i\theta + 2k\pi i \mid k \in \mathbb{Z} \}$$

or we will call  $\log$  is a multiple-valued function. It is quite similar to the definition of argument of a complex number. Therefore we should define logarithmic in a unique way.

**Definition 3.** The principal value of  $\log(z)$  equals to

$$\text{Log}(z) = \log |z| + i\text{Arg}(z)$$

where  $-\pi < \text{Arg}(z) \leq \pi$ .

Remark 1 : Since it is reasonable to define the range of the angle of  $z_0$  in another way, say,  $z_0 = re^{i\theta} \neq 0$  for  $a < \theta \leq 2\pi + a$  for any real number  $a$ . Such a choice of range of the angle of  $z$  is called branch. And we can define another single-value function for  $\log$  by  $\log(r) + i\theta$  with  $a < \theta \leq 2\pi + a$ . (The word "principal" in definition 3 means that  $a = -\pi$ . We would not call the single-valued  $\log$  to be principal if  $a \neq -\pi$ .) And the range  $-\pi < \theta \leq \pi$  is called principal branch.

Remark 2 : The domain of definition of  $\text{Log}(z)$  is  $\mathbb{C} \setminus \{0\}$  because of  $\log |z|$ .

Remark 3 : Although  $\text{Log}(z)$  can be defined on the ray  $\theta = a$ ,  $\text{Log}(z)$  is not continuous there (not analytic).

Remark 4 :  $\text{Log}(z)$  is analytic in the domain  $r > 0$  and  $-\pi < \text{Arg}(z) < \pi$  (or other branch).

## 0.4 Power function

**Definition 4.** Let  $z \neq 0$  and  $c \in \mathbb{C}$ , the power is defined as

$$z^c = e^{c \text{Log}(z)}$$

Clearly it can be defined for other branch.

Remark : The domain of definition of power function is again  $\mathbb{C} \setminus \{0\}$ .

Remark : Some operations which is true in real number turn out is false in complex number:

(a)  $z^{c_1} z^{c_2} = z^{c_1+c_2}$ ; (b)  $(z^{c_1})^{c_2} = z^{c_1 c_2}$  (c)  $(zw)^c = z^c w^c$

## 0.5 Continuity of a Function

**Definition 5.** Let  $\Omega$  be open subset of  $\mathbb{C}$  and  $f : \Omega \rightarrow \mathbb{C}$ . Let  $z_0 \in \Omega$ , we say  $\lim_{z \rightarrow z_0} f(z) = c$  if for all  $\varepsilon > 0$  there is a  $\delta > 0$  such that if  $|z - z_0| < \delta$ , then  $|f(z) - c| < \varepsilon$ .

**Definition 6.** Let  $\Omega$  be open subset of  $\mathbb{C}$  and  $f = f_1 + if_2 : \Omega \rightarrow \mathbb{C}$ . Let  $z_0 \in \Omega$ , then  $f$  is continuous at  $z_0$  if and only if  $f_1$  and  $f_2$  are continuous at  $z_0$ . In other words,  $\lim_{z \rightarrow z_0} f(z) = f(z_0)$  if and only if  $\lim_{z \rightarrow z_0} f_1(z) = f_1(z_0)$  and  $\lim_{z \rightarrow z_0} f_2(z) = f_2(z_0)$ .

## 0.6 Exercise

1. Compute the value of  $\log(-1 + \sqrt{3}i)$  with branch  $-\pi < \text{Arg}(z) \leq \pi$ .
2. Find the domain of  $f(z) = \text{Log}(z - i)$ .
3. Find the principal values of  $(1 + i)^i$ .
4. Describe the image under  $f = e^z$  of the following sets:
  - (a) The set of  $z = x + yi$  such that  $x \leq 1$  and  $0 \leq y \leq \pi$ .
  - (b) The set of  $z = x + yi$  such that  $0 \leq y \leq \pi$ .