

MATH4060 Exercise 2

Deadline: October 6, 2015.

The questions are from Stein and Shakarchi, *Complex Analysis*, unless otherwise stated.

Chapter 3. Exercise 12.

Chapter 4. Exercise 1, 2, 3, 6, 7.

Chapter 5. Exercise 2, 3, 4, 5, 6.

Additional Exercise. Show that if $0 < a < 1$, then

$$\int_0^{\infty} \frac{v^{a-1}}{1+v} dv = \frac{\pi}{\sin \pi a}.$$

(Hint: Consider, for a small $\delta > 0$, the contour $\gamma_1 + \gamma_2 - \gamma_3 - \gamma_4$, where

$$\gamma_1 = \{t + i\delta : \varepsilon \leq t \leq R\},$$

$$\gamma_2 = \{(R^2 + \delta^2)^{1/2} e^{i\theta} : \arctan(\delta/R) \leq \theta \leq 2\pi - \arctan(\delta/R)\},$$

$$\gamma_3 = \{t - i\delta : \varepsilon \leq t \leq R\},$$

$$\gamma_4 = \{(\varepsilon^2 + \delta^2)^{1/2} e^{i\theta} : \arctan(\delta/\varepsilon) \leq \theta \leq 2\pi - \arctan(\delta/\varepsilon)\},$$

and consider a suitable branch of $v \mapsto v^{a-1}$ for v away from the non-negative real axis.)