

1. Let $f_1, f_2, f_3, f_4, f_5, f_6, f_7, f_8 : \mathbb{R} \rightarrow \mathbb{R}$ be the functions defined by


$$f_1(x) = 0.1x^3, \quad f_2(x) = \sqrt[5]{x} - 1, \quad f_3(x) = x^5 - 2x^3 + x, \quad f_4(x) = 0.25x^2 \sin(10x),$$

$$f_5(x) = 1.3^x, \quad f_6(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}, \quad f_7(x) = \frac{1}{x^2 + 1}, \quad f_8(x) = 4^{\sin(4x)}$$


for any $x \in \mathbb{R}$.

Rough sketches of the respective graphs of the above functions:


Graph of f_1 :




Graph of f_2 :




Graph of f_3 :




Graph of f_4 :




Graph of f_5 :




Graph of f_6 :



Graph of f_7 :



Graph of f_8 :



2. Which of f_1, \dots, f_8 is/are surjective? Which not?

• f_1, f_2, f_3, f_4 are surjective.

• f_5, f_6, f_7, f_8 are not surjective.

Question. How to see the answer for such functions from \mathbb{R} to \mathbb{R} ?

Answer (b1). Inspect the graph of f_1, \dots, f_8 on the ‘coordinate plane’.

• $i = 1, 2, 3, 4$. Why surjective?

At each ‘altitude’ $b \in \mathbb{R}$, the horizontal line $y = b$ cuts the graph of f_i at least once.

Some $x_b \in \mathbb{R}$ satisfies $y = f_i(x_b)$.

• $i = 5, 6, 7, 8$. Why not surjective?

At some ‘altitude’ $b_0 \in \mathbb{R}$, the horizontal line $y = b_0$ cuts the graph of f_i nowhere.

No $x \in \mathbb{R}$ satisfies $b_0 = f_i(x)$.

Answer (b2). Re-interpret (b1) in terms of solving equations.

• $i = 1, 2, 3, 4$. Why surjective?

For each $b \in \mathbb{R}$, the equation $b = f_i(u)$ with ‘unknown’ u in \mathbb{R} has at least one solution in \mathbb{R} .

• $i = 5, 6, 7, 8$. Why not surjective?

There is some $b_0 \in \mathbb{R}$ for which the equation $b_0 = f_i(u)$ with ‘unknown’ u in \mathbb{R} has no solution in \mathbb{R} .

Answer (a). Directly verify the condition (S) or its negation respectively.

• $i = 1, 2, 3, 4$. Surjectivity? [Recall the statement (S).]

* How do we check the surjectivity of f_1 , in practice?

$$\forall y \in \mathbb{R}, \exists x \in \mathbb{R} \text{ such that } y = f_1(x).$$

Pick any $y \in \mathbb{R}$. [This y is kept fixed in the discussion below.]

[We name a candidate $x \in \mathbb{R}$ for which $y = f_1(x)$. An appropriate candidate is given by a solution of the equation $y = f_1(u)$ with unknown u in \mathbb{R} .]

Take $x = \sqrt[3]{10y}$.

• By definition, $x \in \mathbb{R}$.

• Also, $f_1(x) = 0.1x^3 = 0.1(\sqrt[3]{10y})^3 = (0.1) \cdot (10y) = y$.

It follows that f_1 is surjective. \square

* How about f_2 ? [Exercise.]

Pick any $y \in \mathbb{R}$.

Take $x = (y+1)^5$.

• By definition, $x \in \mathbb{R}$.

• Also, $f_2(x) = \sqrt[5]{x} - 1 = \sqrt[5]{(y+1)^5} - 1 = (y+1) - 1 = y$.

It follows that f_2 is surjective. \square

Roughwork: Solve $y = f_1(u)$ with unknown $u \in \mathbb{R}$.

$$y = 0.1u^3$$
$$u^3 = 10y$$
$$u = \sqrt[3]{10y}$$

Roughwork: Solve $y = f_2(u)$ with unknown $u \in \mathbb{R}$.

$$y = \sqrt[5]{u} - 1$$
$$\sqrt[5]{u} = y + 1$$
$$u = (y + 1)^5$$

Remark. Things are more difficult in practice for f_3, f_4 , when we do not make use of the calculus. (Why?)

Answer (a). Directly verify the condition (S) or its negation respectively.

- $i = 5, 6, 7, 8$. Non-surjectivity? [Recall the statement $\sim(S)$.]

* How do we check the non-surjectivity of f_8 , in practice? $\exists y_0 \in \mathbb{R}$ such that $(\forall x \in \mathbb{R}, y_0 \neq f_8(x))$.

[Name a candidate $y_0 \in \mathbb{R}$ for which $y_0 \neq f_8(x)$ for any $x \in \mathbb{R}$. We are aware that for any $x \in \mathbb{R}$, $4^{-1} \leq 4^{\sin(4x)} \leq 4$.]

Take $y_0 = 5$.

- Note that $y_0 \in \mathbb{R}$.
- Pick any $x \in \mathbb{R}$. We have $f_8(x) = 4^{\sin(4x)} \leq 4 < 5 = y_0$.

Hence $y_0 \neq f_8(x)$.

It follows that f_8 is not surjective. \square

* How about f_6 ?



Roughwork: Observe that for any $x \in \mathbb{R}$,

$$\left[|f_6(x)| = \left| \frac{e^x - e^{-x}}{e^x + e^{-x}} \right| = \frac{|e^x - e^{-x}|}{e^x + e^{-x}} \leq \frac{|e^x| + |e^{-x}|}{e^x + e^{-x}} = 1. \right]$$

Take $y_0 = 2$.

- Note that $y_0 \in \mathbb{R}$.
 - Pick any $x \in \mathbb{R}$. We have $|f_6(x)| = \left| \frac{e^x - e^{-x}}{e^x + e^{-x}} \right| \leq 1 < 2 = y_0$. Then $f_6(x) \neq y_0$.
- It follows that f_6 is not surjective. \square

* How about f_5, f_7 ? [Exercise.]



3. Which of f_1, \dots, f_8 is/are injective? Which not?

• f_1, f_2, f_5, f_6 are injective.

• f_3, f_4, f_7, f_8 are not injective.

Question. How to see which is injective and which not, for such a real-valued function of one real variable?

Answer (b1). Inspect the graph of f_1, \dots, f_8 .

• $i = 1, 2, 5, 6$. Why injective?

At each 'altitude' $b \in \mathbb{R}$, the horizontal line $y = b$ cuts the graph of f_i at most once: no two distinct x, w satisfy $f_i(x) = f_i(w)$.

• $i = 3, 4, 7, 8$. Why not injective?

At some 'altitude' $b_0 \in \mathbb{R}$, the horizontal line $y = b_0$ cuts the graph of f_i twice or more: some distinct x, w satisfy $f_i(x) = f_i(w)$.

Answer (b2). We re-interpret (b1) in terms of solving equations.

• $i = 1, 2, 5, 6$. Why injective?

For each $b \in \mathbb{R}$, the equation $b = f_i(u)$ with 'unknown' u in \mathbb{R} has at most one solution in \mathbb{R} .

• $i = 3, 4, 7, 8$. Why not injective?

There is some value $b_0 \in \mathbb{R}$ for which the equation $b_0 = f_i(u)$ with 'unknown' u in \mathbb{R} has two or more solutions in \mathbb{R} .

Answer (a). Directly verify the condition (I) or its negation respectively.

- $i = 1, 2, 5, 6$. Injectivity? [Recall the statement (I).]

* How do we check the injectivity of f_6 , in practice?

$$\forall x, w \in \mathbb{R}, \text{ if } f_6(x) = f_6(w) \text{ then } x = w.$$

Pick any $x, w \in \mathbb{R}$. [These x, w are fixed in the discussion below. We verify that if $f_6(x) = f_6(w)$ then $x = w$.]

Suppose $f_6(x) = f_6(w)$.

$$\text{Then } \frac{e^x - e^{-x}}{e^x + e^{-x}} = \frac{e^w - e^{-w}}{e^w + e^{-w}}.$$

$$\text{Therefore } e^{x+w} + e^{x-w} - e^{-x+w} - e^{-x-w} = (e^x - e^{-x})(e^w + e^{-w}) = (e^x + e^{-x})(e^w - e^{-w}) \\ = e^{x+w} - e^{x-w} + e^{-x+w} - e^{-x-w}.$$

$$\text{Hence } 2e^{x-w} = 2e^{-x+w}. \text{ Then } x-w = -x+w. \text{ We have } x=w.$$

It follows that f_6 is injective. \square

* How about f_1 ?

Pick any $x, w \in \mathbb{R}$. Suppose $f_1(x) = f_1(w)$.

$$\text{Then } 0.1x^3 = 0.1w^3.$$

$$\text{Therefore } (x-w)(x^2+xw+w^2) = x^3-w^3 = 0.$$

$\therefore \leftarrow$ [your work.]

$$\text{Hence } x = w.$$

It follows that f_1 is injective. \square

* How about f_2, f_5 ? [Exercise.]

Answer (a). Directly verify the condition (I) or its negation respectively.


- $i = 3, 4, 7, 8$. Non-injectivity? [Recall the statement $\sim(I)$.]

How do we check the non-injectivity of f_7 , in practice?

$$\exists x_0, w_0 \in \mathbb{R} \text{ such that } \underline{x_0 \neq w_0} \text{ and } \underline{f_7(x_0) = f_7(w_0)}.$$

[Name $x_0, w_0 \in \mathbb{R}$ for which $x_0 \neq w_0$ and $f_7(x_0) = f_7(w_0)$. Try this rough-work: Start with the 'relation' ' $f_7(x_0) = f_7(w_0)$ ' to see what may prevent us from obtaining ' $x_0 = w_0$ '.]

Take $x_0 = \frac{1}{2}, w_0 = -\frac{1}{2}$.



• $x_0, w_0 \in \mathbb{R}$.

• $x_0 \neq w_0$.

• $f_7(x_0) = \frac{1}{1+x_0^2} = \frac{1}{1+(\frac{1}{2})^2} = \frac{4}{5}$.

$f_7(w_0) = \frac{1}{1+w_0^2} = \frac{1}{1+(-\frac{1}{2})^2} = \frac{4}{5}$.

So $f_7(x_0) = f_7(w_0)$.

It follows that f_7 is not injective. \square

Roughwork:

Ask: what happens when $\underline{f_7(x_0) = f_7(w_0)}$?

$$f_7(x_0) = f_7(w_0)$$

$$\Rightarrow \frac{1}{1+x_0^2} = \frac{1}{1+w_0^2}$$

$$\Rightarrow 1+x_0^2 = 1+w_0^2$$

$$\Rightarrow x_0^2 = w_0^2$$

$$\Rightarrow |x_0| = |w_0|$$

Now ask: Can we name some distinct $x_0, w_0 \in \mathbb{R}$ satisfying $|x_0| = |w_0|$ and $\underline{f_7(x_0) = f_7(w_0)}$?

How about f_3, f_4, f_8 ? [Exercise.]