

1. Definition.

Let A, B be sets.

The set $\text{Map}(A, B)$ is defined to be the set of all functions from A to B .

Remark.

$\text{Map}(\mathbb{N}, B)$ is the set of all infinite sequences in B :

each $\varphi \in \text{Map}(\mathbb{N}, B)$ is the infinite sequence

$$(\varphi(0), \varphi(1), \varphi(2), \dots, \varphi(n), \varphi(n+1), \dots),$$

with each term being an element of B .

2. A basic example of unequal cardinality: $\mathbb{N} \not\sim \text{Map}(\mathbb{N}, \{0, 1\})$.

Theorem (VI).

There is no surjective function from \mathbb{N} to $\text{Map}(\mathbb{N}, \{0, 1\})$.

Corollary (VII).

There is no bijective function from \mathbb{N} to $\text{Map}(\mathbb{N}, \{0, 1\})$.

3. Idea in the proof of Theorem (VI).

Suppose there were some surjective function, say, Φ , from \mathbb{N} to $\text{Map}(\mathbb{N}, \{0, 1\})$.

We look for a contradiction against this false assumption.

For each $n \in \mathbb{N}$, the mathematical object $\Phi(n)$ is an infinite sequence in $\{0, 1\}$.

Since Φ was surjective, *every* infinite sequence in $\{0, 1\}$ would *appear somewhere* in the (infinite) list of infinite sequences

$$\begin{aligned}\Phi(0) &= ((\Phi(0))(0), (\Phi(0))(1), (\Phi(0))(2), (\Phi(0))(3), \dots), \\ \Phi(1) &= ((\Phi(1))(0), (\Phi(1))(1), (\Phi(1))(2), (\Phi(1))(3), \dots), \\ \Phi(2) &= ((\Phi(2))(0), (\Phi(2))(1), (\Phi(2))(2), (\Phi(2))(3), \dots), \\ \Phi(3) &= ((\Phi(3))(0), (\Phi(3))(1), (\Phi(3))(2), (\Phi(3))(3), \dots), \\ &\vdots \quad \vdots\end{aligned}$$

Out of this list of infinite sequences, we construct an *extra* infinite sequence in $\{0, 1\}$ which would *not appear* in this list. This is the desired contradiction.

The method of construction for this extra sequence is known as **Cantor's diagonal argument**.

4. Illustration of Cantor's diagonal argument through a 'specific' Φ .

For the sake of illustration, let us assume that $\Phi(0), \Phi(1), \Phi(2), \Phi(3), \Phi(4), \Phi(5), \dots$ are respectively given by:

$$\begin{array}{l}
 \Phi(0) = (\quad , \quad , \quad , \quad , \quad , \quad , \dots) \\
 \Phi(1) = (\quad , \quad , \quad , \quad , \quad , \quad , \dots) \\
 \Phi(2) = (\quad , \quad , \quad , \quad , \quad , \quad , \dots) \\
 \hline
 \Phi(3) = (\quad , \quad , \quad , \quad , \quad , \quad , \dots) \\
 \Phi(4) = (\quad , \quad , \quad , \quad , \quad , \quad , \dots) \\
 \Phi(5) = (\quad , \quad , \quad , \quad , \quad , \quad , \dots) \\
 \dots
 \end{array}$$

List out each of the terms of $\Phi(0), \Phi(1), \Phi(2), \Phi(3), \Phi(4), \Phi(5), \dots$ row by row to form the 'infinite' table below:

n	$\Phi(n)$	$(\Phi(n))(0)$	$(\Phi(n))(1)$	$(\Phi(n))(2)$	$(\Phi(n))(3)$	$(\Phi(n))(4)$	$(\Phi(n))(5)$	\dots	'Diagonal'	'Reverse of diagonal'
0	$\Phi(0)$							\dots		
1	$\Phi(1)$							\dots		
2	$\Phi(2)$							\dots		
3	$\Phi(3)$							\dots		
4	$\Phi(4)$							\dots		
5	$\Phi(5)$							\dots		
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
???	λ							\dots		
		$\lambda(0)$	$\lambda(1)$	$\lambda(2)$	$\lambda(3)$	$\lambda(4)$	$\lambda(5)$	\dots		

Expected: Because Φ is surjective, we expect every infinite sequence in $\{0,1\}$ to turn up here.

Question.

- Is there any $z \in \mathbb{N}$ which satisfies $\Phi(z) = \lambda$, really?

Answer.

- Since $\lambda(0)$ is obtained by the 'flip' of $(\Phi(0))(0)$, λ does not agree with $\Phi(0)$ at the 0-th term. Hence $\lambda \neq \Phi(0)$.

Since $\lambda(1)$ is obtained by the 'flip' of $(\Phi(1))(1)$, λ does not agree with $\Phi(1)$ at the 1-st term. Hence $\lambda \neq \Phi(1)$

Since $\lambda(n)$ is obtained by the 'flip' of $(\Phi(n))(n)$, λ does not agree with $\Phi(n)$ at the n -th term. Hence $\lambda \neq \Phi(n)$. Et cetera.

Hence λ is not amongst $\Phi(0), \Phi(1), \Phi(2), \Phi(3), \Phi(4), \Phi(5), \dots$.

This λ is the '*extra*' infinite sequence 'generated' by this 'specific' Φ according to Cantor's diagonal argument.

(No matter which 'specific' Φ we start with, the procedure described above will give you a corresponding 'troublesome' λ . Try your own favourite 'concrete' $\Phi(0), \Phi(1), \Phi(2), \Phi(3), \Phi(4), \Phi(5), \dots$, go through the procedure described above, and have fun.)

5. Formal argument for Theorem (VI).

Suppose there were some surjective function $\Phi : \mathbb{N} \longrightarrow \text{Map}(\mathbb{N}, \{0, 1\})$.

Define the function $\lambda : \mathbb{N} \longrightarrow \{0, 1\}$ by

$$\lambda(x) = \begin{cases} 1 & \text{if } (\Phi(x))(x) = 0 \\ 0 & \text{if } (\Phi(x))(x) = 1 \end{cases}$$

(Is λ well-defined as a function?)

Since Φ was surjective, there would be some $z \in \mathbb{N}$ so that $\Phi(z) = \lambda$.

However, by definition, we have $(\Phi(z))(z) \neq \lambda(z)$.

Therefore $\lambda \neq \Phi(z)$. Contradiction arises.

Hence there is no surjective function from \mathbb{N} to $\text{Map}(\mathbb{N}, \{0, 1\})$ in the first place.

6. Theorem (VIII).

Let A be a set. The following statements hold:

- (1) *There is no surjective function from A to $\text{Map}(A, \{0, 1\})$.*
- (2) *There is no bijective function from A to $\text{Map}(A, \{0, 1\})$.*
- (3) $A \not\sim \text{Map}(A, \{0, 1\})$.

Proof.

The proof for Statement (1) in Theorem (VIII) is almost the same as that for Theorem (VI): just replace \mathbf{N} by A in the formal proof for Theorem (VI).

Statements (2), (3) follow immediately from Statement (1).

Proof of Statement (1) in Theorem (VIII).

~~Formal argument for Theorem (VI).~~

Suppose there were some surjective function $\Phi : \mathbb{N} \rightarrow \text{Map}(\mathbb{N}, \{0, 1\})$.

Define the function $\lambda : \mathbb{N} \rightarrow \{0, 1\}$ by

$$\lambda(x) = \begin{cases} 1 & \text{if } (\Phi(x))(x) = 0 \\ 0 & \text{if } (\Phi(x))(x) = 1 \end{cases}$$

(Is λ well-defined as a function?)

Since Φ was surjective, there would be some $z \in \mathbb{N}$ so that $\Phi(z) = \lambda$.

However, by definition, we have $(\Phi(z))(z) \neq \lambda(z)$.

Therefore $\lambda \neq \Phi(z)$. Contradiction arises.

Hence there is no surjective function from \mathbb{N} to $\text{Map}(\mathbb{N}, \{0, 1\})$ in the first place.

Theorem (VIII).

Let A be a set. The following statements hold:

- (1) There is no surjective function from A to $\text{Map}(A, \{0, 1\})$.
- (2) There is no bijective function from A to $\text{Map}(A, \{0, 1\})$.
- (3) $A \not\sim \text{Map}(A, \{0, 1\})$.

Another formulation of Theorem (VIII).

Let A be a set. The following statements hold:

- (1) There is no surjective function from A to $\mathfrak{P}(A)$.
- (2) There is no bijective function from A to $\mathfrak{P}(A)$.
- (3) $A \not\sim \mathfrak{P}(A)$.

Why? How?

• Recall that $\mathfrak{P}(A) \sim \text{Map}(A, \{0, 1\})$.

$\chi: \mathfrak{P}(A) \rightarrow \text{Map}(A, \{0, 1\})$ given by
 $\chi^A(S) = \chi_S^A$ for any $S \in \mathfrak{P}(A)$
is a bijective function.