

Suggested solution to Assignment 2

1. (a)

$$\begin{array}{cccc} & & & \min \\ & \begin{pmatrix} 5 & 1 & -2 & 6 \\ -1 & 0 & 1 & -2 \\ 3 & 2 & 5 & 4 \end{pmatrix} & & \\ \max & 5 & 2 & 5 & 6 \end{array}$$

Both the maximin and minimax are 2. Therefore the entry  $a_{32} = 2$  is a saddle point. The value of the game is 2.

(b)

$$\begin{array}{cccc} & & & \min \\ & \begin{pmatrix} -4 & 5 & -3 & -3 \\ 0 & 1 & 3 & -1 \\ -3 & -1 & 2 & -5 \\ 2 & -4 & 0 & -2 \end{pmatrix} & & \\ \max & 2 & 5 & 3 & -1 \end{array}$$

Both the maximin and minimax are -1. Therefore the entry  $a_{24} = -1$  is a saddle point. The value of the game is -1.

2. (a)

$$A = \begin{pmatrix} 1 & 7 \\ 2 & -2 \end{pmatrix} \begin{array}{l} -6 \times 4 \\ 4 \times 6 \end{array} \begin{pmatrix} \frac{2}{5} \\ \frac{3}{5} \end{pmatrix}$$

$$\begin{array}{cc} & -1 & 9 \\ & 9 & 1 \\ \begin{pmatrix} \frac{9}{10} \\ \frac{1}{10} \end{pmatrix} & & \end{array}$$

So the maximin strategy for the row player is  $(\frac{2}{5}, \frac{3}{5})$ .  
 The minimax strategy for the column player is  $(\frac{9}{10}, \frac{1}{10})$ .  
 The value of the game is  $v = (\frac{2}{5}, \frac{3}{5}) \begin{pmatrix} 1 & 7 \\ 2 & -2 \end{pmatrix} \begin{pmatrix} \frac{9}{10} \\ \frac{1}{10} \end{pmatrix} = \frac{8}{5}$ .

(b)

$$A = \begin{pmatrix} 3 & -1 \\ -2 & 4 \end{pmatrix} \begin{array}{l} 4 \times 6 \\ -6 \times 4 \end{array} \begin{pmatrix} \frac{3}{5} \\ \frac{2}{5} \end{pmatrix}$$

$$\begin{array}{cc} & 5 & -5 \\ & 5 & 5 \\ \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} & & \end{array}$$

So the maximin strategy for the row player is  $(\frac{3}{5}, \frac{2}{5})$ .  
 The minimax strategy for the column player is  $(\frac{1}{2}, \frac{1}{2})$ .  
 The value of the game is  $v = (\frac{3}{5}, \frac{2}{5}) \begin{pmatrix} 3 & -1 \\ -2 & 4 \end{pmatrix} \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix} = 1$

$$(c) A = \begin{pmatrix} 3 & 2 & 4 & 0 \\ -2 & 1 & -4 & 5 \end{pmatrix}$$


By drawing the lower envelope, the maximum point of the lower envelope is the intersection point of  $C_1$  and  $C_4$ . By solving

$$\begin{cases} C_1: V = 3x - 2(1-x) \\ C_4: V = 5(1-x) \end{cases}$$

$$x = 0.7, V = 1.5$$

For the minimax strategy:  $\begin{pmatrix} 3 & 0 \\ -2 & 5 \end{pmatrix} \begin{pmatrix} y_1 \\ y_4 \end{pmatrix} = \begin{pmatrix} 1.5 \\ 1.5 \end{pmatrix} \Rightarrow y_1 = y_4 = 0.5$ .

Hence the maximin strategy for the row player is  $(0.7, 0.3)$ ; the minimax strategy for the column player is  $(0.5, 0, 0, 0.5)$ ; and the value is 1.5.



$$(d) A = \begin{pmatrix} 1 & 0 & 4 & 2 \\ 0 & 2 & -3 & -2 \end{pmatrix}$$

By drawing the lower envelope, the maximum point of the lower envelope is the intersection point of  $C_1, C_2$  and  $C_4$ . By solving

$$\begin{cases} C_1: V = x \\ C_2: V = 2(1-x) \\ C_4: V = 2x - 2(1-x) \end{cases}$$

$$x = \frac{2}{3}, V = \frac{2}{3}$$

For the minimax strategy:  $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 0 & 2 & -2 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_4 \end{pmatrix} = \begin{pmatrix} 1 \\ \frac{2}{3} \\ \frac{2}{3} \end{pmatrix}$

Note that we have added the equation  $y_1 + y_2 + y_4 = 1$  to exclude the solutions which are not probability vectors. Using row operation, we obtain the row echelon for

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & \frac{2}{3} \\ 1 & 0 & 2 & \frac{2}{3} \\ 0 & 2 & -2 & \frac{2}{3} \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 2 & \frac{2}{3} \\ 0 & 1 & -1 & \frac{1}{3} \\ 0 & 0 & 0 & 0 \end{array} \right)$$


The non-negative solution to the system of equations is  $(y_1, y_2, y_4) = (\frac{2}{3} - 2t, \frac{1}{3} + t, t)$ ,  $0 \leq t \leq \frac{1}{3}$ .

Hence the column player has minimax strategies  $q = (\frac{2}{3} - 2t, \frac{1}{3} + t, t)$  for  $0 \leq t \leq \frac{1}{3}$ .

In particular,  $(\frac{2}{3}, \frac{1}{3}, 0, 0)$  and  $(0, \frac{2}{3}, 0, \frac{1}{3})$  are minimax strategies for the column player;

The maximin strategy for the row player is  $(\frac{2}{3}, \frac{1}{3})$ ;

The value of the game is  $\frac{2}{3}$ .



$$(e) A = \begin{pmatrix} 5 & -3 \\ -3 & 5 \\ 2 & -1 \\ 4 & 0 \end{pmatrix}$$


The upper envelope is shown in the right figure.

$$\text{Solving } \begin{cases} C_2: V = -3y + 5(1-y) \\ C_4: V = 4y \end{cases}$$

$$y = \frac{5}{12}, V = \frac{5}{3}$$

$$\begin{pmatrix} -3 & 5 \\ 4 & 0 \end{pmatrix} \begin{matrix} -8 & 4 \\ 4 & 8 \end{matrix} \begin{pmatrix} \frac{1}{3} \\ \frac{2}{3} \end{pmatrix}$$

$$\begin{matrix} 7 & 5 \\ 5 & 7 \end{matrix} \begin{pmatrix} \frac{5}{12} \\ \frac{7}{12} \end{pmatrix}$$



Therefore the maximin strategy for the row player and the minimax strategy for the column player are  $(0, \frac{1}{3}, 0, \frac{2}{3})$  and  $(\frac{5}{12}, \frac{7}{12})$  respectively; the value is  $\frac{5}{3}$ .

$$(f) A = \begin{pmatrix} 5 & -2 & 3 \\ 3 & -1 & 4 \\ 0 & 3 & 1 \end{pmatrix} \quad A \text{ is nonsingular}$$

Using row operation, we obtain

$$\left( \begin{array}{ccc|ccc} 5 & -2 & 3 & 1 & 0 & 0 \\ 3 & -1 & 4 & 0 & 1 & 0 \\ 0 & 3 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{13}{32} & -\frac{11}{32} & \frac{5}{32} \\ 0 & 1 & 0 & \frac{3}{32} & -\frac{5}{32} & \frac{11}{32} \\ 0 & 0 & 1 & -\frac{9}{32} & \frac{15}{32} & -\frac{1}{32} \end{array} \right)$$

$$\text{Hence } A^{-1} = \begin{pmatrix} \frac{13}{32} & -\frac{11}{32} & \frac{5}{32} \\ \frac{3}{32} & -\frac{5}{32} & \frac{11}{32} \\ -\frac{9}{32} & \frac{15}{32} & -\frac{1}{32} \end{pmatrix}, \quad \mathbf{1}^T A^{-1} \mathbf{1} = \frac{21}{32} \quad (\mathbf{1} = (1, 1, 1)^T)$$

$$\text{By tutorial notes, } v = 1/\mathbf{1}^T A^{-1} \mathbf{1} = \frac{32}{21} \neq 0, \quad p^T = \mathbf{1}^T A^{-1} = \left( \frac{7}{21}, -\frac{1}{21}, \frac{15}{21} \right)^T$$

Since  $p$  has negative component, there is a theorem showing that an arbitrary  $m \times n$  matrix game whose value is not zero may be solved by choosing some suitable square submatrix and checking the resulting optimal strategies for the whole matrix.

$$\text{By checking, we choose the submatrix } A' = \begin{pmatrix} 5 & -2 \\ 0 & 3 \end{pmatrix} \begin{matrix} 7 & 3 \\ -3 & 7 \end{matrix} \begin{pmatrix} \frac{3}{10} \\ \frac{7}{10} \end{pmatrix}$$

$$\begin{matrix} 5 & -5 \\ 5 & 5 \end{matrix} \begin{pmatrix} \frac{5}{2} \\ \frac{5}{2} \end{pmatrix}$$

Hence a maximin strategy for the row player is  $P = (\frac{3}{10}, 0, \frac{7}{10})$ ; a minimax strategy for the column player is  $Q = (\frac{1}{2}, \frac{1}{2}, 0)$ ; the value is  $V = \frac{5 \times 3 - 0}{5 + 3 - (-2)} = \frac{3}{2}$ .

One may check the result by the following calculations.

$$PA = \left(\frac{3}{10}, 0, \frac{7}{10}\right) \begin{pmatrix} 5 & -2 & 3 \\ 3 & -1 & 4 \\ 0 & 3 & 1 \end{pmatrix} = \left(\frac{3}{2}, \frac{3}{2}, \frac{8}{5}\right)$$

$$AQ^T = \begin{pmatrix} 5 & -2 & 3 \\ 3 & -1 & 4 \\ 0 & 3 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{3}{2} \\ \frac{1}{2} \\ \frac{3}{2} \end{pmatrix}$$

One sees that the row player may guarantee that his payoff is at least  $\frac{3}{2}$  by using  $P = (\frac{3}{10}, 0, \frac{7}{10})$  and the column player may guarantee the payoff to the row player is at most  $\frac{3}{2}$  by using  $Q = (\frac{1}{2}, \frac{1}{2}, 0)$ .

$$(9) \quad A = \begin{pmatrix} 5 & 1 & -2 & 6 \\ -1 & 0 & 1 & -2 \\ 3 & 2 & 5 & 4 \end{pmatrix} \begin{array}{l} \min \\ -2 \\ 2 \\ 2 \end{array}$$

max 5 2 5 6

Both the maximin and minimax are 2. Hence  $a_{32} = 2$  is a saddle point.

Then the value of the game is 2.

Obviously, the maximin strategy for the row player is  $(0, 0, 1)$ ;

the minimax strategy for the column player is  $(0, 1, 0, 0)$ .

3. The game matrix is  $A = \begin{pmatrix} -3 & 2 \\ 9 & -8 \end{pmatrix}$ .

$$\begin{pmatrix} -3 & 2 \\ 9 & -8 \end{pmatrix} \begin{array}{l} -5 \\ 17 \end{array} \times \begin{array}{l} 17 \\ 5 \end{array} \begin{pmatrix} \frac{17}{22} \\ \frac{5}{22} \end{pmatrix}$$

~~$\begin{pmatrix} -12 & 10 \\ 10 & 12 \end{pmatrix}$~~

$$\left(\frac{5}{11}, \frac{6}{11}\right)$$

Therefore the value of the game is  $\left(\frac{17}{22}, \frac{5}{22}\right) \begin{pmatrix} -3 & 2 \\ 9 & -8 \end{pmatrix} \begin{pmatrix} \frac{5}{11} \\ \frac{6}{11} \end{pmatrix} = -\frac{3}{11}$ ;

the optimal strategy for Raymond is  $\left(\frac{17}{22}, \frac{5}{22}\right)$ ;

the optimal strategy for Calvin is  $\left(\frac{5}{11}, \frac{6}{11}\right)$ .

4. (a) Alex 1  $\begin{matrix} \text{Becky 1} & \text{Becky 2} \\ \begin{pmatrix} 3 & -1 \\ -1 & 11 \end{pmatrix} \end{matrix}$ , i.e. the game matrix is  $\begin{pmatrix} 3 & -1 \\ -1 & 11 \end{pmatrix}$ .

$$\begin{pmatrix} 3 & -1 \\ -1 & 11 \end{pmatrix} \begin{matrix} 4 & 12 \\ -12 & 4 \end{matrix} \begin{matrix} \frac{3}{4} \\ \frac{1}{4} \end{matrix}$$

$$\begin{matrix} 4 & -12 \\ 12 & 4 \end{matrix}$$

$$\begin{pmatrix} \frac{3}{4} & \frac{1}{4} \end{pmatrix}$$

Hence the optimal strategies for Alex and Becky are both  $(\frac{3}{4}, \frac{1}{4})$ .

(b) By (a), the value of the game is  $v = (\frac{3}{4}, \frac{1}{4}) \begin{pmatrix} 3 & -1 \\ -1 & 11 \end{pmatrix} \begin{pmatrix} \frac{3}{4} \\ \frac{1}{4} \end{pmatrix} = 2$ .

Hence to make the game fair,  $k = v = 2$ .

5. The game matrix is

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

We may delete the first row and the last row since they are dominated by the third and the fifth row respectively to get the reduced matrix

$$A' = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \end{pmatrix}$$

Then we may delete the ~~third~~ and the ~~fifth~~ columns since they are dominated by the first, the second and the fourth the last columns respectively to get the reduced matrix

$$A'' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Finally we may delete the third row since it is dominated by any other row.

Hence the matrix  $A$  is reduced to the  $4 \times 4$  **diagonal** matrix

$$A''' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \text{ where diagonal entries } d_i = 1, i=1, 2, 3, 4.$$

Using the principle of indifference,  $v = \frac{1}{\mathbf{1}^T A''' \mathbf{1}} = \left( \sum_{i=1}^4 1/d_i \right)^{-1} = 1 \cdot (\mathbf{1} = (1, 1, 1, 1)^T)$

And  $p = v A'''^{-T} \mathbf{1} = v (1/d_1, \dots, 1/d_4) = (1, 1, 1, 1)$

Similarly,  $q = v A'''^{-1} \mathbf{1} = v (1/d_1, \dots, 1/d_4) = (1, 1, 1, 1)$ .

Therefore, the value of  $A$  is 1; the optimal strategies for player I and player II are  $(0, 1, 0, 1, 1, 1, 0)$  and  $(1, 1, 0, 0, 0, 1, 1)$  respectively.

6. Assume that II has optimal strategy giving positive weight in each entry.

By principle of indifference, I's optimal strategy  $p$  satisfies

$$\sum_{i=1}^m p_i a_{ij} = v, \quad j=1, 2, \dots, m.$$

Thus  $p_1 = v, -2p_1 + p_2 = v, 3p_1 - 2p_2 + p_3 = v, -4p_1 + 3p_2 - 2p_3 + p_4 = v.$

Solving  $p_1 = v, p_2 = 3v, p_3 = 4v, p_4 = 4v$

Since  $\sum_{i=1}^4 p_i = 1$ , we get  $12v = 1$ , thus  $v = 1/12.$

And  $p = (p_1, p_2, p_3, p_4) = (1/12, 1/4, 1/3, 1/3).$

Similar argument shows that  $q = (1/3, 1/3, 1/4, 1/12).$

Since both  $p$  and  $q$  are nonnegative, both are optimal strategies and  $1/12$  is the value of the game.

7. By the condition, the game matrix is

$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{matrix} & \begin{pmatrix} 0 & -1 & 2 & 2 & 2 \\ 1 & 0 & -1 & 2 & 2 \\ -2 & 1 & 0 & -1 & 2 \\ -2 & -2 & 1 & 0 & -1 \\ -2 & -2 & -2 & 1 & 0 \end{pmatrix} \end{matrix}$$

This game is symmetric, so the value is zero.

Note that row 1 dominates rows 4 and 5, and column 1 dominated columns 4 and 5.

We only need to consider the upper left  $3 \times 3$  submatrix  $\begin{pmatrix} 0 & -1 & 2 \\ 1 & 0 & -1 \\ -2 & 1 & 0 \end{pmatrix}.$

Assume that II has optimal strategy  $q'$  so that  $q'_1 > 0, q'_2 > 0, q'_3 > 0.$

By the principle of indifference, we have

$$p'_2 - 2p'_3 = 0, \quad -p'_1 + p'_3 = 0, \quad 2p'_1 - p'_2 = 0$$

Together with the condition  $p'_1 + p'_2 + p'_3 = 0$ , we have  $p'_1 = 1/4, p'_2 = 1/2, p'_3 = 1/4.$

Similarly, we can get  $q'_1 = 1/4, q'_2 = 1/2, q'_3 = 1/4.$

Hence, the optimal strategies are  $p = q = (1/4, 1/2, 1/4, 0, 0).$

9 (a) 
$$\begin{pmatrix} -3 & 1 \\ c & -2 \end{pmatrix} \begin{matrix} \min \\ -3 \\ x \end{matrix}$$

$$\begin{matrix} \max y \\ 1 \end{matrix}$$

(i) If  $c \leq -3$ , then  $x=c$ , so minimax is  $-3$ ; and  $y=-3$ , so the maximin is  $-3$ .

Hence we see that  $\min \max = \max \min$ , i.e. A has a saddle point if  $c \leq -3$ .

(ii) If  $-3 < c \leq -2$ , then  $x=c$ , so minimax is  $c$ ; and  $y=c$ , so maximin is  $c$ .

Hence A has a saddle point if  $-3 < c \leq -2$ .

(iii) If  $c > -2$ , then  $x=-2$ , so minimax is  $-2$ ; and  $y=c$ , so maximin  $> -2$ .

Hence A has no saddle point if  $c > -2$ .

Therefore, if  $c \leq -2$ , then A has a saddle point.

(b) (i) By the hypothesis, the value of A is 0.

$$\text{Thus } v = \frac{(-3)(-2) - 1 \cdot c}{-3 - 2 - 1 - c} = 0 \implies c = 6.$$

(ii) By (i),  $A = \begin{pmatrix} -3 & 1 \\ 6 & -2 \end{pmatrix}$ .

Hence the maximin strategy for the row player is  $p = \left( \frac{-2-6}{-3-2-6-1}, \frac{-3-1}{-3-2-6-1} \right) = \left( \frac{2}{3}, \frac{1}{3} \right)$ ;  
the minimax strategy for the column player is  $q = \left( \frac{-2-1}{-3-2-6-1}, \frac{-3-6}{-3-2-6-1} \right) = \left( \frac{1}{4}, \frac{3}{4} \right)$ .

10. Method 1:

Since  $A^T = -A \implies A = -A^T$ , the value of A and  $-A^T$  are the same.

Hence  $v = -v$ , i.e.  $2v = 0 \implies v = 0$ .

Therefore, the value of A is zero.

Method 2:

Let  $p$  be an optimal strategy for I. If II uses the same strategy, then

$$p^T A p = \sum_i \sum_j p_i a_{ij} p_j = \sum_i \sum_j p_i (-a_{ji}) p_j = - \sum_j \sum_i p_j a_{ji} p_i = -p^T A p.$$

Hence  $p^T A p = 0$ . This shows that the value  $v \leq p^T A p = 0$ , i.e.  $v \leq 0$ .

A symmetric argument shows that  $v \geq q^T A q^T = 0$ , i.e.  $v \geq 0$ .

Therefore, the value of A is  $v = 0$ .



		Ben	
		Left	Right
8. (a) Aaron	Left	-2	4
	Right	3	-1

(b) value  $v = \frac{-2 \times (-1) - 3 \times 4}{-2 - 4 - 3 - 1} = 1$

Maximin strategy for Aaron is  $\left( \frac{1-3}{-10}, \frac{-2-4}{-10} \right) = \left( \frac{2}{5}, \frac{3}{5} \right)$

Minimax strategy for Ben is  $\left( \frac{-1-4}{-10}, \frac{-2-3}{-10} \right) = \left( \frac{1}{2}, \frac{1}{2} \right)$

11. (a) (1)  $0 \leq Ay^T \leq v \mathbf{1}^T$  (2)  $yA = yA^T = (Ay^T)^T = (v \mathbf{1}^T)^T = v \mathbf{1}$   
 (3)  $yAy^T = y(v \mathbf{1}^T) = v$ ,  $y \in P^n$   
 by Minimax Thm,  $v$  is the value.

(b) Let  $A = \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$   $\begin{matrix} \text{max} \\ -1 \\ -1 \\ 1 \end{matrix}$

Since maximum and minimum are 1

$\Rightarrow$  the value of  $A$  is 1

$$\begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{3} \\ \frac{1}{3} \\ \frac{1}{3} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \text{ ok}$$

13. (a)  $x = (1, 1, 2, 3, 5)$ ,  $a = -2$

(b)  $y = (1, 0, 2, 1, 4)$ ,  $b = -3$

(c)  $2 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 2 \end{pmatrix} + \beta \begin{pmatrix} 1 \\ 1 \\ 1 \\ -3 \\ 1 \end{pmatrix} = \begin{pmatrix} \alpha \\ \alpha \\ \alpha \\ \alpha \\ \alpha \end{pmatrix} \Rightarrow \beta = \alpha, -2\alpha - 3\beta = \alpha$   
 $\Rightarrow \alpha = -2\beta$

Also, since  $y \in P^5$ ,  $\Rightarrow \alpha(x_1 + \dots + x_5) + \beta(y_1 + \dots + y_5) =$

$$12\alpha + 8\beta = 1$$

$$-24V + 8V = 1$$

$$-16V = 1$$

$$\Rightarrow V = -\frac{1}{16}$$

$$\Rightarrow \alpha = \frac{1}{8}, \beta = -\frac{1}{16}$$

$$\begin{aligned} \Rightarrow q &= \frac{1}{8} (1 \ 1 \ 2 \ 3 \ 5) + (-\frac{1}{16}) (1 \ 0 \ 2 \ 1 \ 4) \\ &= \frac{1}{16} (1 \ 2 \ 2 \ 5 \ 6) \end{aligned}$$

For 11(a), Since  $A$  is symmetric and

$$Aq^T = -\frac{1}{16} \mathbf{1}^T$$

$\Rightarrow$  the maximin strategy is  $q$ ,  
minimax strategy is  $q$  and  $V = -\frac{1}{16}$

14. (a) The max strategy of  $A_k$  is  $P = \left( \frac{8k+1}{16k-6}, \frac{8k-5}{16k-6} \right)$   
The min strategy of  $A_k$  is  $q = \left( \frac{8k-2}{16k-6}, \frac{8k-4}{16k-6} \right)$

the value of this game is

$$V = \frac{(4k-3)4k - (4k-2)(4k-1)}{4k-3 + 4k + 4k-2 + 4k-1} = \frac{1}{3-8k}$$

(b) Suppose  $\hat{p} = (p_1, p_2, \dots, p_n)$  is the maximum strategy of  $D$

$$\Rightarrow \hat{p}A = \left( \frac{p_1}{r_1}, \dots, \frac{p_n}{r_n} \right)$$

by principle of indifference,  $p_1/r_1 = \dots = p_n/r_n = V$

$$\Rightarrow p_i = r_i V$$

$$\text{Also, } p_1 + \dots + p_n = 1$$

$$\Rightarrow V(r_1 + \dots + r_n) = 1 \quad \Rightarrow V = \frac{1}{r_1 + r_2 + \dots + r_n}$$

$$(c) V = \frac{1}{r_1 + r_2 + \dots + r_n} = \frac{1}{\sum_{k=1}^{25} \frac{1}{A_k}} = \frac{1}{\sum_{k=1}^{25} (3-8k)}$$