

1.* Show that the uniform limit of a sequence of continuous functions is continuous, and hence that if $m(E) < +\infty$ and $f: E \rightarrow \mathbb{R}$ is measurable then, $\forall \eta > 0$, \exists closed set $F \subseteq E$ with $m(E \setminus F) < \eta$ such that $f|_F: F \rightarrow \mathbb{R}$ is continuous.

2. Let $F = \bigcup_{n=1}^{\infty} F_n$, disjoint closed sets F_1, \dots, F_n .

Let $f: F \rightarrow \mathbb{R}$ be such that $f|_{F_n}$ is cts, $\forall n$.

Show that f is cts.

3.* Let $F_n \subseteq (n, n+1]$ be closed ($\mathbb{R} \setminus F_n$ open) $\forall n \in \mathbb{N}$, and let $F = \bigcup_{n \in \mathbb{N}} F_n$.

Show that $f: F \rightarrow \mathbb{R}$ is continuous if each $f|_{F_n}$ is cts. (Can the condition

$F_n \subseteq (n, n+1]$ be weakened to $F_n \subseteq \mathbb{R}$?)

4. Let $G = \bigcup_{n=1}^{\infty} I_n$, countable disjoint open intervals I_n , and let $F: \mathbb{R} \setminus G$. Let $x < y < z$ with $x, z \in F$ and $y \in I_n := (a_n, b_n)$. Show that $a_n \in F$, $b_n \in F$, $x \leq a_n$, and $b_n \leq z$.

5. Let G, I_n, F be as in Q4, and let $f: \mathbb{R} \rightarrow \mathbb{R}$ be such that $f|_F$ and $f|_{\overline{I_n}}$ be

continuous, $\forall n \in \mathbb{N}$ ($\overline{I_n}$ denotes the closure of I_n). Suppose further that the graph of $f|_{\overline{I_n}}$ is a line-segment.

Show that f is continuous (By symmetry, need only show that f is right-continuous at

each $x_0 \in \mathbb{R}$: $\lim_{x \rightarrow x_0^+} f(x) = f(x_0)$, i.e. $\forall \varepsilon > 0 \exists \delta > 0$ such that $|f(x) - f(x_0)| < \varepsilon \forall x \in (x_0, x_0 + \delta)$.)

This is evident if $x_0 \in G$ (so $\exists n \in \mathbb{N}$ s.t. $x_0 \in I_n$). We may hence assume that $x_0 \in F$, and there are three cases to consider :

$$(a) \exists \delta > 0 \text{ s.t. } (x_0, x_0 + \delta) \subseteq F \text{ (so } [x_0, x_0 + \delta] \subseteq F)$$

$$(b) \exists \delta > 0 \text{ s.t. } (x_0, x_0 + \delta) \subseteq G \text{ (so } (x_0, x_0 + \delta) \subseteq I_n \text{ for some } n)$$

$$(c) (x_0, x_0 + \delta) \text{ intersects } F \text{ and } G, \forall \delta > 0.$$

Hint: For case (a), you use the continuity of $f|_F$.

For case (b), you use the continuity of $f|_{[x_0, x_0 + \delta]}$.

For case (c), let $\varepsilon > 0$. $\exists \delta_0 > 0$ such that $|f(x) - f(x_0)| < \varepsilon \forall x \in F \cap [x_0, x_0 + \delta_0]$ as $f|_F$ is continuous at x_0 . By the assumption in case (c) and consider smaller $\delta_0 > 0$ if necessary, we may assume that $x_0 + \delta_0 \in F$. Show that if $x \in G \cap (x_0, x_0 + \delta_0)$, then $\exists!$ $n \in \mathbb{N}$ with $x \in (a_n, b_n)$. Since $x_0, x_0 + \delta_0 \in F$, one has (?)

$$x_0 \leq a_n < x < b_n \leq x_0 + \delta_0 \text{ and } a_n, b_n \in F,$$

$$|f(\cdot) - f(x_0)| < \varepsilon \text{ at } a_n, b_n \text{ \& so at } x.$$

6.* Do the same as Q5 but check "the left-continuity" in place of "the right-continuity"