MATH4050 Real Analysis
Homework 1

There are 9 questions in this assignment (your works on the questions with * will be marked). The page number and question number for each question correspond to that in Royden's Real Analysis, 3rd or 4th edition.

1* (3rd: P.12, Q6)

Let $f : X \rightarrow Y$ be a mapping of a nonempty space X into Y. Show that f is one-to-one if and only if there is a mapping $g : Y \rightarrow X$ such that $g \circ f$ is the identity map on X, that is, such that $g(f(x)) = x$ for all $x \in X$.

2. (3rd: P.12, Q7)

Let $f : X \rightarrow Y$ be a mapping of X into Y. Show that f is onto if there is a mapping $g : Y \rightarrow X$ such that $f \circ g$ is the identity map in Y, that is, such that $f(g(y)) = y$ for all $y \in Y$.

3. Show that any set X can be "indexed": $\exists$ a set I and a function $f : I \rightarrow X$ such that $\{f(i) : i \in I\} = X$.

4* (3rd: P.16, Q14)

Given a set B and a collection of sets $\mathcal{C}$. Show that

$$B \cap \left[\bigcup_{A \in \mathcal{C}} A \right] = \bigcup_{A \in \mathcal{C}} (B \cap A).$$

5. (3rd: P.16, Q15)

Show that if $\mathcal{A}$ and $\mathcal{B}$ are two collection of sets, then
$[\bigcup \{A : A \in \mathcal{A}\}] \cap [\bigcup \{B : B \in \mathcal{B}\}] = \bigcup \{A \cap B : (A, B) \in \mathcal{A} \times \mathcal{B}\}$.

6. (3rd: P.16, Q16)

Let $f : X \rightarrow Y$ be a function and $\{A_\lambda\}_{\lambda \in \Lambda}$ be a collection of subsets of X.

   a. Show that $f[\bigcup A_\lambda] = \bigcup f[A_\lambda]$.
   b. Show that $f[\bigcap A_\lambda] \subset \bigcap f[A_\lambda]$.
   c. Give an example where $f[\bigcap A_\lambda] \neq \bigcap f[A_\lambda]$.

7* (3rd: P.16, Q17)

Let $f : X \rightarrow Y$ be a function and $\{B_\lambda\}_{\lambda \in \Lambda}$ be a collection of subsets of Y.

   a. Show that $f^{-1}[\bigcup B_\lambda] = \bigcup f^{-1}[B_\lambda]$.
   b. Show that $f^{-1}[\bigcap B_\lambda] = \bigcap f^{-1}[B_\lambda]$.
   c. Show that $f^{-1}[B^c] = (f^{-1}[B])^c$ for $B \subset Y$.

8* (3rd: P.16, Q18)

    a. Show that if f maps X into Y and $A \subset X$, $B \subset Y$, then

$$f[f^{-1}[B]] \subset B$$

    and

$$f^{-1}[f[A]] \supset A.$$

    b. Give examples to show that we need not have equality.

    c. Show that if f maps X onto Y and $B \subset Y$, then

$$f[f^{-1}[B]] = B.$$

9. Show that $f \mapsto \int_0^1 f(x)dx$ is a "monotone" function on $R[0,1]$ (consisting of all Riemann integrable functions on $[0,1]$), and $R[0,1]$ is a linear space. Show further that

$$\lim_{n \to \infty} \int_0^1 f_n(x)dx = \int_0^1 f(x)dx$$

if $f, f_n \in R[0,1]$ such that

$$\lim_{n \to \infty} \left(\sup_{x \in [0,1]} |f_n(x) - f(x)| \right) = 0.$$

2