

§ 8.4 The Trigonometric Functions

Thm 8.4.1 \exists functions $C: \mathbb{R} \rightarrow \mathbb{R}$ and $S: \mathbb{R} \rightarrow \mathbb{R}$ such that


(i) $C''(x) = -C(x)$ and $S''(x) = -S(x)$, $\forall x \in \mathbb{R}$.

(ii) $\begin{cases} C(0) = 1 \\ C'(0) = 0 \end{cases}$ and $\begin{cases} S(0) = 0 \\ S'(0) = 1 \end{cases}$

Pf: Define $C_n(x)$ and $S'_n(x)$ inductively by

$$\begin{cases} C_1(x) = 1 \\ S'_1(x) = x \end{cases}$$

$$\begin{cases} S'_n(x) = \int_0^x C_n(t) dt \\ C_{n+1}(x) = 1 - \int_0^x S'_n(t) dt \end{cases}$$



Then "Induction": C_n & S'_n are continuous, $\forall n$
 \Rightarrow integrable on any bounded interval

\therefore All C_n & S'_n are well-defined.

Moreover, by Fundamental Thm 7.3.5,

$$S'_n(x) = C_n(x) \quad \& \quad C'_{n+1}(x) = -S'_n(x), \quad \forall x \in \mathbb{R}, \forall n$$

Claim :

$$\left\{ \begin{aligned} C_{n+1}(x) &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^n \frac{x^{2n}}{(2n)!} \\ S_{n+1}(x) &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} \end{aligned} \right.$$

Pf : (Ex! By induction)

Let $A > 0$.

If $x \in [-A, A]$ and $m > n > 2A$,
(ie. $|x| \leq A$)

$$\left(\frac{A}{2n}, \frac{A}{2m} < \frac{1}{4} \right)$$

then

$$\begin{aligned} |C_m(x) - C_n(x)| &= \left| (-1)^n \frac{x^{2n}}{(2n)!} + \dots + (-1)^{m-1} \frac{x^{2(m-1)}}{(2(m-1))!} \right| \\ &\leq \frac{A^{2n}}{(2n)!} + \dots + \frac{A^{2m-2}}{(2m-2)!} \\ &= \frac{A^{2n}}{(2n)!} \left[1 + \frac{(2n)!}{(2(n+1))!} A^2 + \frac{(2n)!}{(2(n+2))!} A^4 + \dots + \frac{(2n)!}{(2(m-1))!} A^{2(m-1-n)} \right] \\ &\leq \frac{A^{2n}}{(2n)!} \left[1 + \frac{A^2}{(2n)^2} + \frac{A^4}{(2n)^4} + \dots + \frac{A^{2(m-1-n)}}{(2n)^{2(m-1-n)}} \right] \\ &\leq \frac{A^{2n}}{(2n)!} \left[1 + \left(\frac{1}{4}\right)^2 + \left(\frac{1}{4}\right)^4 + \dots + \left(\frac{1}{4}\right)^{2(m-1-n)} \right] \\ &< \frac{16}{15} \cdot \frac{A^{2n}}{(2n)!} \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \frac{A^{2n}}{(2n)!} = 0$, Cauchy Criterion for Uniform Convergence implies C_n converges uniformly on $[-A, A]$, $\forall A > 0$

And hence, $C_n(x)$ converges $\forall x \in \mathbb{R}$.

$$\text{Let } C(x) = \lim_{n \rightarrow \infty} C_n(x).$$

Then C_n converges uniformly to C on $[-A, A]$, $\forall A > 0$.

Hence Thm 8.2.2 \Rightarrow

C is cts on $[-A, A]$, $\forall A > 0$

and therefore, C is cts on \mathbb{R} .

Moreover, $C_n(0) = 1 \Rightarrow C(0) = 1$.

$$\text{Since } S_n(x) = \int_0^x C_n(t) dt$$

$$S_m(x) - S_n(x) = \int_0^x (C_m(t) - C_n(t)) dt$$

$$\Rightarrow |S_m(x) - S_n(x)| \leq \int_0^x |C_m(t) - C_n(t)| dt \quad \text{if } x \geq 0$$

(Cor 7.3.15) $\left(\int_x^0 |C_m(t) - C_n(t)| dt, \text{ if } x < 0 \right)$

Then for $x \in [-A, A]$ & $m > n > 2A$,

$$|S_m(x) - S_n(x)| \leq \int_0^x \frac{16}{15} \cdot \frac{A^{2n}}{(2n)!} dt$$

$$\leq \frac{16}{15} \cdot \frac{A^{2n}}{(2n)!} \cdot A \quad \left(\text{similarly for } \int_x^0 \dots \right)$$

$$\rightarrow 0 \text{ as } n \rightarrow \infty$$

$\therefore S_n$ converges uniformly on $[-A, A]$, $\forall A > 0$.

$\Rightarrow S_n(x)$ converges $\forall x \in \mathbb{R}$.

Let $S(x) = \lim_{n \rightarrow \infty} S_n(x)$, $\forall x \in \mathbb{R}$

Then S_n converges uniformly to S on $[-A, A]$, $\forall A > 0$.

By Thm 8.2.2, S is cts on \mathbb{R} (as S_n cts on \mathbb{R} , $\forall n$)

Since $S_n(0) = 0$, we have $S(0) = 0$.

(To be cont'd)