


Quick Review on Exponential Function and Logarithmic Functions :

• $y = a^x$ with $a > 0$

Note: $y = a^x$ is well-defined when $a > 0$!

Think: If $a = -1$, when $x = \frac{1}{2}$, $y = a^x = \sqrt{-1}$!



graph of $y = a^x$ for

- 1) $a > 1$ 2) $a = 1$ 3) $0 < a < 1$

Properties :


1) a^x is positive for any $a > 0$ and any real number x .

2) If $a > 1$, $\lim_{x \rightarrow -\infty} a^x = 0$

$1 > a > 0$, $\lim_{x \rightarrow +\infty} a^x = 0$

• $y = \log_a x$ with $a > 1$ or $0 < a < 1$

Note: $y = \log_a x$ is well-defined when $a > 1$ or $0 < a < 1$!



graph of $y = \log_a x$ for

- 1) $a > 1$ 2) $0 < a < 1$

Properties :

1) By definition, if $y = a^x$, then $\log_a y = x$

$y > 0$

independent variable

dependent variable

$y = \log_a x \Rightarrow x > 0$

2) The graphs of $y = a^x$ and $y = \log_a x$ are symmetric along $y = x$.


It suffices to show if (p, q) lies on the graph of $y = a^x$, then (q, p) lies on the graph of $y = \log_a x$ and vice versa.

But it follows from the definition directly:

$$q = a^p \iff \log_a q = p$$

Note: reflecting $y = 1^x = 1$ along $y = x$ is $x = 1$, which is NOT a function.

It suggests why $\log_a x$ is NOT defined if $a = 1$.



Then situation when $a > 1$

(How about $0 < a < 1$?)

Differentiation of Exponential Functions:

Recall:

$$\frac{d}{dx} e^x = e^x$$

In particular, we write $\ln x$ instead of $\log_e x$, which is called **natural logarithm**.

Ex: Apply chain rule to show that $\frac{d}{dx} e^{ax} = ae^{ax}$.

Now, we write $a^x = e^{\ln a^x} = e^{(\ln a)x}$.

$$\text{So } \frac{d}{dx} a^x = \frac{d}{dx} e^{(\ln a)x} = (\ln a) e^{(\ln a)x} = a^x \ln a.$$

Note: If $a > 1$, $\frac{d}{dx} a^x = a^x \ln a > 0$ for all x , so $y = a^x$ is strictly increasing.


If $0 < a < 1$, $\frac{d}{dx} a^x = a^x \ln a < 0$ for all x , so $y = a^x$ is strictly decreasing.

Implicit Differentiation

e.g. $x^2 + y^2 = 2$ — \mathcal{C}

Locus of \mathcal{C} is a circle centered at $(0,0)$ with radius $\sqrt{2}$.

Check: $(1,1)$ is a point lying on the circle.





We want to find the equation of the tangent line l
(i.e. need to know the slope of l)

Note: $x^2 + y^2 = 2$ is NOT a function.

Question: How to find $\frac{dy}{dx}$? (and, actually, is it defined?)

Answer: Yes, roughly speaking,



The small segment of \mathcal{C} containing $(1,1)$ can be regarded as the graph of some function $y = g(x)$. (In fact, $y = \sqrt{2-x^2}$ in this case.)

How to find? Do it as usual!

e.g. $x^2 + y^2 = 2$

differentiate both sides with respect to x .

$$2x + \frac{d}{dx} y^2 = 0$$

$$2x + 2y \frac{dy}{dx} = 0 \quad (\text{Applying chain rule})$$

$$\frac{dy}{dx} = -\frac{x}{y}$$


$$\therefore \frac{dy}{dx} = -1 \quad \text{when } (x,y) = (1,1)$$

$$\text{We denote it by } \left. \frac{dy}{dx} \right|_{(x,y)=(1,1)} = -1$$

Remark :

$\frac{dy}{dx}$ is defined at a point of a curve only if a small arc containing the point can be regarded as the graph of some function $y=g(x)$.

$\therefore \frac{dy}{dx}$ is NOT defined when $(x,y) = (1,0)$ or $(-1,0)$.



No matter how small the arc is,


it cannot be realized as graph of some function $y=g(x)$.

e.g. $x^3 + y^3 - 3xy = 0$ — \mathcal{C}

$$3x^2 + 3y^2 \frac{dy}{dx} - 3y - 3x \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{y-x^2}{y^2-x}$$

If we want to find the slope of the tangent line at A .



putting $x=1$ into \mathcal{C} .

$$y^3 - 3y + 1 = 0$$

NOT easy to solve!

FACT: The above equation has three roots, two roots α_1, α_2 are positive ($\alpha_1 < \alpha_2$) one root is negative.

$A = (1, \alpha_1)$ and what we need is $\left. \frac{dy}{dx} \right|_{(x,y)=(1,\alpha_1)}$

Applications :

e.g. Differentiation of Logarithmic Function

Let $y = \ln x$, $x > 0$. Then $e^y = x$,

differentiate both sides with respect to x . $e^y \frac{dy}{dx} = 1$

$$e^y \frac{dy}{dx} = 1$$

$$\frac{dy}{dx} = \frac{1}{e^y} = \frac{1}{x}$$

$$\therefore \frac{d}{dx} \ln x = \frac{1}{x} \text{ for } x > 0.$$

Ex: By rewriting $\log_a x = \frac{\ln x}{\ln a}$, show that $\frac{d}{dx} \log_a x = \frac{1}{x \ln a}$.

e.g. Let $y = \ln|x|$, $x \neq 0$. Find $\frac{dy}{dx}$.

We can rewrite $y = \begin{cases} \ln x & \text{if } x > 0 \\ \ln(-x) & \text{if } x < 0 \end{cases}$

For $x > 0$, we have just shown that $\frac{dy}{dx} = \frac{1}{x}$



For $x < 0$, $y = \ln(-x)$

$$e^y = -x$$

$$e^y \frac{dy}{dx} = -1$$

$$\frac{dy}{dx} = \frac{-1}{e^y} = \frac{1}{x}$$

$$\therefore \frac{d}{dx} \ln|x| = \frac{1}{x} \text{ for } x \neq 0$$



Note: It is why $\int \frac{1}{x} dx = \ln|x| + C$.

↑
putting absolute sign here.

e.g. If $y = \sqrt[3]{\frac{(x-1)(x-2)^2}{x-4}}$, then find $\frac{dy}{dx}$.

Difficult to differentiate by using chain rule and quotient rule.

$$y^3 = \frac{(x-1)(x-2)^2}{x-4}$$

$$\ln y^3 = \ln \frac{(x-1)(x-2)^2}{x-4}$$

$$3 \ln y = \ln(x-1) + 2 \ln(x-2) - \ln(x-4)$$

$$\frac{3}{y} \frac{dy}{dx} = \frac{1}{x-1} + \frac{2}{x-2} - \frac{1}{x-4}$$

(Apply implicit differentiation)

$$\frac{dy}{dx} = \frac{y}{3} \left(\frac{1}{x-1} + \frac{2}{x-2} - \frac{1}{x-4} \right) = \frac{1}{3} \sqrt[3]{\frac{(x-1)(x-2)^2}{x-4}} \left(\frac{1}{x-1} + \frac{2}{x-2} - \frac{1}{x-4} \right)$$

Application of Differentiation:

e.g. Bacteria in closed environment with nutrition

What happens:

- Number of bacteria increases as plenty of nutrition at the beginning.
- Number of bacteria decreases as nutrition is no longer sufficient to support a large number of bacteria.

Suppose the number of bacteria t hours after the start of the experiment is modeled by the function

$$N(t) = \frac{10t+5}{e^{t+1}} \text{ (thousand)}, \quad t \geq 0.$$

① Number of bacteria at the beginning
 $= N(0) = 5/e \approx 1.84$ (thousand)

$$\textcircled{2} \quad N'(t) = \frac{10e^{t+1} - (10t+5)e^{t+1}}{e^{2(t+1)}}$$

$$= \frac{-10t+5}{e^{t+1}}$$

$$N'(t) > 0 \Leftrightarrow -10t+5 > 0 \Leftrightarrow t < 0.5$$

$$N'(t) < 0 \Leftrightarrow -10t+5 < 0 \Leftrightarrow t > 0.5$$

1st derivative check $\Rightarrow N(t)$ attains max. when $t = 0.5$

Max. number of bacteria = $N(0.5) = 10/e^{1.5} \approx 2.23$ (thousand)

$$\textcircled{3} \quad \lim_{t \rightarrow +\infty} N(t) = \lim_{t \rightarrow +\infty} \frac{1}{e} \frac{10t+5}{e^t}$$

$$= 0$$

Recall: If $p(x)$ is a polynomial,

then $\lim_{x \rightarrow +\infty} \frac{p(x)}{e^x} = 0$.

i.e. Extinct eventually!


$$\textcircled{4} \quad N''(t) = \frac{-10e^{t+1} - (-10t+5)e^{t+1}}{e^{2(t+1)}}$$

$$= \frac{10t-15}{e^{t+1}}$$

$$N''(t) > 0 \Leftrightarrow 10t-15 > 0 \Leftrightarrow t > 1.5$$

$$N''(t) < 0 \Leftrightarrow 10t-15 < 0 \Leftrightarrow t < 1.5$$

\therefore point of inflection = $(1.5, N(1.5)) \approx (1.5, 1.64)$



On the other hand, $N'(t)$ attains min when $t = 1.5$

$$N'(1.5) = -10/e^{1.5} \approx -2.23$$

i.e. Number of bacteria decreases most rapidly at $t = 1.5$ and decreasing rate at $t = 1.5$ is 2.23 thousand/hour.