

DIMENSION ESTIMATES FOR C^1 ITERATED FUNCTION SYSTEMS AND REPELLERS. PART I

DE-JUN FENG AND KÁROLY SIMON

ABSTRACT. This is the first article in a two-part series containing some results on dimension estimates for C^1 iterated function systems and repellers. In this part, we prove that the upper box-counting dimension of the attractor of any C^1 iterated function system (IFS) on \mathbb{R}^d is bounded above by its singularity dimension, and the upper packing dimension of any ergodic invariant measure associated with this IFS is bounded above by its Lyapunov dimension. Similar results are obtained for the repellers for C^1 expanding maps on Riemannian manifolds.

1. INTRODUCTION

The dimension theory of iterated function systems (IFS) and dynamical repellers has developed into an important field of research during the last 40 years. One of the main objectives is to estimate variant notions of dimension of the involved invariant sets and measures. Despite many new and significant developments in recent years, only the cases of conformal repellers and attractors of conformal IFSs under certain separation condition have been completely understood. In such cases, the topological pressure plays a crucial role in the theory. Indeed, the Hausdorff and box-counting dimensions of the repeller X for a C^1 conformal expanding map f are given by the unique number s satisfying the Bowen-Ruelle formula $P(X, f, -s \log \|D_x f\|) = 0$, where the functional P is the topological pressure, see [7, 38, 23]. A similar formula is obtained for the Hausdorff and box-counting dimensions of the attractor of a conformal IFS satisfying the open set condition (see e.g. [34]).

The study of dimension in the non-conformal cases has proved to be much more difficult. In his seminal paper [13], Falconer established a general upper bound on the Hausdorff and box-counting dimensions of a self-affine set (which is the attractor of an IFS consisting of contracting affine maps) in terms of the so-called affinity dimension, and proved that for typical self-affine sets under a mild assumption this upper bound is equal to the dimension. So far substantial progresses have been made towards understanding when the Hausdorff and box-counting dimensions of a concrete planar self-affine set are equal to its affinity dimension; see [3, 24] and the references therein. However very little has been known in the higher dimensional case.

In [15, Theorem 5.3], by developing a sub-additive version of the thermodynamical formalism, Falconer showed that the upper box-counting dimension of a mixing

repeller Λ of a non-conformal C^2 mapping ψ , under the following distortion condition

$$(1.1) \quad \|(D_x\psi)^{-1}\|^2\|D_x\psi\| < 1 \quad \text{for } x \in \Lambda,$$

is bounded above by the zero point of the (sub-additive) topological pressure associated with the singular value functions of the derivatives of iterates of ψ . We write this zero point as $\dim_{S^*} \Lambda$ and call it the *singularity dimension* of Λ (see Definition 6.1 for the detail). The condition (1.1) is used to prove the bounded distortion property of the singular value functions, which enables one to control the distortion of balls after many iterations (see [13, Lemma 5.2]). Examples involved “triangular maps” were constructed in [32] to show that the condition (1.1) is necessary for this bounded distortion property.

Using a quite different approach, Zhang [40] proved that the Hausdorff dimension of the repeller of an arbitrary C^1 expanding map ψ is also bounded above by the singularity dimension. (We remark that the upper bound given by Zhang was defined in a slightly different way, but it is equal to the singularity dimension; see [2, Corollary 2] for a proof.) The basic technique used in Zhang’s proof is to estimate the Hausdorff measure of $\psi(A)$ for small sets A (see [40, Lemma 3]), which is applied to only one iteration so it avoids assuming any distortion condition. However his method does not apply to the box-counting dimension.

Thanks to the results of Falconer and Zhang, it arises a natural question whether the upper box-counting dimension of a C^1 repeller is always bounded above by its singularity dimension. The challenge here is the lack of valid tools to analyse the geometry of the images of balls under a large number of iterations of C^1 maps. In [4, Theorem 3], Barreira made a positive claim to this question, but his proof contains a crucial mistake¹, as found by Manning and Simon [32]. In a recent paper [10, Theorem 3.2], Cao, Pesin and Zhao obtained an upper bound for the upper box-counting dimension of the repellers of $C^{1+\alpha}$ maps satisfying certain dominated splitting property. However that upper bound depends on the involved splitting and is usually strictly larger than the singularity dimension.

In the present paper, we give an affirmative answer to the above question. We also establish an analogous result for the attractors of C^1 non-conformal IFSs. Meanwhile we prove that the upper packing dimension of an ergodic invariant measure supported on a C^1 repeller (resp. the projection of an ergodic measure on the attractor of a C^1 IFS) is bound above by its Lyapunov dimension.

In the continuation of this paper [22] we verify that upper bound estimates of the dimensions of the attractors and ergodic measures in the previous paragraph, give the exact values of the dimensions for some families of C^2 non-conformal IFSs on the plane at least typically. Typically means that the assertions hold for almost all translations of the system. These families include the C^2 non-conformal IFSs on the plane for which all differentials are either diagonal matrices, or all differentials

¹This result was cited/applied in several papers (e.g., [2, Corollary 4], [11, Theorems 4.4-4.7]) without noticing the mistake in [4].

are lower triangular matrices and the contraction in y direction is stronger than in x direction.

We first state our results for C^1 IFSs. To this end, let us introduce some notation and definitions. Let Z be a compact subset of \mathbb{R}^d . A finite family $\{f_i\}_{i=1}^\ell$ of contracting self maps on Z is called a C^1 iterated function system (IFS), if there exists an open set $U \supset Z$ such that each f_i extends to a contracting C^1 -diffeomorphism $f_i : U \rightarrow f_i(U) \subset U$. Let K be the attractor of the IFS, that is, K is the unique non-empty compact subset of \mathbb{R}^d such that

$$(1.2) \quad K = \bigcup_{i=1}^{\ell} f_i(K)$$

(cf. [16]).

Let (Σ, σ) be the one-sided full shift over the alphabet $\{1, \dots, \ell\}$ (cf. [8]). Let $\Pi : \Sigma \rightarrow K$ denote the canonical coding map associated with the IFS $\{f_i\}_{i=1}^\ell$. That is,

$$(1.3) \quad \Pi(x) = \lim_{n \rightarrow \infty} f_{x_1} \circ \dots \circ f_{x_n}(z), \quad x = (x_n)_{n=1}^\infty,$$

with $z \in U$. The definition of Π is independent of the choice of z .

For any compact subset X of Σ with $\sigma X \subset X$, we call (X, σ) a *one-sided subshift* or simply *subshift* over $\{1, \dots, \ell\}$ and let $\dim_S X$ denote the singularity dimension of X with respect to the IFS $\{f_i\}_{i=1}^\ell$ (cf. Definition 2.4).

For a set $E \subset \mathbb{R}^d$, let $\overline{\dim}_B E$ denote the upper box-counting dimension of E (cf. [16]). The first result in this paper is the following, stating that the upper box-counting dimension of $\Pi(X)$ is bounded above by the singularity dimension of X .

Theorem 1.1. *Let $X \subset \Sigma$ be compact and $\sigma X \subset X$. Then $\overline{\dim}_B \Pi(X) \leq \dim_S X$. In particular,*

$$\overline{\dim}_B K \leq \dim_S \Sigma.$$

For an ergodic σ -invariant measure m on Σ , let $\dim_L m$ denote the Lyapunov dimension of m with respect to $\{f_i\}_{i=1}^\ell$ (cf. Definition 2.5). For a Borel probability measure η on \mathbb{R}^d (or a manifold), let $\overline{\dim}_P \eta$ denote the upper packing dimension of η . That is,

$$\overline{\dim}_P \eta = \text{esssup}_{x \in \text{supp}(\eta)} \overline{d}(\eta, x), \quad \text{with } \overline{d}(\eta, x) := \limsup_{r \rightarrow 0} \frac{\log \eta(B(x, r))}{\log r},$$

where $B(x, r)$ denotes the closed ball centered at x of radius r . Equivalently,

$$\overline{\dim}_P \eta = \inf \{ \dim_P F : F \text{ is a Borel set with } \eta(F) = 1 \},$$

where $\dim_P F$ stands for the packing dimension of F (cf. [16]). See e.g. [17] for a proof.

Our second result is the following, which can be viewed as a measure analogue of Theorem 1.1.

Theorem 1.2. *Let m be an ergodic σ -invariant measure on Σ , then*

$$\overline{\dim}_P(m \circ \Pi^{-1}) \leq \dim_L m,$$

where $m \circ \Pi^{-1}$ stands for the push-forward of m by Π .

The above theorem improves a result of Jordan and Pollicott [26, Theorem 1] which states that

$$\overline{\dim}_H(m \circ \Pi^{-1}) \leq \dim_L m$$

under a slightly more general setting, where $\overline{\dim}_H(m \circ \Pi^{-1})$ stands for the upper Hausdorff dimension of $m \circ \Pi^{-1}$. Recall that the upper Hausdorff dimension of a measure is the infimum of the Hausdorff dimension of Borel sets of full measure, which is always less than or equal to the upper packing dimension of the measure.

Next we turn to the case of repellers. Let \mathbf{M} be a smooth Riemannian manifold of dimension d and $\psi : \mathbf{M} \rightarrow \mathbf{M}$ a C^1 -map. Let Λ be a compact subset of \mathbf{M} such that $\psi(\Lambda) = \Lambda$. We say that ψ is *expanding* on Λ and Λ a *repeller* if

- (a) there exists $\lambda > 1$ such that $\|(D_z\psi)v\| \geq \lambda\|v\|$ for all $z \in \Lambda$, $v \in T_z\mathbf{M}$ (with respect to a Riemannian metric on \mathbf{M});
- (b) there exists an open neighborhood V of Λ such that

$$\Lambda = \{z \in V : \psi^n(z) \in V \text{ for all } n \geq 0\}.$$

We refer the reader to [35, Section 20] for more details. In what follows we always assume that Λ is a repeller of ψ . Let $\dim_{S^*}\Lambda$ denote the singular dimension of Λ with respect to ψ (see Definition 6.1). For an ergodic ψ -invariant measure μ on Λ , let $\dim_{L^*}\mu$ be the Lyapunov dimension of μ with respect to ψ (see Definition 6.2). Analogous to Theorems 1.1-1.2, we have the following.

Theorem 1.3. *Let Λ be the repeller of ψ . Then*

$$\overline{\dim}_B\Lambda \leq \dim_{S^*}\Lambda.$$

Theorem 1.4. *Let μ be an ergodic ψ -invariant measure supported on Λ . Then*

$$\overline{\dim}_P\mu \leq \dim_{L^*}\mu.$$

For the estimates of the box-counting dimension of attractors of C^1 non-conformal IFSs (resp. C^1 repellers), the reader may reasonably ask what difficulties arose in the previous work [15] which the present article overcomes. Below we give an explanation and illustrate roughly our strategy of the proof.

Let us give an account of the IFS case. The case of repellers is similar. Let K be the attractor of a C^1 IFS $\{f_i\}_{i=1}^\ell$. To estimate $\overline{\dim}_B K$, by definition one needs to estimate for given $r > 0$ the least number of balls of radius r required to cover K , say $N_r(K)$. To this end, one may iterate the IFS to get $K = \bigcup_{i_1 \dots i_n} f_{i_1 \dots i_n}(K)$ and then estimate $N_r(f_{i_1 \dots i_n}(K))$ separately, where $f_{i_1 \dots i_n} := f_{i_1} \circ \dots \circ f_{i_n}$. For this purpose, one needs to estimate $N_r(f_{i_1 \dots i_n}(B))$, where B is a fixed ball covering K .

Under the strong assumption of distortion property, Falconer was able to show that $f_{i_1 \dots i_n}(B)$ is roughly comparable to the ellipsoid $(D_x f_{i_1 \dots i_n})(B)$ for each $x \in B$

(see [15, Lemma 5.2]), then by cutting the ellipsoid into roughly round pieces he could use certain singular value function to give an upper bound of $N_r(f_{i_1 \dots i_n}(B))$, and then apply the sub-additive thermodynamic formalism to estimate the growth rate of $\sum_{i_1 \dots i_n} N_r(f_{i_1 \dots i_n}(B))$. However in the general C^1 non-conformal case, this approach is no longer feasible, since it seems hopeless to analyse the geometric shape of $f_{i_1 \dots i_n}(B)$ when n is large.

The strategy of our approach is quite different. We use an observation going back to Douady and Oesterlé [12] (see also [40]) that, for a given C^1 map f , when B_0 is an enough small ball in a fixed bounded region, then $f(B_0)$ is close to be an ellipsoid and so it can be covered by certain number of balls controlled by the singular values of the differentials of f (see Lemma 4.1). Since the maps f_i in the IFS are contracting, we may apply this fact to the maps $f_{i_n}, f_{i_{n-1}}, \dots, f_{i_1}$ recursively. Roughly speaking, suppose that B_0 is a ball of small radius r_0 , then $f_{i_n}(B_0)$ can be covered by N_1 balls of radius r_1 , and the image of each of them under $f_{i_{n-1}}$ can be covered by N_2 balls of radius r_2 , and so on, where $N_j, r_j/r_{j-1}$ ($j = 1, \dots, n$) can be controlled by the singular values of the differentials of $f_{i_{n-j+1}}$. In this way we get an estimate that $N_{r_n}(f_{i_1 \dots i_n}(B_0)) \leq N_1 \dots N_n$ (see Proposition 4.2 for a more precise statement), which is in spirit analogous to the corresponding estimate for the Hausdorff measure by Zhang [40]. In this process, we don't need to consider the differentials of $f_{i_1 \dots i_n}$ and so no distortion property is required. By developing a key technique from the thermodynamic formalism (see Proposition 3.4), we can get an upper bound for $\bar{\dim}_B K$, say s_1 . Replacing the IFS $\{f_i\}_{i=1}^\ell$ by its n -th iteration $\{f_{i_1 \dots i_n}\}$, we get other upper bounds s_n . Again using some technique in the thermodynamic formalism (see Proposition 3.1), we manage to show that $\liminf s_n$ is bounded above by the singularity dimension.

The paper is organized as follows. In Section 2, we give some preliminaries about the sub-additive thermodynamic formalism and give the definitions of the singularity and Lyapunov dimensions with respect to a C^1 IFS. In Section 3, we prove two auxiliary results (Propositions 3.1 and 3.4) which play a key role in the proof of Theorem 1.1 (and that of Theorem 1.3). The proofs of Theorems 1.1 and 1.2 are given in Sections 4-5, respectively. In Section 6, we give the definitions of the singularity and Lyapunov dimensions in the repeller case and prove Theorems 1.3-1.4. For the convenience of the reader, in the appendix we summarize the main notation and typographical conventions used in this paper.

2. PRELIMINARIES

2.1. Variational principle for sub-additive pressure. In order to define the singularity and Lyapunov dimensions and prove our main results, we require some elements from the sub-additive thermodynamic formalism.

Let (X, d) be a compact metric space and $T : X \rightarrow X$ a continuous mapping. We call (X, T) a *topological dynamical system*. For $x, y \in X$ and $n \in \mathbb{N}$, we define

$$(2.1) \quad d_n(x, y) := \max_{0 \leq i \leq n-1} d(T^i(x), T^i(y)).$$

A set $E \subset X$ is called (n, ε) -separated if for every distinct $x, y \in E$ we have $d_n(x, y) > \varepsilon$.

Let $C(X)$ denote the set of real-valued continuous functions on X . Let $\mathcal{G} = \{g_n\}_{n=1}^\infty$ be a *sub-additive potential* on X , that is, $g_n \in C(X)$ for all $n \geq 1$ such that

$$(2.2) \quad g_{m+n}(x) \leq g_n(x) + g_m(T^n x) \quad \text{for all } x \in X \text{ and } n, m \in \mathbb{N}.$$

Following [9], below we define the topological pressure of \mathcal{G} .

Definition 2.1. For given $n \in \mathbb{N}$ and $\varepsilon > 0$ we define

$$(2.3) \quad P_n(X, T, \mathcal{G}, \varepsilon) := \sup \left\{ \sum_{x \in E} \exp(g_n(x)) : E \text{ is an } (n, \varepsilon)\text{-separated set} \right\}.$$

Then the *topological pressure of \mathcal{G} with respect to T* is defined by

$$(2.4) \quad P(X, T, \mathcal{G}) := \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_n(X, T, \mathcal{G}, \varepsilon).$$

If the potential \mathcal{G} is additive, i.e. $g_n = S_n g := \sum_{k=0}^{n-1} g \circ T^k$ for some $g \in C(X)$, then $P(X, T, \mathcal{G})$ recovers the classical topological pressure $P(X, T, g)$ of g (see e.g. [39]).

Let $\mathcal{M}(X)$ denote the set of Borel probability measures on X , and $\mathcal{M}(X, T)$ the set of T -invariant Borel probability measures on X . For $\mu \in \mathcal{M}(X, T)$, let $h_\mu(T)$ denote the measure-theoretic entropy of μ with respect to T (cf. [39]). Moreover, for $\mu \in \mathcal{M}(X, T)$, by sub-additivity we have

$$(2.5) \quad \mathcal{G}_*(\mu) := \lim_{n \rightarrow \infty} \frac{1}{n} \int g_n d\mu = \inf_n \frac{1}{n} \int g_n d\mu \in [-\infty, \infty).$$

See e.g. [39, Theorem 10.1]. We call $\mathcal{G}_*(\mu)$ the *Lyapunov exponent* of \mathcal{G} with respect to μ .

The proofs of our main results rely on the following general variational principle for the topological pressure of sub-additive potentials.

Theorem 2.2 ([9], Theorem 1.1). *Let $\mathcal{G} = \{g_n\}_{n=1}^\infty$ be a sub-additive potential on a topological dynamical system (X, T) . Suppose that the topological entropy of T is finite. Then*

$$(2.6) \quad P(X, T, \mathcal{G}) = \sup \{h_\mu(T) + \mathcal{G}_*(\mu) : \mu \in \mathcal{M}(X, T)\}.$$

Particular cases of the above result, under stronger assumptions on the dynamical systems and the potentials, were previously obtained by many authors, see for example [14, 21, 18, 28, 33, 5] and references therein.

Measures that achieve the supremum in (2.6) are called *equilibrium measures* for the potential \mathcal{G} . There exists at least one ergodic equilibrium measure when the entropy map $\mu \mapsto h_\mu(T)$ is upper semi-continuous; this is the case when (X, T) is a subshift, see e.g. [19, Proposition 3.5] and the remark there.

The following well-known result is also needed in our proofs.

Lemma 2.3. *Let $X_i, i = 1, 2$ be compact metric spaces and let $T_i : X_i \rightarrow X_i$ be continuous. Suppose $\pi : X_1 \rightarrow X_2$ is a continuous surjection such that the following diagram commutes:*

$$\begin{array}{ccc} X_1 & \xrightarrow{T_1} & X_1 \\ \pi \downarrow & & \downarrow \pi \\ X_2 & \xrightarrow{T_2} & X_2 \end{array}$$

Then $\pi_ : \mathcal{M}(X_1, T_1) \rightarrow \mathcal{M}(X_2, T_2)$ (defined by $\mu \mapsto \mu \circ \pi^{-1}$) is surjective. If furthermore there is an integer $q > 0$ so that $\pi^{-1}(y)$ has at most q elements for each $y \in X_2$, then*

$$h_\mu(T_1) = h_{\mu \circ \pi^{-1}}(T_2)$$

for each $\mu \in \mathcal{M}(X_1, T_1)$.

Proof. The first part of the result is the same as [31, Chapter IV, Lemma 8.3]. The second part follows from the Abramov-Rokhlin formula (see [6]). \square

2.2. Subshifts. In this subsection, we introduce some basic notation and definitions about subshifts.

Let (Σ, σ) be the one-sided full shift over the alphabet $\mathcal{A} = \{1, \dots, \ell\}$. That is, $\Sigma = \mathcal{A}^{\mathbb{N}}$ endowed with the product topology, and $\sigma : \Sigma \rightarrow \Sigma$ is the left shift defined by $(x_i)_{i=1}^{\infty} \mapsto (x_{i+1})_{i=1}^{\infty}$. The topology of Σ is compatible with the following metric on Σ :

$$d(x, y) = 2^{-\inf\{k: x_{k+1} \neq y_{k+1}\}} \quad \text{for } x = (x_i)_{i=1}^{\infty}, y = (y_i)_{i=1}^{\infty}.$$

For $x = (x_i)_{i=1}^{\infty} \in \Sigma$ and $n \in \mathbb{N}$, write $x|n = x_1 \dots x_n$.

Let X be a non-empty compact subset of Σ satisfying $\sigma X \subset X$. We call (X, σ) a *one-sided subshift* or simply *subshift* over \mathcal{A} . We denote the collection of finite words allowed in X by X^* , and the subset of X^* of words of length n by X_n^* . In particular, define for $n \in \mathbb{N}$,

$$(2.7) \quad X^{(n)} := \left\{ (x_i)_{i=1}^{\infty} \in \mathcal{A}^{\mathbb{N}} : x_{kn+1}x_{kn+2} \dots x_{(k+1)n} \in X_n^* \text{ for all } k \geq 0 \right\}.$$

Let $\mathcal{G} = \{g_n\}_{n=1}^{\infty}$ be a sub-additive potential on a subshift (X, σ) . It is known that in such case, the topological pressure of \mathcal{G} can be alternatively defined by

$$(2.8) \quad P(X, \sigma, \mathcal{G}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_{\mathbf{i} \in X_n^*} \sup_{x \in [\mathbf{i}] \cap X} \exp(g_n(x)) \right),$$

where $[\mathbf{i}] := \{x \in \Sigma : x|n = \mathbf{i}\}$ for $\mathbf{i} \in \mathcal{A}^n$; see [9, p. 649]. The limit can be seen to exist by using a standard sub-additivity argument. We remark that (2.8) was first introduced by Falconer in [14] for the definition of the topological pressure of sub-additive potentials on a mixing repeller.

2.3. Singularity dimension and Lyapunov dimension with respect to C^1 IFSs. In this subsection, we define the singularity and Lyapunov dimensions with respect to C^1 IFSs. The corresponding definitions with respect to C^1 repellers will be given in Section 5.

Let $\{f_i\}_{i=1}^\ell$ be a C^1 IFS on \mathbb{R}^d with attractor K . Let (Σ, σ) be the one-sided full shift over the alphabet $\{1, \dots, \ell\}$ and let $\Pi : \Sigma \rightarrow K$ denote the corresponding coding map defined as in (1.3). For a differentiable function $f : U \subset \mathbb{R}^d \rightarrow \mathbb{R}^d$, let $D_z f$ denote the differential of f at $z \in U$.

For $T \in \mathbb{R}^{d \times d}$, let $\alpha_1(T) \geq \dots \geq \alpha_d(T)$ denote the singular values of T . Following [13], for $s \geq 0$ we define the *singular value function* $\phi^s : \mathbb{R}^{d \times d} \rightarrow [0, \infty)$ as

$$(2.9) \quad \phi^s(T) = \begin{cases} \alpha_1(T) \cdots \alpha_k(T) \alpha_{k+1}^{s-k}(T) & \text{if } 0 \leq s \leq d, \\ \det(T)^{s/d} & \text{if } s > d, \end{cases}$$

where $k = [s]$ is the integral part of s .

Definition 2.4. For a compact subset X of Σ with $\sigma(X) \subset X$, the *singularity dimension of X with respect to $\{f_i\}_{i=1}^\ell$* , written as $\dim_S X$, is the unique non-negative value s for which

$$P(X, \sigma, \mathcal{G}^s) = 0,$$

where $\mathcal{G}^s = \{g_n^s\}_{n=1}^\infty$ is the sub-additive potential on Σ defined by

$$(2.10) \quad g_n^s(x) = \log \phi^s(D_{\Pi \sigma^n x} f_{x|n}), \quad x \in \Sigma,$$

with $f_{x|n} := f_{x_1} \circ \dots \circ f_{x_n}$ for $x = (x_n)_{n=1}^\infty$.

Definition 2.5. Let m be an ergodic σ -invariant Borel probability measure on Σ . For any $i \in \{1, \dots, d\}$, the i -th Lyapunov exponent of m is

$$(2.11) \quad \lambda_i(m) := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log(\alpha_i(D_{\Pi \sigma^n x} f_{x|n})) \, dm(x).$$

The *Lyapunov dimension of m with respect to $\{f_i\}_{i=1}^\ell$* , written as $\dim_L m$, is the unique non-negative value s for which

$$h_m(\sigma) + \mathcal{G}_*^s(m) = 0,$$

where $\mathcal{G}_*^s = \{g_n^s\}_{n=1}^\infty$ is defined as in (2.10) and $\mathcal{G}_*^s(m) := \lim_{n \rightarrow \infty} \frac{1}{n} \int g_n^s \, dm$. See Figure 1 for the mapping $s \mapsto -\mathcal{G}_*^s(m)$ in the case when $d = 2$.

It follows from the definition of the singular value function ϕ^s that for an ergodic measure m we have

$$\mathcal{G}_*^s(m) = \begin{cases} \lambda_1(m) + \dots + \lambda_{[s]}(m) + (s - [s])\lambda_{[s]+1}(m), & \text{if } s < d \\ \frac{s}{d}(\lambda_1(m) + \dots + \lambda_d(m)), & \text{if } s \geq d. \end{cases}$$

Observe that in the special case when all the Lyapunov exponents are equal to the same λ then $\dim_L m = \frac{h_m(\sigma)}{-\lambda}$.

Remark 2.6. (i) The concept of singularity dimension was first introduced by Falconer [13, 15]; see also [29]. It is also called *affinity dimension* when the IFS $\{f_i\}_{i=1}^\ell$ is affine, i.e. each map f_i is affine.




FIGURE 1. The connection between Lyapunov dimension, entropy and the function $s \mapsto -\mathcal{G}_*^s(m)$ when $d = 2$.

- (ii) The definition of Lyapunov dimension of ergodic measures with respect to an IFS presented above was taken from [26]. It is a generalization of that given in [27] for affine IFSs.

2.4. A special consequence of Kingman's subadditive ergodic theorem. Here we state a special consequence of Kingman's subadditive ergodic theorem which will be needed in the proof of Lemma 5.1.

Lemma 2.7. *Let T be a measure preserving transformation of the probability space (X, \mathcal{B}, m) , and let $\{g_n\}_{n \in \mathbb{N}}$ be a sequence of L^1 functions satisfying the following sub-additivity relation:*

$$g_{n+m}(x) \leq g_n(x) + g_m(T^n x) \quad \text{for all } x \in X.$$

Suppose that there exists $C > 0$ such that

$$(2.12) \quad |g_n(x)| \leq Cn \quad \text{for all } x \in X \text{ and } n \in \mathbb{N}.$$

Then

$$\lim_{n \rightarrow \infty} E\left(\frac{g_n}{n} | \mathcal{C}_n\right)(x) = g(x) := \lim_{n \rightarrow \infty} \frac{g_n(x)}{n}$$

for m -a.e. x , where

$$\mathcal{C}_n := \{B \in \mathcal{B} : T^{-n}B = B \text{ a.e.}\},$$

$E(\cdot | \cdot)$ denotes the conditional expectation and $g(x)$ is T -invariant.

Proof. By Kingman's subadditive ergodic theorem, $\frac{g_n}{n}$ converges pointwisely to a T -invariant function g a.e. In the meantime, by Birkhoff's ergodic theorem, for each $n \in \mathbb{N}$,

$$E(g_n | \mathcal{C}_n)(x) = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{j=0}^{k-1} g_n(T^{jn} x) \text{ a.e.}$$

Since $\sum_{j=0}^{k-1} g_n(T^{jn}x) \geq g_{kn}(x)$ by subadditivity, it follows that

$$(2.13) \quad E\left(\frac{g_n}{n} \middle| \mathcal{C}_n\right)(x) \geq \lim_{k \rightarrow \infty} \frac{g_{nk}(x)}{nk} = g(x) \text{ a.e.}$$

Since the sequence $\{g_n\}$ is sub-additive and satisfies (2.12), by [28, Lemma 2.2], for any $0 < k < n$,

$$\frac{g_n(x)}{n} \leq \frac{1}{kn} \sum_{j=0}^{n-1} g_k(T^j x) + \frac{3kC}{n}, \quad x \in X.$$

As a consequence,

$$(2.14) \quad E\left(\frac{g_n}{n} \middle| \mathcal{C}_n\right)(x) \leq \frac{1}{kn} E\left(\sum_{j=0}^{n-1} g_k \circ T^j \middle| \mathcal{C}_n\right)(x) + \frac{3kC}{n}.$$

Notice that for each $f \in L^1$ and $n \in \mathbb{N}$,

$$(2.15) \quad E\left(\sum_{j=0}^{n-1} f \circ T^j \middle| \mathcal{C}_n\right) = nE(f | \mathcal{C}_1) \text{ a.e.}$$

To see the above identity, one simply applies Birkhoff's ergodic theorem (with respect to the transformations T^n and T , respectively) to the following limits

$$\lim_{k \rightarrow \infty} \frac{1}{k} \sum_{j=0}^{k-1} (f + f \circ T + \dots + f \circ T^{n-1})(T^{nj}x) = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{j=0}^{nk-1} f(T^j x).$$

Now applying the identity (2.15) (in which taking $f = g_k$) to (2.14) yields

$$E\left(\frac{g_n}{n} \middle| \mathcal{C}_n\right) \leq E\left(\frac{g_k}{k} \middle| \mathcal{C}_1\right) + \frac{3kC}{n} \text{ a.e. for } 0 < k < n.$$

It follows that

$$\limsup_{n \rightarrow \infty} E\left(\frac{g_n}{n} \middle| \mathcal{C}_n\right) \leq E\left(\frac{g_k}{k} \middle| \mathcal{C}_1\right) \text{ a.e. for each } k,$$

so by the dominated convergence theorem,

$$\limsup_{n \rightarrow \infty} E\left(\frac{g_n}{n} \middle| \mathcal{C}_n\right) \leq \lim_{k \rightarrow \infty} E\left(\frac{g_k}{k} \middle| \mathcal{C}_1\right) = E(g | \mathcal{C}_1) = g \text{ a.e.}$$

Combining it with (2.13) yields the desired result $\lim_{n \rightarrow \infty} E\left(\frac{g_n}{n} \middle| \mathcal{C}_n\right) = g$ a.e. \square

3. SOME AUXILIARY RESULTS

In this section we give two auxiliary results (Propositions 3.1 and 3.4) which are needed in the proof of Theorem 1.1.

Proposition 3.1. *Let (X, σ) be a one-sided subshift over a finite alphabet \mathcal{A} and $\mathcal{G} = \{g_n\}_{n=1}^{\infty}$ a sub-additive potential on $\mathcal{A}^{\mathbb{N}}$. Then*

$$(3.1) \quad P(X, \sigma, \mathcal{G}) = \lim_{n \rightarrow \infty} \frac{1}{n} P(X^{(n)}, \sigma^n, g_n) = \inf_{n \geq 1} \frac{1}{n} P(X^{(n)}, \sigma^n, g_n),$$

where $X^{(n)}$ is defined as in (2.7).

Remark 3.2. Instead of (3.1), it was proved in [2, Proposition 2.2] that

$$P(X, \sigma, \mathcal{G}) = \lim_{n \rightarrow \infty} \frac{1}{n} P(X, \sigma^n, g_n)$$

under a more general setting. We remark that the proof of (3.1) is more subtle.

To prove the Proposition 3.1, we need the following.

Lemma 3.3 ([9], Lemma 2.3). *Under the assumptions of Proposition 3.1, suppose that $\{\nu_n\}_{n=1}^\infty$ is a sequence in $\mathcal{M}(\mathcal{A}^\mathbb{N})$, where $\mathcal{M}(\mathcal{A}^\mathbb{N})$ denotes the space of all Borel probability measures on $\mathcal{A}^\mathbb{N}$ with the weak* topology. We form the new sequence $\{\mu_n\}_{n=1}^\infty$ by $\mu_n = \frac{1}{n} \sum_{i=0}^{n-1} \nu_n \circ \sigma^{-i}$. Assume that μ_{n_i} converges to μ in $\mathcal{M}(\mathcal{A}^\mathbb{N})$ for some subsequence $\{n_i\}$ of natural numbers. Then $\mu \in \mathcal{M}(\mathcal{A}^\mathbb{N}, \sigma)$, and moreover*

$$\limsup_{i \rightarrow \infty} \frac{1}{n_i} \int g_{n_i}(x) d\nu_{n_i}(x) \leq \mathcal{G}_*(\mu) := \lim_{n \rightarrow \infty} \frac{1}{n} \int g_n d\mu.$$

Proof of Proposition 3.1. We first prove that for each $n \in \mathbb{N}$,

$$(3.2) \quad P(X, \sigma, \mathcal{G}) \leq \frac{1}{n} P(X^{(n)}, \sigma^n, g_n).$$

To see this, fix $n \in \mathbb{N}$ and let μ be an equilibrium measure for the potential \mathcal{G} . Then

$$\begin{aligned} P(X, \sigma, \mathcal{G}) &= h_\mu(\sigma) + \mathcal{G}_*(\mu) \\ &\leq h_\mu(\sigma) + \frac{1}{n} \int g_n d\mu \quad (\text{by (2.5)}) \\ &= \frac{1}{n} \left(h_\mu(\sigma^n) + \int g_n d\mu \right) \\ &\leq \frac{1}{n} P(X^{(n)}, \sigma^n, g_n), \end{aligned}$$

where in the last inequality, we use the fact that $\mu \in \mathcal{M}(X^{(n)}, \sigma^n)$ and the classical variational principle for the topological pressure of additive potentials. This proves (3.2).

In what follows we prove that

$$(3.3) \quad P(X, \sigma, \mathcal{G}) \geq \limsup_{n \rightarrow \infty} \frac{1}{n} P(X^{(n)}, \sigma^n, g_n).$$

Clearly (3.2) and (3.3) imply (3.1). To prove (3.3), by the classical variational principle we can take a subsequence $\{n_i\}$ of natural numbers and $\nu_{n_i} \in \mathcal{M}(X^{(n_i)}, \sigma^{n_i})$ such that

$$(3.4) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} P(X^{(n)}, \sigma^n, g_n) = \lim_{i \rightarrow \infty} \frac{1}{n_i} \left(h_{\nu_{n_i}}(\sigma^{n_i}) + \int g_{n_i} d\nu_{n_i} \right).$$

Set $\mu^{(i)} = \frac{1}{n_i} \sum_{k=0}^{n_i-1} \nu_{n_i} \circ \sigma^{-k}$ for each i . Taking a subsequence if necessary we may assume that $\mu^{(i)}$ converges to an element $\mu \in \mathcal{M}(\mathcal{A}^\mathbb{N})$ in the weak* topology. By

Lemma 3.3, $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{N}}, \sigma)$ and moreover

$$(3.5) \quad \limsup_{i \rightarrow \infty} \frac{1}{n_i} \int g_{n_i}(x) d\nu_{n_i}(x) \leq \mathcal{G}_*(\mu).$$

Next we show further that μ is supported on X . For this we adopt some arguments from the proof of [30, Theorem 1.1]. Notice that for each i , $\mu^{(i)}$ is σ -invariant supported on

$$\bigcup_{k=0}^{n_i-1} \sigma^k X^{(n_i)} = \bigcup_{k=1}^{n_i} \sigma^{n_i-k} X^{(n_i)}.$$

Hence μ is supported on

$$\bigcap_{N=1}^{\infty} \overline{\bigcup_{i=N}^{\infty} \bigcup_{k=1}^{n_i} \sigma^{n_i-k} X^{(n_i)}}.$$

If x is in this set, then for each $N \geq 1$ there exist integers $i(N) \geq N$ and $k(N) \in [1, n_{i(N)}]$ for which $d(x, \sigma^{n_{i(N)}-k(N)} X^{(n_{i(N)})}) < 1/N$, hence

$$(3.6) \quad d(x, X) < \frac{1}{N} + \sup_{y \in \sigma^{n_{i(N)}-k(N)} X^{(n_{i(N)})}} d(y, X) \leq \frac{1}{N} + 2^{-k(N)}$$

and

$$(3.7) \quad d(\sigma^{k(N)} x, X) < \frac{2^{k(N)}}{N} + \sup_{y \in \sigma^{n_{i(N)}-k(N)} X^{(n_{i(N)})}} d(\sigma^{k(N)} y, X) \leq \frac{2^{k(N)}}{N} + 2^{-n_{i(N)}}.$$

If the values $k(N)$ are unbounded as $N \rightarrow \infty$, then (3.6) yields $x \in X$ while if they are bounded then some value of k recurs infinitely often as $k(N)$ which implies that $\sigma^k x \in X$ by (3.7). Thus μ is supported on

$$\bigcup_{k=0}^{\infty} \sigma^{-k} X.$$

Since $\sigma X \subset X$, the set $(\sigma^{-1} X) \setminus X$ is wandering under σ^{-1} (i.e. its preimages under powers of σ are disjoint), so it must have zero μ -measure. Consequently, $\mu \in \mathcal{M}(X, \sigma)$.

Notice that $h_{\mu^{(i)}}(\sigma) = \frac{1}{n_i} h_{\nu_{n_i}}(\sigma^{n_i})$ (see Lemma A.1). By the upper semi-continuity of the entropy map,

$$h_{\mu}(\sigma) \geq \limsup_{i \rightarrow \infty} h_{\mu^{(i)}}(\sigma) = \limsup_{i \rightarrow \infty} \frac{1}{n_i} h_{\nu_{n_i}}(\sigma^{n_i}),$$

which, together with (3.5), yields that

$$\begin{aligned} h_{\mu}(\sigma) + \mathcal{G}_*(\mu) &\geq \limsup_{i \rightarrow \infty} \frac{1}{n_i} \left(h_{\nu_{n_i}}(\sigma^{n_i}) + \int g_{n_i} d\nu_{n_i} \right) \\ &= \limsup_{n \rightarrow \infty} \frac{1}{n} P(X^{(n)}, \sigma^n, g_n). \end{aligned}$$

Applying Theorem 2.2, we obtain (3.3). This completes the proof of the proposition. \square

Next we present another auxiliary result.

Proposition 3.4. *Let (X, σ) be a one-sided subshift over a finite alphabet \mathcal{A} and $g, h \in C(X)$. Assume in addition that $h(x) < 0$ for all $x \in X$. Let*

$$r_0 = \sup_{x \in X} \exp(h(x)).$$

Set for $0 < r < r_0$,

$$\mathcal{A}_r := \left\{ i_1 \dots i_n \in X^* : \sup_{x \in [i_1 \dots i_n] \cap X} \exp(S_n h(x)) < r \leq \sup_{y \in [i_1 \dots i_{n-1}] \cap X} \exp(S_{n-1} h(y)) \right\},$$

where X^* is the collection of finite words allowed in X and $S_n h(x) := \sum_{k=0}^{n-1} h(\sigma^k x)$. Then

$$(3.8) \quad \lim_{r \rightarrow 0} \frac{\log \left(\sum_{I \in \mathcal{A}_r} \sup_{x \in [I] \cap X} \exp(S_{|I|} g(x)) \right)}{\log r} = -t,$$

where t is the unique real number so that $P(X, \sigma, g + th) = 0$, and $|I|$ stands for the length of I .

To prove the above result, we need the following.

Lemma 3.5. *Let (X, σ) be a one-sided subshift over a finite alphabet \mathcal{A} and $f \in C(X)$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sup \{ |S_n f(x) - S_n f(y)| : x_i = y_i \text{ for all } 1 \leq i \leq n \} = 0.$$

Proof. The result is well-known. For the reader's convenience, we include a proof.

Define for $n \in \mathbb{N}$,

$$\text{var}_n f = \sup \{ |f(x) - f(y)| : x_i = y_i \text{ for all } 1 \leq i \leq n \}.$$

Since f is uniformly continuous, $\text{var}_n f \rightarrow 0$ as $n \rightarrow \infty$. It follows that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \text{var}_i f = 0.$$

This concludes the result of the lemma since

$$\sup \{ |S_n f(x) - S_n f(y)| : x_i = y_i \text{ for all } 1 \leq i \leq n \}$$

is bounded above by $\sum_{i=1}^n \text{var}_i f$. □

Proof of Proposition 3.4. Set

$$\Theta_r = \sum_{I \in \mathcal{A}_r} \sup_{x \in [I] \cap X} \exp(S_{|I|} g(x)), \quad r \in (0, r_0).$$

Let $\epsilon > 0$. It is enough to show that

$$(3.9) \quad r^{-t+\epsilon} \leq \Theta_r \leq r^{-t-\epsilon}$$

for sufficiently small r .

To this end, set for $0 < r < r_0$,

$$m(r) = \min\{|I| : I \in \mathcal{A}_r\}, \quad M(r) = \max\{|I| : I \in \mathcal{A}_r\}.$$

From the definition of \mathcal{A}_r and the negativity of h , it follows that there exist two positive constants a, b such that

$$(3.10) \quad a \log(1/r) \leq m(r) \leq M(r) \leq b \log(1/r), \quad \forall r \in (0, r_0).$$

Define

$$\Gamma_r = \sum_{I \in \mathcal{A}_r} \sup_{x \in [I] \cap X} \exp(S_{|I|}(g + th)(x)), \quad r \in (0, r_0).$$

By Lemma 3.5 and (3.10), it is readily checked that

$$(3.11) \quad r^{t+\epsilon/2} \Theta_r \leq \Gamma_r \leq r^{t-\epsilon/2} \Theta_r \quad \text{for sufficiently small } r,$$

Hence to prove (3.9), it suffices to prove that $r^{\epsilon/2} \leq \Gamma_r \leq r^{-\epsilon/2}$ for small r .

We first prove $\Gamma_r > r^{\epsilon/2}$ when r is small. Suppose on the contrary this is not true. Then by (3.10) we can find some $r > 0$ and $\lambda > 0$ such that $Z(r, \lambda) < 1$, where

$$(3.12) \quad Z(r, \lambda) := \sum_{I \in \mathcal{A}_r} \exp(\lambda|I|) \sup_{x \in [I] \cap X} \exp(S_{|I|}(g + th)(x)).$$

From Lemma 2.14 of [8] it follows that $P(X, \sigma, g + th) \leq -\lambda$, contradicting the fact that $P(X, \sigma, g + th) = 0$. Hence we have $\Gamma_r > r^{\epsilon/2}$ when r is sufficiently small.

Next we prove the inequality $\Gamma_r \leq r^{-\epsilon/2}$ for small r . To do this, fix $\lambda \in (0, \epsilon/(2b))$, where b is the constant in (3.10). We claim that there exists $0 < r_1 < r_0$ such that

$$(3.13) \quad Z(r, -\lambda) < 1 \quad \text{for all } r \in (0, r_1),$$

where Z is defined as in (3.12). Since $\lambda \in (0, \epsilon/(2b))$, it follows from (3.10) that $\exp(-\lambda|I|) \geq r^{\epsilon/2}$ for any $I \in \mathcal{A}_r$. Hence (3.13) implies that $\Gamma_r \leq r^{-\epsilon/2}$ for $0 < r < r_1$.

Now it remains to prove (3.13). Since $P(X, \sigma, g + th) = 0$, by definition we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_{I \in X^* : |I|=n} \sup_{x \in [I] \cap X} \exp(S_{|I|}(g + th)(x)) \right) = 0.$$

Hence there exists a large N such that $e^{-\lambda N/2} < 1 - e^{-\lambda/2}$ and for any $n > N$,

$$\gamma_n := \sum_{I \in X^* : |I|=n} \exp(-\lambda|I|) \sup_{x \in [I] \cap X} \exp(S_{|I|}(g + th)(x)) \leq \exp(-\lambda n/2).$$

Take a small $r_1 \in (0, r_0)$ so that $m(r) \geq N$ for any $0 < r < r_1$. Hence by (3.10), for any $0 < r < r_1$, $\mathcal{A}_r \subset \{I \in X^* : |I| \geq N\}$ and so

$$Z(r, -\lambda) \leq \sum_{n=N}^{\infty} \gamma_n \leq \sum_{n=N}^{\infty} \exp(-\lambda n/2) = \frac{e^{-\lambda N/2}}{1 - e^{-\lambda/2}} < 1.$$

This proves (3.13) and we are done. □

4. THE PROOF OF THEOREM 1.1

Recall that for $T \in \mathbb{R}^{d \times d}$, $\alpha_1(T) \geq \dots \geq \alpha_d(T)$ are the singular values of T , and $\phi^s(T)$ ($s \geq 0$) is defined as in (2.9). We begin with an elementary but important lemma.

Lemma 4.1. *Let $E \subset U \subset \mathbb{R}^d$, where E is compact and U is open. Let $k \in \{0, 1, \dots, d-1\}$. Then for any non-degenerate C^1 map $f : U \rightarrow \mathbb{R}^d$, there exists $r_0 > 0$ so that for any $y \in E$, $z \in B(y, r_0)$ and $0 < r < r_0$, the set $f(B(z, r))$ can be covered by*

$$(2d)^d \cdot \frac{\phi^k(D_y f)}{(\alpha_{k+1}(D_y f))^k}$$

many balls of radius $\alpha_{k+1}(D_y f)r$.

Proof. The result was implicitly proved in [40, Lemma 3] by using an idea of [12]. For the convenience of the reader, we provide a detailed proof.

Set

$$\gamma = \min_{y \in E} \alpha_d(D_y f).$$

Then

$$(4.1) \quad B(0, \gamma) \leq D_y f(B(0, 1)) \text{ for all } y \in E.$$

Since f is C^1 , non-degenerate on U and E is compact, it follows that $\gamma > 0$. Take $\epsilon = (\sqrt{d} - 1)/2$. Then there exists a small $r_0 > 0$ such that for $u, v, w \in V_{2r_0}(E) := \{x : d(x, E) < 2r_0\}$,

$$(4.2) \quad |f(u) - f(v) - D_v f(u - v)| \leq \epsilon \gamma |u - v| \quad \text{if } |u - v| \leq r_0,$$

and

$$(4.3) \quad D_v f(B(0, 1)) \subset ((1 + \epsilon)D_w f)(B(0, 1)) \quad \text{if } |v - w| \leq r_0.$$

Now let $y \in E$ and $z \in B(y, r_0)$. For any $0 < r < r_0$ and $x \in B(z, r)$, taking $u = x$ and $v = z$ in (4.2) gives

$$f(x) - f(z) - D_z f(x - z) \in B(0, \epsilon \gamma r),$$

so by (4.3) and (4.1),

$$\begin{aligned} f(x) - f(z) &\in D_z f(B(0, r)) + B(0, \epsilon \gamma r) \\ &\subset ((1 + \epsilon)D_y f)(B(0, r)) + B(0, \epsilon \gamma r) && \text{(by (4.3))} \\ &\subset D_y f(B(0, (1 + \epsilon)r)) + B(0, \epsilon r) && \text{(by (4.1))} \\ &\subset D_y f(B(0, (1 + 2\epsilon)r)) \\ &= D_y f\left(B\left(0, \sqrt{d}r\right)\right), \end{aligned}$$

where $A + A' := \{u + v : u \in A, v \in A'\}$. Therefore

$$f(B(z, r)) \subset f(z) + D_y f\left(B\left(0, \sqrt{d}r\right)\right).$$

That is, $f(B(z, r))$ is contained in an ellipsoid which has principle axes of lengths $2\sqrt{d}\alpha_i(D_y f)r$, $i = 1, \dots, d$. Hence $f(B(z, r))$ is contained in a rectangular parallelepiped of side lengths $2\sqrt{d}\alpha_i(D_y f)r$, $i = 1, \dots, d$. Now we can divide such a parallelepiped into at most

$$\left(\prod_{i=1}^{k+1} \frac{2d\alpha_i(D_y f)}{\alpha_{k+1}(D_y f)} \right) \cdot (2d)^{d-k-1} \leq (2d)^d \cdot \frac{\phi^k(D_y f)}{(\alpha_{k+1}(D_y f))^k}$$

cubes of side $\frac{2}{\sqrt{d}} \cdot \alpha_{k+1}(D_y f)r$. Therefore this parallelepiped (and $f(B(z, r))$ as well) can be covered by

$$(2d)^d \cdot \frac{\phi^k(D_y f)}{(\alpha_{k+1}(D_y f))^k}$$

balls of radius $\alpha_{k+1}(D_y f)r$. □

In the remaining part of this section, let $\{f_i\}_{i=1}^\ell$ be a C^1 IFS on \mathbb{R}^d with attractor K . Let (Σ, σ) be the one-sided full shift over the alphabet $\{1, \dots, \ell\}$ and $\Pi : \Sigma \rightarrow K$ the canonical coding map associated with the IFS (cf. (1.3)). As a consequence of Lemma 4.1, we obtain the following.

Proposition 4.2. *Let $k \in \{0, 1, \dots, d-1\}$. Set $C = (2d)^d$. Then there exists $C_1 > 0$ such that for $\mathbf{i} = (i_p)_{p=1}^\infty \in \Sigma$ and $n \in \mathbb{N}$, the set $f_{\mathbf{i}|n}(K)$ can be covered by $C_1 \prod_{p=0}^{n-1} G(\sigma^p \mathbf{i})$ balls of radius $\prod_{p=0}^{n-1} H(\sigma^p \mathbf{i})$, where*

$$(4.4) \quad G(\mathbf{i}) := \frac{C\phi^k(D_{\Pi\sigma\mathbf{i}}f_{i_1})}{\alpha_{k+1}(D_{\Pi\sigma\mathbf{i}}f_{i_1})^k}, \quad H(\mathbf{i}) := \alpha_{k+1}(D_{\Pi\sigma\mathbf{i}}f_{i_1}).$$

Proof. Since $\{f_i\}_{i=1}^\ell$ is a C^1 IFS, there exists an open set $U \supset K$ such that each f_i extends to a C^1 diffeomorphism $f_i : U \rightarrow f_i(U)$. Applying Lemma 4.1 to the mappings f_i , we see that there exists $r_0 > 0$ such that for any $y \in K$, $z \in B(y, r_0)$, $0 < r < r_0$ and $i \in \{1, \dots, \ell\}$, the set $f_i(B(z, r))$ can be covered by

$$\theta(y, i) := \frac{C\phi^k(D_y f_i)}{\alpha_{k+1}(D_y f_i)^k}$$

balls of radius $\alpha_{k+1}(D_y f_i)r$.

Since f_1, \dots, f_ℓ are contracting on U , there exists $\gamma \in (0, 1)$ such that

$$|f_i(x) - f_i(y)| \leq \gamma|x - y| \quad \text{for all } x, y \in U, i \in \{1, \dots, \ell\}.$$

It implies that $\alpha_1(D_y f_i) \leq \gamma$ for any $y \in K$ and $i \in \{1, \dots, \ell\}$. Take a large integer n_0 so that

$$\gamma^{n_0} \max\{1, \text{diam}(K)\} < r_0/2.$$

Clearly there exists a large number C_1 so that the conclusion of the proposition holds for any positive integer $n \leq n_0$ and $\mathbf{i} \in \Sigma$, i.e. the set $f_{\mathbf{i}|n}(K)$ can be covered by $C_1 \prod_{p=0}^{n-1} G(\sigma^p \mathbf{i})$ balls of radius $\prod_{p=0}^{n-1} H(\sigma^p \mathbf{i})$. Below we show by induction that this holds for all $n \in \mathbb{N}$ and $\mathbf{i} \in \Sigma$.

Suppose for some $m \geq n_0$, the conclusion of the proposition holds for any positive integer $n \leq m$ and $\mathbf{i} \in \Sigma$. Then for $\mathbf{i} \in \Sigma$, $f|_{(\sigma\mathbf{i})|_m(K)}$ can be covered by $C_1 \prod_{p=0}^{m-1} G(\sigma^{p+1}\mathbf{i})$ balls of radius $\prod_{p=0}^{m-1} H(\sigma^{p+1}\mathbf{i})$. Let B_1, \dots, B_N denote these balls. We may assume that $B_j \cap f|_{(\sigma\mathbf{i})|_m(K)} \neq \emptyset$ for each j . Since

$$\prod_{p=0}^{m-1} H(\sigma^{p+1}\mathbf{i}) \leq \gamma^m \leq \gamma^{n_0} < r_0/2$$

and

$$d(\Pi\sigma\mathbf{i}, B_j \cap f|_{(\sigma\mathbf{i})|_m(K)}) \leq \text{diam}(f|_{(\sigma\mathbf{i})|_m(K)}) \leq \gamma^{n_0} \text{diam}(K) < r_0/2,$$

so the center of B_j is in $B(\Pi\sigma\mathbf{i}, r_0)$. Therefore $f_{i_1}(B_j)$ can be covered by $\theta(\Pi\sigma\mathbf{i}, i_1) = G(\mathbf{i})$ balls of radius

$$H(\mathbf{i}) \cdot \prod_{p=0}^{m-1} H(\sigma^{p+1}\mathbf{i}) = \prod_{p=0}^m H(\sigma^p\mathbf{i}).$$

Since $f_{i_1(m+1)}(K) \subset \bigcup_{j=1}^N f_{i_1}(B_j)$, it follows that $f_{i_1(m+1)}(K)$ can be covered by

$$G(\mathbf{i})N \leq G(\mathbf{i}) \cdot C_1 \prod_{p=0}^{m-1} G(\sigma^{p+1}\mathbf{i}) = C_1 \prod_{p=0}^m G(\sigma^p\mathbf{i})$$

balls of radius $\prod_{p=0}^m H(\sigma^p\mathbf{i})$. Thus the proposition also holds for $n = m + 1$ and all $\mathbf{i} \in \Sigma$, as desired. \square

Next we provide an upper bound of the upper box-counting dimension of the attractor K of the IFS $\{f_i\}_{i=1}^\ell$.

Proposition 4.3. *Let $k \in \{0, 1, \dots, d-1\}$. Let $G, H : \Sigma \rightarrow \mathbb{R}$ be defined as in (4.4). Let t be the unique real number so that*

$$P(\Sigma, \sigma, (\log G) + t(\log H)) = 0.$$

Then $\overline{\dim}_B K \leq t$.

Proof. For short, we write $g = \log G$ and $h = \log H$. Define

$$r_{\min} = \min_{x \in \Sigma} \alpha_{k+1}(D_{\Pi\sigma x} f_{x_1}), \quad r_{\max} = \max_{x \in \Sigma} \alpha_{k+1}(D_{\Pi\sigma x} f_{x_1}).$$

Then $0 < r_{\min} \leq r_{\max} < 1$. For $0 < r < r_{\min}$, define

$$(4.5) \quad \mathcal{A}_r = \left\{ i_1 \dots i_n \in \Sigma^* : \sup_{x \in [i_1 \dots i_n]} S_n h(x) < \log r \leq \sup_{y \in [i_1 \dots i_{n-1}]} S_{n-1} h(y) \right\};$$

clearly $\{[I] : I \in \mathcal{A}_r\}$ is a partition of Σ . By Proposition 4.2, there exists a constant $C_1 > 0$ such that for each $0 < r < r_{\min}$, every $I \in \mathcal{A}_r$ and $x \in [I]$, $f_I(K)$ can be covered by

$$C_1 \exp(S_{|I|} g(x)) \leq C_1 \exp\left(\sup_{y \in [I]} S_{|I|} g(y)\right)$$

many balls of radius

$$\exp(S_{|I|}h(x)) \leq \exp\left(\sup_{y \in [I]} S_{|I|}h(y)\right) < r.$$

It follows that K can be covered by

$$C_1 \sum_{I \in \mathcal{A}_r} \exp\left(\sup_{y \in [I]} S_{|I|}g(y)\right)$$

many balls of radius r . Hence by Proposition 3.4,

$$\overline{\dim}_B K \leq \limsup_{r \rightarrow 0} \frac{\log\left(\sum_{I \in \mathcal{A}_r} \exp\left(\sup_{y \in [I]} S_{|I|}g(y)\right)\right)}{\log(1/r)} = t.$$

This completes the proof of the proposition. \square

As an application of Proposition 4.3, we may estimate the upper box-counting dimension of the projections of a class of σ -invariant sets under the coding map.

Proposition 4.4. *Let X be a compact subset of Σ satisfying $\sigma X \subset X$, and $k \in \{0, \dots, d-1\}$. Then for each $n \in \mathbb{N}$,*

$$\overline{\dim}_B \Pi(X^{(n)}) \leq t_n,$$

where $X^{(n)}$ is defined as in (2.7), t_n is the unique number for which

$$P(X^{(n)}, \sigma^n, (\log G_n) + t_n(\log H_n)) = 0,$$

and G_n, H_n are continuous functions on Σ defined by

$$(4.6) \quad G_n(y) := \frac{C\phi^k(D_{\Pi\sigma^n y} f_{y|n})}{\alpha_{k+1}(D_{\Pi\sigma^n y} f_{y|n})^k}, \quad H_n(y) := \alpha_{k+1}(D_{\Pi\sigma^n y} f_{y|n}),$$

with $C = (2d)^d$.

Proof. The result is obtained by applying Proposition 4.3 to the IFS $\{f_I : I \in X_n^*\}$ instead of $\{f_i\}_{i=1}^\ell$, where X_n^* stands for the collection of words of length n allowed in X . \square

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Write $s = \dim_S X$. We may assume that $s < d$; otherwise we have nothing left to prove. Set $k = [s]$, i.e. k is the largest integer less than or equal to s . Let $\mathcal{U} = \{u_n\}_{n=1}^\infty$ be the sub-additive potential on Σ defined by

$$u_n(x) = \log \phi^s(D_{\Pi\sigma^n x} f_{x|n}).$$

Then $P(X, \sigma, \mathcal{U}) = 0$ by the definition of $\dim_S X$. Hence by Proposition 3.1,

$$\lim_{n \rightarrow \infty} \frac{1}{n} P(X^{(n)}, \sigma^n, u_n) = \inf_{n \geq 1} \frac{1}{n} P(X^{(n)}, \sigma^n, u_n) = 0.$$

It follows that for each $\epsilon > 0$, there exists $N_\epsilon > 0$ such that

$$(4.7) \quad 0 \leq P(X^{(n)}, \sigma^n, u_n) \leq n\epsilon \quad \text{for } n > N_\epsilon.$$

For $n \in \mathbb{N}$, let s_n be the unique real number so that $P(X^{(n)}, \sigma^n, v_n) = 0$, where v_n is a continuous function on Σ defined by

$$\begin{aligned} v_n(x) &= \log G_n(x) + s_n \log H_n(x) \\ &= \log (C \phi^k(D_{\Pi \sigma^n x} f_{x|n}) \alpha_{k+1}(D_{\Pi \sigma^n x} f_{x|n})^{s_n - k}), \end{aligned}$$

where G_n, H_n are defined in (4.6) and $C = (2d)^d$. By Proposition 4.4,

$$\overline{\dim}_B \Pi(X) \leq \overline{\dim}_B \Pi(X^{(n)}) \leq s_n.$$

If $s \geq s_n$ for some n , then $\overline{\dim}_B \Pi(X) \leq s_n \leq s$ and we are done. In what follows, we assume that $s < s_n$ for each n . Then for each $x \in \Sigma$,

$$\begin{aligned} u_n(x) - v_n(x) &= -\log C + (s - s_n) \log \alpha_{k+1}(D_{\Pi \sigma^n x} f_{x|n}) \\ &\geq -\log C + (s - s_n) \log \alpha_1(D_{\Pi \sigma^n x} f_{x|n}) \\ &\geq -\log C + n(s - s_n) \log \theta, \end{aligned}$$

where

$$\theta := \max_{y \in \Sigma} \alpha_1(D_{\Pi \sigma y} f_{y_1}) < 1.$$

Hence

$$\begin{aligned} P(X^{(n)}, \sigma^n, u_n) &= P(X^{(n)}, \sigma^n, u_n) - P(X^{(n)}, \sigma^n, v_n) \\ &\geq \inf_{x \in \Sigma} (u_n(x) - v_n(x)) \\ &\geq -\log C + n(s - s_n) \log \theta, \end{aligned}$$

where in the second inequality, we used [39, Theorem 9.7(iv)]. Combining it with (4.7) yields that for $n \geq N_\epsilon$,

$$n\epsilon \geq -\log C + n(s - s_n) \log \theta,$$

so

$$s \geq s_n + \frac{\epsilon + n^{-1} \log C}{\log \theta} \geq \overline{\dim}_B \Pi(X) + \frac{\epsilon + n^{-1} \log C}{\log \theta}.$$

Letting $n \rightarrow \infty$ and then $\epsilon \rightarrow 0$, we obtain $s \geq \overline{\dim}_B \Pi(X)$, as desired. \square

5. THE PROOF OF THEOREM 1.2

Let $\Pi : \Sigma \rightarrow \mathbb{R}^d$ be the coding map associated with a C^1 IFS $\{f_i\}_{i=1}^\ell$ on \mathbb{R}^d (cf. (1.3)). For $E \subset \mathbb{R}^d$ and $\delta > 0$, let $N_\delta(E)$ denote the smallest integer N for which E can be covered by N closed balls of radius δ . For $T \in \mathbb{R}^{d \times d}$, let $\alpha_1(T) \geq \dots \geq \alpha_d(T)$ denote the singular values of T , and let $\phi^s(T)$ be the singular value function defined as in (2.9).

The following geometric counting lemma plays an important role in the proof of Theorem 1.2. It is of independent interest as well.

Lemma 5.1. *Let m be an ergodic σ -invariant Borel probability measure on Σ . Set*

$$(5.1) \quad \lambda_i := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log (\alpha_i(D_{\Pi\sigma^n x} f_{x|n})) \, dm(x), \quad i = 1, \dots, d.$$

Let $k \in \{0, \dots, d-1\}$. Write $u := \exp(\lambda_{k+1})$. Then for m -a.e. $x \in \Sigma$,

$$(5.2) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \log N_{u^n}(\Pi([x|n])) \leq (\lambda_1 + \dots + \lambda_k) - k\lambda_{k+1}.$$

Proof. It is known (see, e.g. [1, Theorem 3.3.3]) that for m -a.e. x ,

$$\lim_{p \rightarrow \infty} \frac{1}{p} \log (\alpha_i(D_{\Pi\sigma^p x} f_{x|p})) = \lambda_i, \quad i = 1, \dots, d,$$

and

$$\lim_{p \rightarrow \infty} \frac{1}{p} \log (\phi^i(D_{\Pi\sigma^p x} f_{x|p})) = \lambda_1 + \dots + \lambda_i, \quad i = 1, \dots, d.$$

For $i \in \{1, \dots, d\}$, $x \in \Sigma$ and $p \in \mathbb{N}$, set

$$\begin{aligned} w_p^{(i)}(x) &= \log \phi^i(D_{\Pi\sigma^p x} f_{x|p}), \\ v_p^{(i)}(x) &= \log \alpha_i(D_{\Pi\sigma^p x} f_{x|p}). \end{aligned}$$

Then by the definition of ϕ^i ,

$$(5.3) \quad w_p^{(i)} = v_p^{(1)} + \dots + v_p^{(i)}.$$

Since ϕ^i is sub-multiplicative on $\mathbb{R}^{d \times d}$ (cf. [13, Lemma 2.1]), $\{w_p^{(i)}\}_{p=1}^{\infty}$ is a sub-additive potential satisfying

$$|w_p^{(i)}(x)| \leq pC, \quad p \in \mathbb{N}, \quad x \in \Sigma,$$

for some constant $C > 0$. Set $\mathcal{C}_p := \{B \in \mathcal{B}(\Sigma) : \sigma^{-p}B = B \text{ a.e.}\}$. Then by Lemma 2.7, for m -a.e. x ,

$$(5.4) \quad \lim_{p \rightarrow \infty} E \left(\frac{w_p^{(i)}}{p} \middle| \mathcal{C}_p \right) (x) = \lim_{p \rightarrow \infty} \frac{1}{p} w_p^{(i)}(x) = \lambda_1 + \dots + \lambda_i, \quad i = 1, \dots, d,$$

and so by (5.3),

$$(5.5) \quad \lim_{p \rightarrow \infty} E \left(\frac{v_p^{(i)}}{p} \middle| \mathcal{C}_p \right) (x) = \lambda_i, \quad i = 1, \dots, d,$$

Let $p \in \mathbb{N}$. Applying Proposition 4.2 to the IFS $\{f_I : I \in \mathcal{A}^p\}$, we see that there exists a positive number $C_1(p)$ such that for any $x \in \Sigma$ and $n \in \mathbb{N}$, the set $\Pi([x|np]) = f_{x|np}(K)$ can be covered by $C_1(p) \prod_{i=0}^{n-1} G_p(\sigma^{pi}x)$ balls of radius $\prod_{i=0}^{n-1} H_p(\sigma^{ip}x)$, where

$$(5.6) \quad G_p(x) := \frac{C\phi^k(D_{\Pi\sigma^p x} f_{x|p})}{\alpha_{k+1}(D_{\Pi\sigma^p x} f_{x|p})^k}, \quad H_p(x) := \alpha_{k+1}(D_{\Pi\sigma^p x} f_{x|p}),$$

with $C = (2d)^d$. By the Birkhoff's ergodic theorem, for m -a.e. $x \in \Sigma$,

$$(5.7) \quad \lim_{n \rightarrow \infty} \frac{1}{np} \log \left(C_1(p) \prod_{i=0}^{n-1} G_p(\sigma^{pi} x) \right) = \frac{\log C}{p} + E \left(\frac{w_p^{(k)}}{p} \middle| \mathcal{C}_p \right) (x) - kE \left(\frac{v_p^{(k+1)}}{p} \middle| \mathcal{C}_p \right) (x)$$

and

$$(5.8) \quad \lim_{n \rightarrow \infty} \frac{1}{np} \log \left(\prod_{i=0}^{n-1} H_p(\sigma^{ip} x) \right) = E \left(\frac{v_p^{(k+1)}}{p} \middle| \mathcal{C}_p \right) (x),$$

Let $\epsilon > 0$. By (5.4), (5.5), (5.7), (5.8), for m -a.e. x there exists a positive integer $p_0(x)$ such that for any $p \geq p_0(x)$,

$$(5.9) \quad \prod_{i=0}^{n-1} H_p(\sigma^{ip} x) \leq (u + \epsilon)^{np} \quad \text{for large enough } n,$$

and

$$(5.10) \quad \frac{1}{np} \log \left(C_1(p) \prod_{i=0}^{n-1} G_p(\sigma^{pi} x) \right) \leq \lambda_1 + \dots + \lambda_k - k\lambda_{k+1} + \epsilon \quad \text{for large enough } n,$$

Fix such an x and let $p \geq p_0(x)$. By (5.9),

$$N_{(u+\epsilon)^{np}}(\Pi([x|pn])) \leq C_1(p) \prod_{i=0}^{n-1} G_p(\sigma^{pi} x) \quad \text{for large enough } n.$$

Notice that there exists a constant $C_2 = C_2(d) > 0$ such that a ball of radius $(u + \epsilon)^{np}$ in \mathbb{R}^d can be covered by $C_2(1 + \epsilon/u)^{dnp}$ balls of radius u^{np} . It follows that for large enough n ,

$$\begin{aligned} N_{u^{np}}(\Pi([x|pn])) &\leq C_2(1 + \epsilon/u)^{dnp} N_{(u+\epsilon)^{np}}(\Pi([x|pn])) \\ &\leq C_1(p) C_2(1 + \epsilon/u)^{dnp} \prod_{i=0}^{n-1} G_p(\sigma^{pi} x). \end{aligned}$$

Hence by (5.10),

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \log N_{u^n}(\Pi([x|n])) &= \limsup_{n \rightarrow \infty} \frac{1}{np} \log N_{u^{np}}(\Pi([x|pn])) \\ &\leq d \log(1 + \epsilon/u) + \limsup_{n \rightarrow \infty} \frac{1}{np} \log \left(\prod_{i=0}^{n-1} G_p(\sigma^{pi} x) \right) \\ &\leq d \log(1 + \epsilon/u) + \epsilon + \lambda_1 + \dots + \lambda_k - k\lambda_{k+1}, \end{aligned}$$

where the first equality follows from the fact that for $pn \leq m < p(n+1)$,

$$\begin{aligned} N_{u^m}(\Pi([x|m])) &\leq N_{u^m}(\Pi([x|pn])) \leq 4^d (u^{pn-m})^d N_{u^{np}}(\Pi([x|pn])) \\ &\leq 4^d u^{-pd} N_{u^{np}}(\Pi([x|pn])), \end{aligned}$$

using the fact that for $R > r > 0$, a ball of radius R in \mathbb{R}^d can be covered by $(4R/r)^d$ balls of radius r . Letting $\epsilon \rightarrow 0$ yields the desired inequality (5.2). \square

The following result is also needed in the proof of Theorem 1.2.

Lemma 5.2. *Let m be a Borel probability measure on Σ . Let $\rho, \epsilon \in (0, 1)$. Then for m -a.e. $x = (x_n)_{n=1}^\infty \in \Sigma$,*

$$(5.11) \quad m \circ \Pi^{-1}(B(\Pi x, 2\rho^n)) \geq (1 - \epsilon)^n \frac{m([x_1 \dots x_n])}{N_{\rho^n}(\Pi([x_1 \dots x_n]))} \quad \text{for large enough } n.$$

Proof. The formulation and the proof of the above lemma are adapted from an argument given by Jordan [25]. A similar idea was also employed in the proof of [36, Theorem 2.2].

For $n \in \mathbb{N}$, let Λ_n denote the set of the points $x = (x_n)_{n=1}^\infty \in \Sigma$ such that

$$m \circ \Pi^{-1}(B(\Pi x, 2\rho^n)) < (1 - \epsilon)^n \frac{m([x_1 \dots x_n])}{N_{\rho^n}(\Pi([x_1 \dots x_n]))}.$$

To prove that (5.11) holds almost everywhere, by the Borel-Cantelli lemma it suffices to show that

$$(5.12) \quad \sum_{n=1}^{\infty} m(\Lambda_n) < \infty.$$

For this purpose, let us estimate $m(\Lambda_n)$. Fix $n \in \mathbb{N}$ and $I \in \mathcal{A}^n$. Notice that $\Pi([I])$ can be covered by $N_{\rho^n}(\Pi([I]))$ balls of radius ρ^n . As a consequence, there exists $L \leq N_{\rho^n}(\Pi([I]))$ such that $\Pi(\Lambda_n \cap [I])$ can be covered by L balls of radius ρ^n , say, B_1, \dots, B_L . We may assume that $\Pi(\Lambda_n \cap [I]) \cap B_i \neq \emptyset$ for each $1 \leq i \leq L$. Hence for each i , we may pick $x^{(i)} \in \Lambda_n \cap [I]$ such that $\Pi x^{(i)} \in B_i$. Clearly $B_i \subset B(\Pi x^{(i)}, 2\rho^n)$. Since $x^{(i)} \in \Lambda_n \cap [I]$, by the definition of Λ_n we obtain

$$m \circ \Pi^{-1}(B(\Pi x^{(i)}, 2\rho^n)) < (1 - \epsilon)^n \frac{m([I])}{N_{\rho^n}(\Pi([I]))}.$$

It follows that

$$\begin{aligned} m(\Lambda_n \cap [I]) &\leq m \circ \Pi^{-1}(\Pi(\Lambda_n \cap [I])) \\ &\leq m \circ \Pi^{-1}\left(\bigcup_{i=1}^L B_i\right) \\ &\leq m \circ \Pi^{-1}\left(\bigcup_{i=1}^L B(\Pi x^{(i)}, 2\rho^n)\right) \\ &\leq L(1 - \epsilon)^n \frac{m([I])}{N_{\rho^n}(\Pi([I]))} \\ &\leq (1 - \epsilon)^n m([I]). \end{aligned}$$

Summing over $I \in \mathcal{A}^n$ yields that $m(\Lambda_n) \leq (1 - \epsilon)^n$, which implies (5.12). \square

Remark 5.3. Lemma 5.2 remains valid when the coding map $\Pi : \Sigma \rightarrow \mathbb{R}^d$ is replaced by any Borel measurable map from Σ to \mathbb{R}^d .

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. We may assume that $s := \dim_L m < d$; otherwise there is nothing left to prove. Set $k = [s]$. Let λ_i , $i = 1, \dots, d$, be defined as in (5.1). Then by Definition 2.5,

$$h_m(\sigma) + \lambda_1 + \dots + \lambda_k + (s - k)\lambda_{k+1} = 0.$$

Let $u = \exp(\lambda_{k+1})$ and $\epsilon \in (0, 1)$. Applying Lemma 5.2 yields that for m -a.e. $x = (x_n)_{n=1}^\infty \in \Sigma$,

$$(5.13) \quad m \circ \Pi^{-1}(B(\Pi x, 2u^n)) \geq (1 - \epsilon)^n \frac{m([x_1 \dots x_n])}{N_{u^n}(\Pi([x_1 \dots x_n]))} \quad \text{for large enough } n.$$

It follows that for m -a.e. $x = (x_n)_{n=1}^\infty \in \Sigma$,

$$\begin{aligned} & \bar{d}(m \circ \Pi^{-1}, \Pi x) \\ &= \limsup_{n \rightarrow \infty} \frac{\log(m \circ \Pi^{-1}(B(\Pi x, 2u^n)))}{n \log u} \\ &\leq \frac{\log(1 - \epsilon)}{\log u} + \limsup_{n \rightarrow \infty} \left(\frac{\log m([x_1 \dots x_n])}{n \log u} - \frac{\log N_{u^n}(\Pi([x_1 \dots x_n]))}{n \log u} \right) \\ &\leq \frac{\log(1 - \epsilon) - h_m(\sigma) - (\lambda_1 + \dots + \lambda_k) + k\lambda_{k+1}}{\log u} \\ &= \frac{\log(1 - \epsilon) + s\lambda_{k+1}}{\lambda_{k+1}}, \end{aligned}$$

where in the third inequality, we used the Shannon-McMillan-Breiman theorem (cf. [39, Page. 93]) and Lemma 5.1 (keeping in mind that $\log u = \lambda_{k+1} < 0$). Letting $\epsilon \rightarrow 0$ yields the desired result. \square

6. UPPER BOUND FOR THE BOX-COUNTING DIMENSION OF C^1 -REPELLERS AND THE LYAPUNOV DIMENSIONS OF ERGODIC INVARIANT MEASURES

Throughout this section let \mathbf{M} be a smooth Riemannian manifold of dimension d and $\psi : \mathbf{M} \rightarrow \mathbf{M}$ a C^1 -map. Let Λ be a compact subset of \mathbf{M} such that $\psi(\Lambda) = \Lambda$, and assume that Λ is a repeller of ψ (cf. Section 1). Below we first introduce the definitions of singularity and Lyapunov dimensions for the case of repellers, which are somehow slightly different from that for the case of IFs.

Definition 6.1. The *singularity dimension* of Λ with respect to ψ , written as $\dim_{S^*} \Lambda$, is the unique real value s for which

$$P(\Lambda, \psi, \mathcal{G}^s) = 0,$$

where $\mathcal{G}^s = \{g_n^s\}_{n=1}^\infty$ is the sub-additive potential on Λ defined by

$$(6.1) \quad g_n^s(z) = \log \phi^s((D_z \psi^n)^{-1}), \quad z \in \Lambda.$$

Definition 6.2. For an ergodic ψ -invariant measure μ supported on Λ , the *Lyapunov dimension* of μ with respect to ψ , written as $\dim_{L^*} \mu$, is the unique real value s for which

$$h_\mu(\psi) + \mathcal{G}_*^s(\mu) = 0,$$

where $\mathcal{G}^s = \{g_n^s\}_{n=1}^\infty$ is defined as in (6.1) and $\mathcal{G}_*^s(\mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \int g_n^s d\mu$.

Before proving Theorems 1.3-1.4, we recall some definitions and necessary facts about C^1 repellers.

A finite closed cover $\{R_1, \dots, R_\ell\}$ of Λ is called a *Markov partition* of Λ with respect to ψ if:

- (i) $\overline{\text{int}R_i} = R_i$ for each $i = 1, \dots, \ell$;
- (ii) $\text{int}R_i \cap \text{int}R_j = \emptyset$ for $i \neq j$; and
- (iii) each $\psi(R_i)$ is the union of a subfamily of $\{R_j\}_{j=1}^\ell$.

It is well-known that any repeller of an expanding map has Markov partitions of arbitrary small diameter (see [37], p.146). Let $\{R_1, \dots, R_\ell\}$ be a Markov partition of Λ with respect to ψ . It is known that this dynamical system induces a subshift space of finite type (Σ_A, σ) over the alphabet $\{1, \dots, \ell\}$, where $A = (a_{ij})$ is the transfer matrix of the Markov partition, namely, $a_{ij} = 1$ if $\text{int}R_i \cap \psi^{-1}(\text{int}R_j) \neq \emptyset$ and $a_{ij} = 0$ otherwise [37]. This gives the coding map $\Pi : \Sigma_A \rightarrow \Lambda$ such that

$$(6.2) \quad \Pi(\mathbf{i}) = \bigcap_{n \geq 1} \psi^{-(n-1)}(R_{i_n}), \quad \forall \mathbf{i} = (i_1 i_2 \dots) \in \Sigma_A,$$

and the following diagram

$$(6.3) \quad \begin{array}{ccc} \Sigma_A & \xrightarrow{\sigma} & \Sigma_A \\ \Pi \downarrow & & \downarrow \Pi \\ \Lambda & \xrightarrow{\psi} & \Lambda \end{array}$$

commutes. (Keep in mind that throughout this section, Π denotes the coding map for the repeller Λ (no longer for the coding map for an IFS used in the previous sections.)

The coding map Π is a Hölder continuous surjection. Moreover there is a positive integer q so that $\Pi^{-1}(z)$ has at most q elements for each $z \in \Lambda$ (see [37], p.147).

For $n \geq 1$, define

$$\Sigma_{A,n} := \{i_1 \dots i_n \in \{1, \dots, \ell\}^n : a_{i_k i_{k+1}} = 1 \text{ for } 1 \leq k \leq n-1\}.$$

For any word $I = i_1 \dots i_n \in \Sigma_{A,n}$, the set $\bigcap_{k=1}^n \psi^{-(k-1)}(R_{i_k})$ is called a *basic set* and is denoted by R_I .

The proof of Theorem 1.3 is similar to that of Theorem 1.1. We begin with the following lemma, which is a slight variant of Lemma 4.1.

Lemma 6.3. *Let $E \subset U \subset \mathbf{M}$, where E is compact and U is open. Let $k \in \{0, 1, \dots, d-1\}$. Then for any non-degenerate C^1 map $f : U \rightarrow \mathbf{M}$, there exists $r_0 > 0$ so that for any $y \in E$, $z \in B(y, r_0)$ and $0 < r < r_0$, the set $f(B(z, r))$ can be covered by*

$$C_{\mathbf{M}} \cdot \frac{\phi^k(D_y f)}{(\alpha_{k+1}(D_y f))^k}$$

many balls of radius $\alpha_{k+1}(D_y f)r$, where $C_{\mathbf{M}}$ is a positive constant depending on \mathbf{M} .

Proof. This can be done by routinely modifying the proof of Lemma 4.1 and using similar arguments of the proof of [40, Corollary]. \square

Let $\delta > 0$ be small enough so that $\psi : B(z, \delta) \rightarrow f(B(z, \delta))$ is a diffeomorphism for each z in the δ -neighborhood of Λ . Suppose that $\{R_1, \dots, R_\ell\}$ is a Markov partition of Λ with diameter less than δ .

The following result is an analogue of Proposition 4.2.

Proposition 6.4. *Let $k \in \{0, 1, \dots, d-1\}$. Set $C_{\mathbf{M}}$ be the constant in Lemma 6.3. Then there exists $C_1 > 0$ such that for all $\mathbf{i} = (i_p)_{p=1}^\infty \in \Sigma_A$ and $n \in \mathbb{N}$, the basic set $R_{\mathbf{i}|n}$ can be covered by $C_1 \prod_{p=0}^{n-1} G(\sigma^p \mathbf{i})$ balls of radius $\prod_{p=0}^{n-1} H(\sigma^p \mathbf{i})$, where*

$$(6.4) \quad G(\mathbf{i}) := \frac{C_{\mathbf{M}} \phi^k((D_{\Pi \mathbf{i}} \psi)^{-1})}{\alpha_{k+1}((D_{\Pi \mathbf{i}} \psi)^{-1})^k}, \quad H(\mathbf{i}) := \alpha_{k+1}((D_{\Pi \mathbf{i}} \psi)^{-1}).$$

Remark 6.5. The definitions of the functions G and H in the above proposition are slightly different from that in Proposition 4.2.

Proof of Proposition 6.4. The proof is adapted from that of Proposition 4.2. For the reader's convenience, we provide the full details.

First we construct a local inverse $f_{i,j}$ of ψ for each pair (i, j) with $ij \in \Sigma_{A,2}$. To do so, notice that $\psi(R_{ij}) = R_j$ for each $ij \in \Sigma_{A,2}$. Since ψ is a diffeomorphism restricted on a small neighborhood of R_i , we can find open sets \widetilde{R}_{ij} and \widetilde{R}_j such that $\widetilde{R}_{ij} \supset R_{ij}$, $\widetilde{R}_j \supset R_j$, $\psi(\widetilde{R}_{ij}) = \widetilde{R}_j$ and $\psi : \widetilde{R}_{ij} \rightarrow \widetilde{R}_j$ is diffeomorphic. Then we take $f_{i,j} : \widetilde{R}_j \rightarrow \widetilde{R}_{ij}$ to be the inverse of $\psi : \widetilde{R}_{ij} \rightarrow \widetilde{R}_j$, and the construction is done.

For any $\mathbf{i} = (i_n)_{n=1}^\infty \in \Sigma_A$, we see that $\Pi \sigma \mathbf{i} \in R_{i_2} \subset \widetilde{R}_{i_2}$ and $(\psi \circ f_{i_1, i_2})|_{\widetilde{R}_{i_2}}$ is the identity restricted on \widetilde{R}_{i_2} . Since $\psi(\Pi \mathbf{i}) = \Pi \sigma \mathbf{i}$, it follows that $f_{i_1, i_2}(\Pi \sigma \mathbf{i}) = \Pi \mathbf{i}$. Differentiating $\psi \circ f_{i_1, i_2}$ at $\Pi \sigma \mathbf{i}$ and applying the chain rule, we get

$$(D_{\Pi \mathbf{i}} \psi)(D_{\Pi \sigma \mathbf{i}} f_{i_1, i_2}) = \text{Identity},$$

so

$$(6.5) \quad D_{\Pi \sigma \mathbf{i}} f_{i_1, i_2} = (D_{\Pi \mathbf{i}} \psi)^{-1}.$$

According to Lemma 6.3, there exists $r_0 > 0$ such that for each $ij \in \Sigma_{A,2}$, $y \in R_j$, $z \in B(y, r_0)$ and $0 < r < r_0$, the set $f_{i,j}(B(z, r))$ can be covered by

$$C_{\mathbf{M}} \cdot \frac{\phi^k(D_y f_{i,j})}{(\alpha_{k+1}(D_y f_{i,j}))^k}$$

many balls of radius $\alpha_{k+1}(D_y f_{i,j})r$.

Since ψ is expanding on Λ , there exists $\gamma \in (0, 1)$ such that $\sup_{\mathbf{i} \in \Sigma_A} \alpha_1((D_{\Pi \mathbf{i}} \psi)^{-1}) < \gamma$. Then $\sup_{\mathbf{i} \in \Sigma_A} H(\mathbf{i}) < \gamma$ and

$$(6.6) \quad \text{diam}(R_{\mathbf{i}|(n+1)}) \leq \gamma \text{diam}(R_{(\sigma \mathbf{i})|n})$$

for all $\mathbf{i} \in \Sigma_A$ and $n \in \mathbb{N}$. Take a large integer n_0 so that

$$(6.7) \quad \gamma^{n_0-1} \max\{1, \text{diam}(\Lambda)\} < r_0/2.$$

By (6.6)-(6.7), $\text{diam}(R_{\mathbf{i}|n}) < r_0/2$ for all $\mathbf{i} \in \Sigma_A$ and $n \geq n_0$.

Clearly there exists a large number C_1 so that the conclusion of the proposition holds for any positive integer $n \leq n_0$ and $\mathbf{i} \in \Sigma_A$, i.e. the set $R_{\mathbf{i}|n}$ can be covered by $C_1 \prod_{p=0}^{n-1} G(\sigma^p \mathbf{i})$ balls of radius $\prod_{p=0}^{n-1} H(\sigma^p \mathbf{i})$. Below we show by induction that this holds for all $n \in \mathbb{N}$ and $\mathbf{i} \in \Sigma_A$.

Suppose for some $m \geq n_0$, the conclusion of the proposition holds for any positive integer $n \leq m$ and $\mathbf{i} \in \Sigma_A$. Then for given $\mathbf{i} = (i_n)_{n=1}^\infty \in \Sigma_A$, $R_{(\sigma \mathbf{i})|m}$ can be covered by $C_1 \prod_{p=0}^{m-1} G(\sigma^{p+1} \mathbf{i})$ balls of radius $\prod_{p=0}^{m-1} H(\sigma^{p+1} \mathbf{i})$. Let B_1, \dots, B_N denote these balls. We may assume that $B_j \cap R_{(\sigma \mathbf{i})|m} \neq \emptyset$ for each j . Since

$$\prod_{p=0}^{m-1} H(\sigma^{p+1} \mathbf{i}) \leq \gamma^m \leq \gamma^{n_0} < r_0/2$$

and

$$d(\Pi \sigma \mathbf{i}, B_j \cap R_{(\sigma \mathbf{i})|m}) \leq \text{diam}(R_{(\sigma \mathbf{i})|m}) < r_0/2,$$

so the center of B_j is in $B(\Pi \sigma \mathbf{i}, r_0)$. Therefore by Lemma 6.3 and (6.5), $f_{i_1, i_2}(B_j)$ can be covered by

$$C_M \cdot \frac{\phi^k(D_{\Pi \sigma \mathbf{i}} f_{i_1, i_2})}{(\alpha_{k+1}(D_{\Pi \sigma \mathbf{i}} f_{i_1, i_2}))^k} = G(\mathbf{i})$$

balls of radius

$$\alpha_{k+1}(D_{\Pi \sigma \mathbf{i}} f_{i_1, i_2}) \cdot \prod_{p=0}^{m-1} H(\sigma^{p+1} \mathbf{i}) = H(\mathbf{i}) \cdot \prod_{p=0}^{m-1} H(\sigma^{p+1} \mathbf{i}) = \prod_{p=0}^m H(\sigma^p \mathbf{i}).$$

Since $\psi(R_{\mathbf{i}|(m+1)}) \subset R_{(\sigma \mathbf{i})|m}$, it follows that

$$R_{\mathbf{i}|(m+1)} \subset f_{i_1, i_2}(R_{(\sigma \mathbf{i})|m}) \subset \bigcup_{j=1}^N f_{i_1, i_2}(B_j),$$

hence $R_{\mathbf{i}|(m+1)}$ can be covered by

$$G(\mathbf{i})N \leq G(\mathbf{i}) \cdot C_1 \prod_{p=0}^{m-1} G(\sigma^{p+1} \mathbf{i}) = C_1 \prod_{p=0}^m G(\sigma^p \mathbf{i})$$

balls of radius $\prod_{p=0}^m H(\sigma^p \mathbf{i})$. Thus the proposition also holds for $n = m + 1$ and all $\mathbf{i} \in \Sigma_A$, as desired. \square

Proposition 6.6. *Let $k \in \{0, 1, \dots, d-1\}$. Let $G, H : \Sigma_A \rightarrow \mathbb{R}$ be defined as in (6.4). Let t be the unique real number so that*

$$P(\Sigma_A, \sigma, (\log G) + t(\log H)) = 0.$$

Then $\overline{\dim}_B \Lambda \leq t$.

Proof. Here we use similar arguments as that in the proof of Proposition 4.3. Write $g = \log G$ and $h = \log H$. Define

$$r_{\min} = \min_{\mathbf{i} \in \Sigma_A} h(\mathbf{i}), \quad r_{\max} = \max_{\mathbf{i} \in \Sigma_A} h(\mathbf{i}).$$

Then $0 < r_{\min} \leq r_{\max} < 1$. For $0 < r < r_{\min}$, define

$$(6.8) \quad \mathcal{A}_r = \left\{ i_1 \dots i_n \in \Sigma_A^* : \sup_{x \in [i_1 \dots i_n] \cap \Sigma_A} S_n h(x) < \log r \leq \sup_{y \in [i_1 \dots i_{n-1}] \cap \Sigma_A} S_{n-1} h(y) \right\},$$

where Σ_A^* denotes the set of all finite words allowed in Σ_A . Clearly $\{[I] : I \in \mathcal{A}_r\}$ is a partition of Σ_A . By Proposition 6.4, there exists a constant $C_1 > 0$ such that for each $0 < r < r_{\min}$, every $I \in \mathcal{A}_r$ and $x \in [I]$, R_I can be covered by

$$C_1 \exp(S_{|I|} g(x)) \leq C_1 \exp \left(\sup_{y \in [I]} S_{|I|} g(y) \right)$$

many balls of radius

$$\exp(S_{|I|} h(x)) \leq \exp \left(\sup_{y \in [I]} S_{|I|} h(y) \right) < r.$$

It follows that Λ can be covered by

$$C_1 \sum_{I \in \mathcal{A}_r} \exp \left(\sup_{y \in [I]} S_{|I|} g(y) \right)$$

many balls of radius r . Hence by Proposition 3.4,

$$\overline{\dim}_B \Lambda \leq \limsup_{r \rightarrow 0} \frac{\sum_{I \in \mathcal{A}_r} \exp \left(\sup_{y \in [I]} S_{|I|} g(y) \right)}{\log(1/r)} = t.$$

This completes the proof of the proposition. \square

For $n \in \mathbb{N}$, applying Proposition 6.6 to the mapping ψ^n instead of ψ , we obtain the following.

Proposition 6.7. *Let $k \in \{0, \dots, d-1\}$. Then for each $n \in \mathbb{N}$,*

$$\overline{\dim}_B \Lambda \leq t_n,$$

where t_n is the unique number for which $P(\Sigma_A, \sigma^n, (\log G_n) + t_n(\log H_n)) = 0$, and G_n, H_n are continuous functions on Σ_A defined by

$$(6.9) \quad G_n(y) := \frac{C \phi^k((D_{\Pi y} \psi^n)^{-1})}{\alpha_{k+1}((D_{\Pi y} \psi^n)^{-1})^k}, \quad H_n(y) := \alpha_{k+1}((D_{\Pi y} \psi^n)^{-1}),$$

with $C = C_{\mathbf{M}}$ being the constant in Lemma 6.3.

Now we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. We follow the proof of Theorem 1.1 with slight modifications. Write $s = \dim_{S^*}(\Lambda)$. We may assume that $s < d$; otherwise we have nothing left to prove. Set $k = [s]$. Let $\mathcal{G}^s = \{g_n^s\}_{n=1}^\infty$ be the sub-additive potential on Λ defined by

$$g_n^s(z) = \log \phi^s((D_z \psi^n)^{-1}).$$

Then $P(\Lambda, \psi, \mathcal{G}^s) = 0$ by the definition of $\dim_{S^*}(\Lambda)$. Let $\widehat{\mathcal{G}}^s := \{\widehat{g}_n^s\}_{n=1}^\infty$, where $\widehat{g}_n^s \in C(\Sigma_A)$ is defined by

$$\widehat{g}_n^s(\mathbf{i}) = g_n^s(\Pi \mathbf{i}) = \log \phi^s((D_{\Pi \mathbf{i}} \psi^n)^{-1}), \quad \mathbf{i} \in \Sigma_A.$$

Clearly, $\widehat{\mathcal{G}}^s$ is a sub-additive potential on Σ_A and

$$(\widehat{\mathcal{G}}^s)_*(m) = (\widehat{\mathcal{G}}^s)_*(m \circ \Pi^{-1}), \quad m \in \mathcal{M}(\Sigma_A, \sigma).$$

Since the factor map $\Pi : \Sigma_A \rightarrow \Lambda$ is onto and finite-to-one, by Lemma 2.3, $m \rightarrow m \circ \Pi^{-1}$ is a surjective map from $\mathcal{M}(\Sigma_A, \sigma)$ to $\mathcal{M}(\Lambda, \psi)$ and moreover, $h_m(\sigma) = h_{m \circ \Pi^{-1}}(\psi)$ for $m \in \mathcal{M}(\Sigma_A, \sigma)$. By the variational principle for the sub-additive pressure (see Theorem 2.2),

$$\begin{aligned} P(\Lambda, \psi, \mathcal{G}^s) &= \sup \{h_\mu(\psi) + (\mathcal{G}^s)_*(\mu) : \mu \in \mathcal{M}(\Lambda, \psi)\} \\ &= \sup \{h_{m \circ \Pi^{-1}}(\psi) + (\mathcal{G}^s)_*(m \circ \Pi^{-1}) : m \in \mathcal{M}(\Sigma_A, \sigma)\} \\ &= \sup \{h_m(\sigma) + (\widehat{\mathcal{G}}^s)_*(m) : m \in \mathcal{M}(\Sigma_A, \sigma)\} \\ &= P(\Sigma_A, \sigma, \widehat{\mathcal{G}}^s). \end{aligned}$$

It follows that $P(\Sigma_A, \sigma, \widehat{\mathcal{G}}^s) = 0$.

By Proposition 2.2 in [2],

$$\lim_{n \rightarrow \infty} \frac{1}{n} P(\Sigma_A, \sigma^n, \widehat{g}_n^s) = \inf_{n \geq 1} \frac{1}{n} P(\Sigma_A, \sigma^n, \widehat{g}_n^s) = P(\Sigma_A, \sigma, \widehat{\mathcal{G}}^s).$$

Since $P(\Sigma_A, \sigma, \widehat{\mathcal{G}}^s) = 0$, the above equalities imply that for each $\epsilon > 0$, there exists $N_\epsilon > 0$ such that

$$(6.10) \quad 0 \leq P(\Sigma_A, \sigma^n, \widehat{g}_n^s) \leq n\epsilon \quad \text{for } n > N_\epsilon.$$

For $n \in \mathbb{N}$, let s_n be the unique real number so that $P(\Sigma_A, \sigma^n, v_n^{s_n}) = 0$, where $v_n^{s_n}$ is a continuous function on Σ_A defined by

$$\begin{aligned} v_n^{s_n}(x) &= \log G_n(x) + s_n \log H_n(x) \\ &= \log (C \phi^k((D_{\Pi x} \psi^n)^{-1}) \alpha_{k+1}((D_{\Pi x} \psi^n)^{-1})^{s_n - k}) \end{aligned}$$

where G_n, H_n are defined in (6.9) and $C = C_{\mathbf{M}}$. By Proposition 6.7,

$$\overline{\dim}_B \Lambda \leq s_n.$$

If $s \geq s_n$ for some n , then $\overline{\dim}_B \Lambda \leq s_n \leq s$ and we are done. In what follows, we assume that $s < s_n$ for each n . Then for each $x \in \Sigma_A$,

$$\begin{aligned} \widehat{g}_n^s(x) - v_n^{s_n}(x) &= -\log C + (s - s_n) \log \alpha_{k+1}((D_{\Pi x} \psi^n)^{-1}) \\ &\geq -\log C + (s - s_n) \log \alpha_1((D_{\Pi x} \psi^n)^{-1}) \\ &\geq -\log C + n(s - s_n) \log \theta, \end{aligned}$$

where

$$\theta := \max_{x \in \Sigma_A} \alpha_1((D_{\Pi x} \phi)^{-1}) < 1.$$

Hence

$$\begin{aligned} P(\Sigma_A, \sigma^n, \widehat{g}_n^s) &= P(\Sigma_A, \sigma^n, \widehat{g}_n^s) - P(\Sigma_A, \sigma^n, v_n^{s_n}) \\ &\geq \inf_{x \in \Sigma_A} (\widehat{g}_n^s(x) - v_n^{s_n}(x)) \\ &\geq -\log C + n(s - s_n) \log \theta. \end{aligned}$$

where in the second inequality, we used [39, Theorem 9.7(iv)]. Combining it with (6.10) yields that for $n \geq N_\epsilon$,

$$n\epsilon \geq -\log C + n(s - s_n) \log \theta,$$

so

$$s \geq s_n + \frac{\epsilon + n^{-1} \log C}{\log \theta} \geq \overline{\dim}_B \Lambda + \frac{\epsilon + n^{-1} \log C}{\log \theta}.$$

Letting $n \rightarrow \infty$ and then $\epsilon \rightarrow 0$, we obtain $s \geq \overline{\dim}_B \Lambda$, as desired. \square

In the remaining part of this section, we prove Theorem 1.4. To this end, we need the following two lemmas, which are the analogues of Lemmas 5.1-5.2 for C^1 repellers.

Lemma 6.8. *Let m be an ergodic σ -invariant Borel probability measure on Σ_A . Set*

$$(6.11) \quad \lambda_i := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log(\alpha_i((D_{\Pi x} \psi^n)^{-1})) \, dm(x), \quad i = 1, \dots, d.$$

Let $k \in \{0, \dots, d-1\}$. Write $u := \exp(\lambda_{k+1})$. Then for m -a.e. $x \in \Sigma_A$,

$$(6.12) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \log N_{u^n}(R_{x|n}) \leq (\lambda_1 + \dots + \lambda_k) - k\lambda_{k+1},$$

where $N_\delta(E)$ is the smallest integer N for which E can be covered by N closed balls of radius δ .

Lemma 6.9. *Let m be a Borel probability measure on Σ_A . Let $\rho, \epsilon \in (0, 1)$. Then for m -a.e. $x = (x_n)_{n=1}^\infty \in \Sigma_A$,*

$$(6.13) \quad m \circ \Pi^{-1}(B(\Pi x, 2\rho^n)) \geq (1 - \epsilon)^n \frac{m([x_1 \dots x_n])}{N_{\rho^n}(R_{x_1 \dots x_n})} \quad \text{for large enough } n.$$

The proofs of these two lemmas are essentially identical to that of Lemmas 5.1-5.2. So we simply omit them.

Proof of Theorem 1.4. Here we adapt the proof of Theorem 1.2. We may assume that $s := \dim_{L^*} \mu < d$; otherwise there is nothing left to prove. Since $\Pi : \Sigma_A \rightarrow \Lambda$ is surjective and finite-to-one, by Lemma 2.3, there exists a σ -invariant ergodic measure m on Σ_A so that $m \circ \Pi^{-1} = \mu$ and $h_m(\sigma) = h_\mu(\psi)$.

Set $k = [s]$. Let λ_i , $i = 1, \dots, d$, be defined as in (6.11). Then by Definition 6.2,

$$h_m(\sigma) + \lambda_1 + \dots + \lambda_k + (s - k)\lambda_{k+1} = 0.$$

Let $u = \exp(\lambda_{k+1})$ and $\epsilon \in (0, 1)$. Applying Lemma 6.9 (in which we take $\rho = u$) yields that for m -a.e. $x = (x_n)_{n=1}^\infty \in \Sigma_A$,

$$(6.14) \quad \mu(B(\Pi x, 2u^n)) \geq (1 - \epsilon)^n \frac{m([x_1 \dots x_n])}{N_{u^n}(R_{x_1 \dots x_n})} \quad \text{for large enough } n.$$

It follows that for m -a.e. $x = (x_n)_{n=1}^\infty \in \Sigma_A$,

$$\begin{aligned} \bar{d}(\mu, \Pi x) &= \limsup_{n \rightarrow \infty} \frac{\log(\mu(B(\Pi x, 2u^n)))}{n \log u} \\ &\leq \frac{\log(1 - \epsilon)}{\log u} + \limsup_{n \rightarrow \infty} \left(\frac{\log m([x_1 \dots x_n])}{n \log u} - \frac{\log N_{u^n}(\Pi([x_1 \dots x_n]))}{n \log u} \right) \\ &\leq \frac{\log(1 - \epsilon) - h_m(\sigma) - (\lambda_1 + \dots + \lambda_k) + k\lambda_{k+1}}{\log u} \\ &= \frac{\log(1 - \epsilon) + s\lambda_{k+1}}{\lambda_{k+1}}, \end{aligned}$$

where in the third inequality, we used the Shannon-McMillan-Breiman theorem (cf. [39, Page. 93]) and Lemma 6.8 (keeping in mind that $\log u = \lambda_{k+1} < 0$). Letting $\epsilon \rightarrow 0$ yields the desired result. \square

Remark 6.10. There is an alternative way to prove Theorem 1.3 in the case when \mathbf{M} is an open set of \mathbb{R}^d . In such case, we can construct a C^1 IFS $\{f_i\}_{i=1}^\ell$ on \mathbb{R}^d so that Λ is the projection of a shift invariant set under the coding map associated with the IFS. Then we can use Theorem 1.1 to get an upper bound for $\overline{\dim}_B \Lambda$. An additional effort is then required to justify that this upper bound is indeed equal to $\dim_{S^*} \Lambda$. The details of this approach will be given in a forthcoming survey paper.

Acknowledgement. The research of Feng was partially supported by a HKRGC GRF grant and the Direct Grant for Research in CUHK. The research of Simon was partially supported by the grant OTKA K104745.

APPENDIX A.

In this part, we prove the following.

Lemma A.1. *Let $(\mathcal{A}^\mathbb{N}, \sigma)$ be the one-sided full shift space over a finite alphabet \mathcal{A} . Let $\nu \in \mathcal{M}(\mathcal{A}^\mathbb{N}, \sigma^m)$ for some $m \in \mathbb{N}$. Set $\mu = \frac{1}{m} \sum_{k=0}^{m-1} \nu \circ \sigma^{-k}$. Then $\mu \in \mathcal{M}(\mathcal{A}^\mathbb{N}, \sigma)$ and $h_\mu(\sigma) = \frac{1}{m} h_\nu(\sigma^m)$.*

Proof. The lemma might be well-known. However we are not able to find a reference, so for the convenience of the reader we provide a self-contained proof. The σ -invariance of μ follows directly from its definition, and we only need to prove that $h_\mu(\sigma) = \frac{1}{m} h_\nu(\sigma^m)$.

Clearly, $\nu \circ \sigma^{-k} \in \mathcal{M}(\mathcal{A}^\mathbb{N}, \sigma^m)$ for $k = 0, \dots, m-1$. We claim that $h_{\nu \circ \sigma^{-k}}(\sigma^m) = h_\nu(\sigma^m)$ for $k = 1, \dots, m-1$. Without loss of generality we prove this in the case when $k = 1$. For $n \in \mathbb{N}$, let \mathcal{P}_n denote the partition of $\mathcal{A}^\mathbb{N}$ consisting of the n -th

cylinders of $\mathcal{A}^{\mathbb{N}}$, i.e. $\mathcal{P}_n = \{[I] : I \in \mathcal{A}^n\}$, and set $\sigma^{-1}\mathcal{P}_n = \{\sigma^{-1}([I]) : I \in \mathcal{A}^n\}$. Then it is direct to see that any element in \mathcal{P}_n intersects at most $\#\mathcal{A}$ elements in $\sigma^{-1}\mathcal{P}_n$, and vice versa. Hence

$$|H_\nu(\mathcal{P}_n) - H_\nu(\sigma^{-1}\mathcal{P}_n)| \leq \log \#\mathcal{A};$$

see e.g. [20, Lemma 4.6]. It follows that

$$h_{\nu \circ \sigma^{-1}}(\sigma^m) = \lim_{n \rightarrow \infty} \frac{1}{n} H_{\nu \circ \sigma^{-1}}(\mathcal{P}_{nm}) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\nu(\sigma^{-1}\mathcal{P}_{nm}) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\nu(\mathcal{P}_{nm}) = h_\nu(\sigma^m).$$

This prove the claim.

By the affinity of the measure-theoretic entropy $h_{(\cdot)}(\sigma^m)$ (see [39, Theorem 8.1]), we have

$$h_\mu(\sigma^m) = \frac{1}{m} \sum_{k=0}^{m-1} h_{\nu \circ \sigma^{-k}}(\sigma^m) = h_\nu(\sigma^m),$$

where the second equality follows from the above claim. Hence we have $h_\mu(\sigma) = \frac{1}{m} h_\mu(\sigma^m) = \frac{1}{m} h_\nu(\sigma^m)$. \square

APPENDIX B. MAIN NOTATION AND CONVENTIONS

For the reader's convenience, we summarize in Table 1 the main notation and typographical conventions used in this paper.

TABLE 1. Main notation and conventions

| | |
|------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| $\overline{\dim}_B$ | Upper box-counting dimension |
| $\dim_{\mathcal{P}}$ | Packing dimension |
| $\{f_i\}_{i=1}^\ell$ | A C^1 IFS (Section 1) |
| (Σ, σ) | One-sided full shift over the alphabet $\{1, \dots, \ell\}$ |
| $\Pi : \Sigma \rightarrow K$ | Coding map associated with $\{f_i\}_{i=1}^\ell$ (Section 1) |
| $D_x f$ | Differential of f at x |
| $\dim_S X$ | Singular dimension of X w.r.t. $\{f_i\}_{i=1}^\ell$ (cf. Definition 2.4) |
| $\dim_L m$ | Lyapunov dimension of m w.r.t. $\{f_i\}_{i=1}^\ell$ (cf. Definition 2.5) |
| $P(X, T, \{g_n\}_{n=1}^\infty)$ | Topological pressure of a sub-additive potential $\{g_n\}_{n=1}^\infty$ on a topological dynamical system (X, T) (cf. Definition 2.1) |
| $h_\mu(T)$ | Measure-theoretic entropy of μ w.r.t. T |
| $\mathcal{G}_*(\mu)$ | Lyapunov exponent of a sub-additive potential \mathcal{G} w.r.t. μ (cf. (2.5)) |
| $\alpha_i(T), i = 1, \dots, d$ | The i -th singular value of $T \in \mathbb{R}^{d \times d}$ (Section 2) |
| ϕ^s | Singular value function (cf. 2.9) |
| $S_n g$ | $g + g \circ T + \dots + g \circ T^{n-1}$ for $g \in C(X)$ |
| $\mathcal{G}^s = \{g_n^s\}_{n=1}^\infty$ | (cf. (2.10)) |
| $\mathcal{G}_*^s(m)$ | $\lim_{n \rightarrow \infty} \frac{1}{n} \int g_n^s dm$ |
| $\lambda_i(m), i = 1, \dots, d$ | The i -th Lyapunov exponent of m w.r.t. $\{f_i\}_{i=1}^\ell$ (cf. (2.11)) |
| $N_\delta(E)$ | Smallest number of closed balls of radius δ required to cover E |
| $\dim_{S^*} \Lambda$ | Singularity dimension of Λ w.r.t. ψ (cf. Definition 6.1) |
| $\dim_{L^*} \mu$ | Lyapunov dimension of μ w.r.t. ψ (cf. Definition 6.2) |

REFERENCES

- [1] Ludwig Arnold. *Random dynamical systems*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.
- [2] Jungchao Ban, Yongluo Cao, and Huyi Hu. The dimensions of a non-conformal repeller and an average conformal repeller. *Trans. Amer. Math. Soc.*, 362(2):727–751, 2010.
- [3] Balázs Bárány, Michael Hochman, and Ariel Rapaport. Hausdorff dimension of planar self-affine sets and measures. *Invent. Math.*, 216(3):601–659, 2019.
- [4] Luis Barreira. Dimension estimates in non-conformal hyperbolic dynamics. *Nonlinearity*, 16(5):1657–1672, 2003.
- [5] Luis Barreira. Almost additive thermodynamic formalism: some recent developments. *Rev. Math. Phys.*, 22(10):1147–1179, 2010.
- [6] Thomas Bogenschütz and Hans Crauel. The Abramov-Rokhlin formula. In *Ergodic theory and related topics, III (Güstrow, 1990)*, volume 1514 of *Lecture Notes in Math.*, pages 32–35. Springer, Berlin, 1992.
- [7] Rufus Bowen. Hausdorff dimension of quasicircles. *Inst. Hautes Études Sci. Publ. Math.*, (50):11–25, 1979.
- [8] Rufus Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, volume 470 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, revised edition, 2008. With a preface by David Ruelle, Edited by Jean-René Chazottes.
- [9] Yong-Luo Cao, De-Jun Feng, and Wen Huang. The thermodynamic formalism for sub-additive potentials. *Discrete Contin. Dyn. Syst.*, 20(3):639–657, 2008.
- [10] Yongluo Cao, Yakov Pesin, and Yun Zhao. Dimension estimates for non-conformal repellers and continuity of sub-additive topological pressure. *Geom. Funct. Anal.*, 29(5):1325–1368, 2019.
- [11] Jianyu Chen and Yakov Pesin. Dimension of non-conformal repellers: a survey. *Nonlinearity*, 23(4):R93–R114, 2010.
- [12] Adrien Douady and Joseph Oesterlé. Dimension de Hausdorff des attracteurs. *C. R. Acad. Sci. Paris Sér. A-B*, 290(24):A1135–A1138, 1980.
- [13] Kenneth J. Falconer. The Hausdorff dimension of self-affine fractals. *Math. Proc. Cambridge Philos. Soc.*, 103(2):339–350, 1988.
- [14] Kenneth J. Falconer. A subadditive thermodynamic formalism for mixing repellers. *J. Phys. A*, 21(14):L737–L742, 1988.
- [15] Kenneth J. Falconer. Bounded distortion and dimension for nonconformal repellers. *Math. Proc. Cambridge Philos. Soc.*, 115(2):315–334, 1994.
- [16] Kenneth J. Falconer. *Fractal geometry, Mathematical foundations and applications.* John Wiley & Sons, Inc., Hoboken, NJ, second edition, 2003.
- [17] Ai-Hua Fan, Ka-Sing Lau, and Hui Rao. Relationships between different dimensions of a measure. *Monatsh. Math.*, 135(3):191–201, 2002.
- [18] De-Jun Feng. The variational principle for products of non-negative matrices. *Nonlinearity*, 17(2):447–457, 2004.
- [19] De-Jun Feng. Equilibrium states for factor maps between subshifts. *Adv. Math.*, 226(3):2470–2502, 2011.
- [20] De-Jun Feng and Huyi Hu. Dimension theory of iterated function systems. *Comm. Pure Appl. Math.*, 62(11):1435–1500, 2009.
- [21] De-Jun Feng and Ka-Sing Lau. The pressure function for products of non-negative matrices. *Math. Res. Lett.*, 9(2-3):363–378, 2002.
- [22] De-Jun Feng and Karoly Simon. Dimension estimates for C^1 iterated function systems and repellers. Part II. In preparation. 2020
- [23] Dimitrios Gatzouras and Yuval Peres. Invariant measures of full dimension for some expanding maps. *Ergodic Theory Dynam. Systems*, 17(1):147–167, 1997.

- [24] Michael Hochman and Ariel Rapaport. Hausdorff dimension of planar self-affine sets and measures with overlaps. *arXiv preprint arXiv:1904.09812*, 2019.
- [25] Thomas Jordan. Unpublished note. 2011.
- [26] Thomas Jordan and Mark Pollicott. The Hausdorff dimension of measures for iterated function systems which contract on average. *Discrete Contin. Dyn. Syst.*, 22(1-2):235–246, 2008.
- [27] Thomas Jordan, Mark Pollicott, and Károly Simon. Hausdorff dimension for randomly perturbed self affine attractors. *Comm. Math. Phys.*, 270(2):519–544, 2007.
- [28] Antti Käenmäki. On natural invariant measures on generalised iterated function systems. *Ann. Acad. Sci. Fenn. Math.*, 29(2):419–458, 2004.
- [29] Antti Käenmäki and Markku Vilppolainen. Dimension and measures on sub-self-affine sets. *Monatsh. Math.*, 161(3):271–293, 2010.
- [30] R. Kenyon and Y. Peres. Measures of full dimension on affine-invariant sets. *Ergodic Theory Dynam. Systems*, 16(2):307–323, 1996.
- [31] Ricardo Mañé. *Ergodic theory and differentiable dynamics*. Springer-Verlag, Berlin, 1987.
- [32] A. Manning and K. Simon. Subadditive pressure for triangular maps. *Nonlinearity*, 20(1):133–149, 2007.
- [33] Anna Mummert. The thermodynamic formalism for almost-additive sequences. *Discrete Contin. Dyn. Syst.*, 16(2):435–454, 2006.
- [34] Norbert Patzschke. Self-conformal multifractal measures. *Adv. in Appl. Math.*, 19(4):486–513, 1997.
- [35] Yakov B. Pesin. *Dimension theory in dynamical systems. Contemporary views and applications*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1997.
- [36] Eino Rossi. Local dimensions of measures on infinitely generated self-affine sets. *J. Math. Anal. Appl.*, 413(2):1030–1039, 2014.
- [37] David Ruelle. *Thermodynamic formalism*, volume 5 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Co., Reading, Mass., 1978.
- [38] David Ruelle. Repellers for real analytic maps. *Ergodic Theory Dynam. Systems*, 2(1):99–107, 1982.
- [39] Peter Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [40] Yingjie Zhang. Dynamical upper bounds for Hausdorff dimension of invariant sets. *Ergodic Theory Dynam. Systems*, 17(3):739–756, 1997.

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG,

Email address: `djfeng@math.cuhk.edu.hk`

(Károly Simon) BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, DEPARTMENT OF STOCHASTICS, INSTITUTE OF MATHEMATICS AND MTA-BME STOCHASTICS RESEARCH GROUP, 1521 BUDAPEST, P.O.BOX 91, HUNGARY

Email address: `simonk@math.bme.hu`