

Fourier Analysis

Feb. 22, 2022

Review:

A convergence Thm: Let f be integrable on the circle.

Suppose f is differentiable at x_0 . Then

$$S_N f(x_0) \rightarrow f'(x_0) \quad \text{as } N \rightarrow \infty.$$

The above result is based on the Riemann - Lebesgue lemma.

Remark: Using the same proof, we can show that the above result holds under the weaker assumption that

f is Lipschitz continuous at x_0 , this is,

$\exists M > 0$ such that

$$(*) \quad |f(x) - f(x_0)| \leq M \cdot |x - x_0| \quad \text{for all } x \in [-\pi, \pi]$$

Corollary 1. Let $f, g \in R$. Suppose that

f and g coincide on an open interval I which contains x_0 . Then

$$S_N f(x_0) - S_N g(x_0) \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

pf. Apply the convergence thm to $f - g$ at x_0 .

④.

FACT: The above result implies that

the convergence of $S_N f$ at x_0 only depends on the behavior of f on a neighborhood of x_0 .


This fact is called "Riemann's localization principle".

§ 3.2

A continuous function on the circle whose Fourier series diverges at some point

Let us start from a special function

$$f(x) = \begin{cases} i(\pi - x) & \text{if } 0 \leq x \leq \pi \\ i(-\pi - x) & \text{if } -\pi \leq x < 0 \end{cases}$$



A direct calculation shows that

$$f(x) \sim \sum_{n \neq 0} \frac{1}{n} e^{inx} \quad \text{on } [-\pi, \pi]$$

For $N \in \mathbb{N}$, define

$$f_N(x) = \sum_{0 < |n| \leq N} \frac{1}{n} e^{inx} \quad (= S_N f(x))$$

$$\tilde{f}_N(x) = \sum_{n=-N}^{-1} \frac{1}{n} e^{inx}$$

Lem 2 : (1) \exists a constant $M > 0$ such that

$$|f_N(x)| \leq M \text{ for all } N \in \mathbb{N} \text{ and } x \in \mathbb{R}$$

$$(2) |\tilde{f}_N(0)| \geq \log N$$

Pf. We first prove (1).

Let us consider the Abel mean of f ,

$$A_r f(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(x-y) f(y) dy,$$

$$\text{where } P_r(x) := \frac{1-r^2}{1-2r \cos x + r^2}, \quad 0 \leq r < 1.$$

Notice that for $0 \leq r < 1$,

$$\begin{aligned}
 |A_r f(x)| &\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(x-y) |f(y)| dy \\
 &\leq \frac{\|f\|_{\infty}}{2\pi} \int_{-\pi}^{\pi} P_r(x-y) dy \\
 &= \|f\|_{\infty} \quad (\|f\|_{\infty} := \sup_{y \in [-\pi, \pi]} |f(y)|)
 \end{aligned}$$

However

$$\begin{aligned}
 &|f_N(x) - A_r f(x)| \\
 &= \left| \sum_{0 < |n| \leq N} \frac{1}{n} e^{inx} - \sum_{|n| > 0} \frac{r^{|n|}}{n} e^{inx} \right| \\
 &= \left| \sum_{0 < |n| \leq N} \frac{1 - r^{|n|}}{n} e^{inx} - \sum_{|n| \geq N+1} \frac{r^{|n|}}{n} e^{inx} \right| \\
 &\leq \left| \sum_{0 < |n| \leq N} \frac{1 - r^{|n|}}{n} e^{inx} \right| + \left| \sum_{|n| \geq N+1} \frac{r^{|n|}}{n} e^{inx} \right|
 \end{aligned}$$

$$\leq \sum_{0 < |n| \leq N} \frac{1 - r^{|n|}}{|n|} + \sum_{|n| > N+1} \frac{r^{|n|}}{|n|}$$

$$\leq 2 \cdot \sum_{n=1}^N \frac{1 - r^n}{n} + 2 \cdot \sum_{n=N+1}^{\infty} \frac{r^n}{N}$$

Notice that $\frac{1 - r^n}{n} = \frac{(1-r)(1+r+r^2+\dots+r^{n-1})}{n}$

$$\leq 1-r.$$

Hence

$$|f_N(x) - Arf(x)| \leq 2N(1-r) + 2 \sum_{n=N+1}^{\infty} \frac{r^n}{N}$$

$$\leq 2N(1-r) + 2 \cdot \frac{r^{N+1}}{N(1-r)}$$

$$\leq 2N(1-r) + \frac{2}{N(1-r)}$$

Let us take $\tilde{r} = 1 - \frac{1}{N}$, then $N(1-\tilde{r}) = 1$

Then $|f_N(x) - A\tilde{r}f(x)| \leq 4$

It follows that

$$\begin{aligned}|f_N(x)| &\leq 4 + |A \tilde{f}(x)| \\ &\leq 4 + \|f\|_{\infty}\end{aligned}$$

This prove (1).

Next we prove (2), i.e. $|\widehat{f}_N(0)| > \log N$

Notice that

$$\widehat{f}_N(0) = \sum_{n=-N}^{-1} \frac{1}{n} e^{in \cdot 0} = - \left[1 + \frac{1}{2} + \dots + \frac{1}{N} \right]$$

So

$$|\widehat{f}_N(0)| = 1 + \frac{1}{2} + \dots + \frac{1}{N}$$

$$\begin{aligned}&\text{(using } \frac{1}{k} \geq \int_k^{k+1} \frac{1}{x} dx \text{ for all } k > 0 \text{)} \\ &\geq \sum_{k=1}^N \int_k^{k+1} \frac{1}{x} dx\end{aligned}$$

$$= \sum_{k=1}^N (\log(k+1) - \log k)$$

$$= \log(N+1) > \log N.$$

□

Next for $N \in \mathbb{N}$, define

$$P_N(x) = e^{2inx} \cdot f_N(x)$$

$$= \sum_{\substack{n=-N \\ n \neq 0}}^N \frac{1}{n} e^{i(n+2N)x}$$

$$= \sum_{\substack{n=N \\ n \neq 2N}}^{3N} \frac{1}{n-2N} e^{inx}$$

$$\tilde{P}_N(x) = e^{2inx} \cdot \tilde{f}_N(x) = \sum_{n=N}^{2N-1} \frac{1}{n-2N} e^{inx}$$

Clearly $|P_N(x)| = |f_N(x)| \leq M$,

$$|\tilde{P}_N(x)| = |\tilde{f}_N(x)|, \quad |\tilde{P}_N(0)| > \log N.$$

Construct a sequence $(N_k)_{k=1}^{\infty}$ of positive integers and a sequence $(d_k)_{k=1}^{\infty}$ of positive numbers such that

$$(1) \quad N_{k+1} > 3N_k$$

$$(2) \quad \sum_{k=1}^{\infty} d_k < \infty$$

$$(3) \quad d_k \log N_k \rightarrow \infty \text{ as } k \rightarrow \infty.$$

(For instance, we can take

$$N_k = 4^{4^k} \text{ and } d_k = 3^{-k})$$

Finally set

$$g(x) = \sum_{k=1}^{\infty} d_k P_{N_k}(x), \quad x \in [-\pi, \pi].$$

$$\text{Since } |P_{N_k}(x)| \leq M \text{ and } \sum_{k=1}^{\infty} d_k M < \infty,$$

by the Weierstrass M-Test, we have

$$\sum_{k=1}^{\ell} \alpha_k P_{N_k}(x) \xrightarrow{\quad} g \quad \text{on the circle}$$

as $\ell \rightarrow \infty$.

and g is continuous on the circle.

Next we claim that

$$S_N g(0) \not\rightarrow g(0) \quad \text{as } N \rightarrow \infty \quad (**)$$

Lem 3.

$$\hat{g}(n) = \begin{cases} \frac{\alpha_k}{n - 2N_k} & \text{if } n \in [N_k, 3N_k] \\ & \text{and } n \neq 2N_k \\ 0 & \text{otherwise.} \end{cases}$$

Pf. Recall that for given N ,

$$P_N(x) = \sum_{\substack{N \leq n \leq 3N \\ n \neq 2N}} \frac{1}{n-2N} e^{inx}$$

Hence $\hat{P}_N(n) = \begin{cases} \frac{1}{n-2N} & \text{if } n \in [N, 3N] \\ 0 & \text{otherwise} \end{cases}$

Now fix $n \in \mathbb{Z}$. Let $\varepsilon > 0$. Then $\exists L \in \mathbb{N}$ such that

$$\left| g(x) - \sum_{j=1}^l \alpha_j \widehat{P}_{N_j}(x) \right| < \varepsilon \quad \text{if } l \geq L.$$

Observe that for $h \in \mathcal{R}$,

$$\begin{aligned} |h(n)| &\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} |h(x)| e^{-inx} dx \\ &\leq \|h\|_\infty. \end{aligned}$$

Hence

$$(\ast\ast\ast) \quad \left| \widehat{g}(n) - \sum_{j=1}^l \alpha_j \widehat{P}_{N_j}(n) \right| < \varepsilon \quad \text{if } l \geq L$$

Now suppose that

$$n \notin \bigcup_{j=1}^{\infty} [N_j, 3N_j],$$

Then

$$\alpha_j \widehat{P}_{N_j}(n) = 0 \quad \text{for all } j \quad \text{so}$$

$$\sum_{j=1}^l \alpha_j \widehat{P}_{N_j}(n) = 0 \quad \text{for all } l.$$

So by (***) ,

$$|\widehat{g}(n)| \leq \varepsilon.$$

Since ε is arbitrarily given , $\widehat{g}(n) = 0$.

Next assume $n \in [N_k, 3N_k]$ for some k .

Then for any $l \geq k$,


$$\sum_{j=1}^l \alpha_j \widehat{P}_{N_j}(n) = \alpha_k \widehat{P}_{N_k}(n)$$
$$= \begin{cases} \alpha_k \cdot \frac{1}{n-2N_k} & \text{if } n \neq 2N_k \\ 0 & \text{if } n = 2N_k \end{cases}$$

By (***) , we see that

$$\widehat{g}(n) = \alpha_k \widehat{P}_{N_k}(n)$$

This proves Lem 3. □

$$\bullet \quad g(x) = \alpha_1 P_{N_1}(x) + \alpha_2 P_{N_2}(x) + \dots + \alpha_k P_{N_k}(x) + \dots$$



Let us consider $S_{2N_k} g$

We notice that

$$S_{2N_k} g(x) = \alpha_1 P_{N_1}(x) + \dots + \alpha_{k-1} P_{N_{k-1}}(x) \\ + \alpha_k \cdot \tilde{P}_{N_k}(x)$$

$$S_{2N_k} g(0) = (\alpha_1 P_{N_1}(0) + \dots + \alpha_{k-1} P_{N_{k-1}}(0)) \\ + \alpha_k \cdot \tilde{P}_{N_k}(0) \\ = (I) + (II)$$

$$|(I)| \leq \sum_{i=1}^{k-1} \alpha_i \cdot M < \sum_{i=1}^{\infty} \alpha_i$$

$$\text{but } |(II)| \geq \alpha_k \log N_k \rightarrow +\infty$$

Hence

$$\left| \sum_{n_k} g(0) \right| \rightarrow +\infty \neq g(0)$$

