

Founer Analysis

Feb 24, 2022.

Review:

- We construct a cts function on the circle, say g , such that

$$S_N g(0) \not\rightarrow g(0)$$

Chap 4. Applications of Fourier series.

We are going to give the application of Fourier series in geometry, number theory, analysis and PDE.


§ 4.1 Isoperimetric inequality.

Thm 1. Let Γ be a C^1 simple closed curve in the plane.

Let l denote the length of Γ , let A denote the Area of the region bounded by Γ . Then

$$A \leq \frac{l^2}{4\pi}.$$

Moreover " $=$ " holds if and only if Γ is a circle.



$$A = \text{Area}(\Omega)$$

$$L = \text{length}(\Gamma)$$

$$\text{Then } A \leq \frac{L^2}{4\pi}.$$

Def. A parametrized curve in \mathbb{R}^2 is a mapping

$$\gamma: [a, b] \rightarrow \mathbb{R}^2$$

The image of γ , $\{\gamma(t) : t \in [a, b]\}$, is called a curve, denoted by Γ .

Def. We say $\gamma = \gamma(t)$, $t \in [a, b]$, is C^1 if

$$t \mapsto \gamma(t) \text{ is } C^1 \text{ on } [a, b], \gamma'(t) \neq 0 \text{ on } [a, b]$$

More precisely, writing $\gamma(t) = (x(t), y(t))$,
 γ is $C^1 \Leftrightarrow$ both $x(t)$ and $y(t)$ are C^1 on $[a, b]$,
and $(x'(t), y'(t)) \neq (0, 0)$ on $[a, b]$.

Fact: The length of a C^1 parametrized curve γ is given by


$$\text{length}(\gamma) = \int_a^b |\gamma'(t)| dt$$

where $|\gamma'(t)| = \sqrt{x'(t)^2 + y'(t)^2}$

for $\gamma(t) = (x(t), y(t))$, $t \in [a, b]$.

Def. A parametrized curve $\gamma: [0, l] \rightarrow \mathbb{R}^2$ is said to be parametrized by arclength if

$$|\gamma'(t)| = 1 \quad \text{for } 0 \leq t \leq l.$$



The length of the curve between $\gamma(0)$ and $\gamma(s)$

$$= \int_0^s |\gamma'(t)| dt = \int_0^s 1 dt = s$$

It is not difficult to check that any C^1 simple curve can be reparametrized by arclength.

Lem 2. Let f, g be integrable on the circle.

$$\text{Then } \langle f, g \rangle = \sum_{n=-\infty}^{\infty} \hat{f}(n) \overline{\hat{g}(n)}, \quad (*)$$

where

$$\langle f, g \rangle = \frac{1}{2\pi} \int_0^{2\pi} f(x) \overline{g(x)} dx.$$

Pf. This result is a generalization of the Parseval identity.

Indeed if $f = g$, then $(*)$ becomes the Parseval identity

To prove $(*)$, let us use the following identity

$$\langle f, g \rangle = \frac{1}{4} \left(\|f+g\|^2 - \|f-g\|^2 + i \left(\|f+ig\|^2 - \|f-ig\|^2 \right) \right)$$

which can be checked directly. Then by the Parseval identity

$$\langle f, g \rangle = \frac{1}{4} \left(\sum_{n=-\infty}^{\infty} \left(|\hat{f}(n) + \hat{g}(n)|^2 - |\hat{f}(n) - \hat{g}(n)|^2 \right) + i \left(|\hat{f}(n) + i\hat{g}(n)|^2 - |\hat{f}(n) - i\hat{g}(n)|^2 \right) \right)$$

$$= \sum_{n=-\infty}^{\infty} \hat{f}(n) \overline{\hat{g}(n)}.$$


Proof of the isoperimetric inequality:

By taking a possible transformation

$$(x, y) \rightarrow (sx, sy)$$

if necessary, we may assume that

$$\ell(\Gamma) = 2\pi$$



Parametrize Γ by its arclength,

$$\gamma = \gamma(t) : 0 \leq t \leq 2\pi$$

such that

$$|\gamma'(t)| = \sqrt{X'(t)^2 + Y'(t)^2} = 1 \text{ for } t \in [0, 2\pi]$$

To estimate $A = \text{Area}(\Omega)$, let us use Green Thm:

Green Thm: For C^1 functions $P(x,y)$ and $Q(x,y)$,

$$\oint_{\Gamma} P(x,y) dx + Q(x,y) dy = \iint_{\Omega} \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dx dy$$

In this theorem, taking $P(x,y) = 0$ and $Q(x,y) = x$ gives

$$\oint_{\Gamma} x dy = \iint_{\Omega} 1 dx dy = \text{Area}(\Omega) = A.$$

That's,

$$A = \oint_{\Gamma} x dy = \int_0^{2\pi} x(t) y'(t) dt$$

Next we need to show that

$$A = \int_0^{2\pi} x(t) y'(t) dt \leq \frac{l^2}{4\pi} = \frac{(2\pi)^2}{4\pi} = \pi.$$

Under the condition $x'(t)^2 + y'(t)^2 = 1$.