

Suggested Solution 6

- (1) In the proof of Lusin's Theorem (Theorem 2.12), it was assumed that f is non-negative, bounded and A is compact. Complete the proof by showing the conclusion still holds when f is finite a.e. and A is of finite measure.

Solution: We divide the proof into three steps.

Step 1. Assume that f is bounded and supported on a compact set A . Write $f = f^+ - f^-$. Then both f^+ and f^- are bounded and supported on A . Then by what is proved in Theorem 2.12, the conclusion of Lusin's Theorem holds in this situation.

Step 2. Assume that f is bounded and vanishes outside a measurable set A with $\mu(A) < \infty$. Let $\epsilon > 0$ be fixed. By the regularity of μ , there exists a compact set K and an open set G such that $K \subset A \subset G$ and $\mu(G \setminus K) < \frac{\epsilon}{2}$. By Urysohn's Lemma, there exists $h \in C_c(X)$ such that $K < h < G$.

Now we apply Step 1 to $f|_K$, we have there exists $g \in C_c(X)$ such that

$$\mu(\{x \in X : g(x) \neq f|_K(x)\}) < \frac{\epsilon}{2}.$$

Observe that $gh \in C_c(x)$, $gh \equiv g$ on K and $gh \equiv 0$ outside G . Hence we have

$$\{x : g(x)h(x) \neq f(x)\} \subseteq \{x : g(x) \neq f|_K(x)\} \cup (G \setminus K).$$

Therefore,

$$\begin{aligned} \mu(\{x : g(x)h(x) \neq f(x)\}) &\leq \mu(\{x : g(x) \neq f|_K(x)\}) + \mu(G \setminus K) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Step 3. Assume that f is finite a.e. and vanishes outside a measurable set A with $\mu(A) < \infty$.

For each $n \geq 1$, we define

$$f_n(x) := \begin{cases} f(x) & \text{if } |f(x)| \leq n, \\ n \cdot \text{sign} f(x) & \text{otherwise.} \end{cases}$$

Then we have $f_n(x) \rightarrow f(x)$ for every $x \in X$. Note that

$$\{x : f_n(x) \neq f(x)\} \subseteq \{x : |f(x)| > n\}.$$

Since f is finite a.e., supported on A and $\mu(A) < \infty$, we have

$$\mu(\{x : |f(x)| > n\}) \downarrow 0, \text{ as } n \rightarrow \infty.$$

Hence there exists n_0 , such that

$$\mu(\{x : f_{n_0}(x) \neq f(x)\}) < \frac{\epsilon}{2}.$$

Apply the result of Step 2 to f_{n_0} , we get a $g \in C_c(X)$ such that

$$\mu(\{x : g(x) \neq f_{n_0}(x)\}) < \frac{\epsilon}{2}.$$

Note that

$$\{x : g(x) \neq f(x)\} \subseteq \{x : g(x) = f_{n_0}(x), f_{n_0}(x) \neq f(x)\} \cup \{x : g(x) \neq f_{n_0}(x)\}.$$

Hence we have $\mu(\{x : g(x) \neq f(x)\}) \leq \epsilon$, completing the proof.

- (2) Let μ be a Riesz measure on \mathbb{R}^n . Show that for every measurable function f , there exists a sequence of continuous functions $\{f_n\}$ such that $f_n \rightarrow f$ almost everywhere.

Solution: For each $k \geq 1$, we define a set $B_k := \{x \in \mathbb{R}^n : |x| \leq k\}$ and a function

$$f_k(x) := \begin{cases} f(x) & \text{if } x \in B_k \text{ and } |f(x)| \leq k, \\ k \cdot \text{sign} f(x) & \text{if } x \in B_k \text{ and } |f(x)| > k, \\ 0 & \text{otherwise.} \end{cases}$$

Then it is easy to see that $f_k(x) \rightarrow f(x)$ at every $x \in \mathbb{R}^n$. Note that f_k is bounded and supported on a set of finite measure, we can apply the result of Exercise (1) to get a $g_k \in C_c(\mathbb{R}^n)$, such that

$$\mu(\{x \in \mathbb{R}^n : f_k(x) \neq g_k(x)\}) < \frac{1}{2^k}.$$

Let $A_k = \{x \in \mathbb{R}^n : g_k(x) \neq f_k(x)\}$. Then by the Borel-Cantelli Lemma, we have for almost every $x \in \mathbb{R}^n$, $x \in A_k$ for finite many k . As a consequence, we have $g_k \rightarrow f$ a.e..

- (3) Here we construct a Cantor-like set, or a Cantor set with positive measure, with positive measure by modifying the construction of the Cantor set as follows. Let $\{a_k\}$ be a sequence of positive numbers satisfying

$$\gamma \equiv \sum_{k=1}^{\infty} 2^{k-1} a_k < 1.$$

Construct the set \mathcal{S} so that at the k th stage of the construction one removes 2^{k-1} centrally situated open intervals each of length a_k . Establish the facts:

- (a) $\mathcal{L}^1(\mathcal{S}) = 1 - \gamma$,
- (b) \mathcal{S} is compact and nowhere dense.
- (c) \mathcal{S} is perfect hence uncountable.

Note. A set A is perfect if for every $x \in A$ and $\epsilon > 0$, $(B_\epsilon(x) \setminus \{x\}) \cap A \neq \emptyset$, that is, every point in A is an accumulation point of A . It is known that a perfect set must be uncountable.

Solution:

- (a) As the intervals removed at the same stage or different stages are mutually disjoint, we have

$$\begin{aligned} \mathcal{L}^1(\mathcal{S}) &= 1 - \sum_{k=1}^{\infty} 2^{k-1} \text{length of interval removed in the } k \text{ th stage} \\ &= 1 - \sum_{k=1}^{\infty} 2^{k-1} a_k \\ &= 1 - \gamma. \end{aligned}$$

- (b) Let S_n be the set of points left in $[0, 1]$ after the n -th level construction. Then S_n is descending and $\mathcal{S} = \bigcap_{n=1}^{\infty} S_n$. Notice that S_n is a union of 2^n mutually disjoint closed intervals hence is compact. Hence \mathcal{S} is compact. The 2^n components of S_n are of the same length

$$b_n = 2^{-n} \left(1 - \sum_{k=1}^n 2^{k-1} a_k \right).$$

Clearly $b_n \rightarrow 0$ as $n \rightarrow \infty$. Hence \mathcal{S} does not have an interior point, since otherwise \mathcal{S} will contain an open interval which is also contained in every S_n , which is impossible since $b_n \rightarrow 0$ as $n \rightarrow \infty$. Hence \mathcal{S} is nowhere dense.

- (c) If $x \in \mathcal{S}$, then x belongs some connected component of $S_n, \forall n \in \mathbb{N}$. Observe that the

end points of the 2^n intervals of S_n are in S , so $\exists y_n$ end point of one of the interval s.t.

$$|y_n - x| \leq b_n \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We have S is a perfect set.

(4) Let $0 < \varepsilon < 1$. Construct an open set $G \subset [0, 1]$ which is dense in $[0, 1]$ but $\mathcal{L}^1(G) = \varepsilon$.

Solution: Similar to the construction of Cantor's familiar "middle thirds" set. Define $K_0 = [0, 1]$ and inductively define $K_n \subset K_{n-1}$ by removing an open interval of length $2(1 - \varepsilon)2^{-2n}$. By the construction each K_n has 2^n connected components with length a_n which satisfy

$$\begin{cases} a_n = \frac{1}{2}(a_{n-1} - 2\varepsilon 2^{-2n}), & n = 1, 2, \dots \\ a_0 = 1, \end{cases}$$

from which we get $a_n = (1 - \varepsilon)2^{-n} + \varepsilon 2^{-2n}$. Thus

$$\mathcal{L}^1(K) = \lim_{n \rightarrow \infty} \mathcal{L}^1(K_n) = \lim_{n \rightarrow \infty} 2^n a_n = 1 - \varepsilon.$$

Take $G = [0, 1] \setminus K$, then $\mathcal{L}^1(G) = \varepsilon$. On the other hand, G is dense in $[0, 1]$ since the interior of K is empty.

(5) Let A be the subset of $[0, 1]$ which consists of all numbers which do not have the digit 4 appearing in their decimal expansion. Find $\mathcal{L}^1(A)$.

Solution: Let $B = \{0, 1, 2, 3, 5, 6, 7, 8, 9\}$, the set $F_0 = \{x \in [0, 1] : x = 0.a_1 a_2 a_3 \dots, a_j = 0, 1, 2, \dots, 9\} = [\frac{4}{10}, \frac{5}{10}]$ is of Lebesgue measure $\frac{1}{10}$. Fix $y_1 \in B$, $|B| = 9^1 = 9$, the set $F_{y_1} = \{x \in [0, 1] : x = 0.y_1 a_2 a_3 \dots, a_j = 0, 1, 2, \dots, 9 \forall j \geq 3\} = [\frac{y_1}{10} + \frac{4}{100}, \frac{y_1}{10} + \frac{5}{100}]$ is of Lebesgue measure $\frac{1}{100}$. Fix $(y_1, y_2) \in B^2$, $|B^2| = 9^2 = 81$, the set $F_{(y_1, y_2)} = \{x \in [0, 1] : x = 0.y_1 y_2 a_3 a_4 \dots, a_j = 0, 1, 2, \dots, 9 \forall j \geq 4\}$ is of measure $\frac{1}{1000}$. Continuing the process, we have

$$A = [0, 1] \setminus \left(\bigcup_{n=1}^{\infty} \bigcup_{(y_1, y_2, \dots, y_n) \in B^n} F_{(y_1, y_2, \dots, y_n)} \cup F_0 \right)$$

and as all $F_{(y_1, y_2, \dots, y_n)}, F_0$ are disjoint, we have

$$\begin{aligned} \mathcal{L}^1(A) &= 1 - \frac{1}{10} - \sum_{n=1}^{\infty} \sum_{(y_1, y_2, \dots, y_n) \in B^n} \frac{1}{10^{n+1}} \\ &= 1 - \frac{1}{10} - \sum_{n=1}^{\infty} \frac{9^n}{10^{n+1}} \\ &= 0. \end{aligned}$$

- (6) Let \mathcal{N} be a Vitali set in $[0, 1]$. Show that $\mathcal{M} = [0, 1] \setminus \mathcal{N}$ has measure 1 and hence deduce that

$$\mathcal{L}^1(\mathcal{N}) + \mathcal{L}^1(\mathcal{M}) > \mathcal{L}^1(\mathcal{N} \cup \mathcal{M}).$$

Remark: I have no idea what $\mathcal{L}^1(\mathcal{N})$ is, except that it is positive.

Solution: We first prove that every Lebesgue measurable subset of \mathcal{N} must be of measure zero. Let A be a Lebesgue measurable subset of \mathcal{N} , $\{A + q\}_{q \in \mathbb{Q} \cap [0, 1]}$ is a sequence of disjoint measurable set contained inside $[-1, 2]$. By translational invariance of Lebesgue measure,

$$\mathcal{L}^1\left(\bigcup_{q \in \mathbb{Q} \cap [0, 1]} A + q\right) = \sum_{q \in \mathbb{Q} \cap [0, 1]} \mathcal{L}^1(A + q) = \sum_{q \in \mathbb{Q} \cap [0, 1]} \mathcal{L}^1(A) < \infty,$$

Therefore we must have

$$\mathcal{L}^1(A) = 0.$$

We try to prove by contradiction, suppose there is an open set G s.t. $\mathcal{L}^1(G) = 1 - \varepsilon < 1$ and $G \supseteq \mathcal{N}^c$. Then $[0, 1] \setminus G$ is a measurable subset of \mathcal{N} satisfying

$$0 < \varepsilon = \mathcal{L}^1([0, 1]) - \mathcal{L}^1(G) \leq \mathcal{L}^1([0, 1] \setminus G).$$

Contradicting to our previous result.

- (7) Let E be a subset of \mathbb{R} with positive Lebesgue measure. Prove that for each $\alpha \in (0, 1)$, there exists an open interval I so that

$$\mathcal{L}^1(E \cap I) \geq \alpha \mathcal{L}^1(I).$$

It shows that E contains almost a whole interval. Hint: Choose an open G containing E

such that $\mathcal{L}^1(E) \geq \alpha \mathcal{L}^1(G)$ and note that G can be decomposed into disjoint union of open intervals. One of these intervals satisfies our requirement.

Solution: As $\exists n \in \mathbb{N}$ s.t. $\mathcal{L}^1(E \cap (-n, n)) > 0$, WLOG we may assume that E has finite outer measure, then $\forall \alpha \in (0, 1)$, \exists open G s.t. $G \supseteq E$ and

$$\mathcal{L}^1(E) + \frac{(1-\alpha)}{\alpha} \mathcal{L}^1(E) \geq \mathcal{L}^1(G),$$

Hence

$$\mathcal{L}^1(E) \geq \alpha \mathcal{L}^1(G).$$

we can write $G = \bigcup_{i=1}^{\infty} I_i$ where I_i are disjoint open intervals. Then one of these I_i must satisfy the desired property, otherwise

$$\mathcal{L}^1(E) \leq \sum_{i=1}^{\infty} \mathcal{L}^1(E \cap I_i) < \alpha \sum_{i=1}^{\infty} \mathcal{L}^1(I_i) = \alpha \mathcal{L}^1(G) < \infty,$$

contradicting the above inequality.

(8) Let E be a measurable set in \mathbb{R} with respect to \mathcal{L}^1 and $\mathcal{L}^1(E) > 0$. Show that $E - E$ contains an interval $(-a, a)$, $a > 0$. Hint:

- (a) U, V open, with finite measure, $x \mapsto \mathcal{L}^1((x+U) \cap V)$ is continuous on \mathbb{R} .
 (b) A, B measurable, $\mu(A), \mu(B) < \infty$, then $x \mapsto \mathcal{L}^1((x+A) \cap B)$ is continuous. For $A \subset U, B \subset V$, try

$$|\mathcal{L}^1((x+U) \cap V) - \mathcal{L}^1((x+A) \cap B)| \leq \mathcal{L}^1(U \setminus A) + \mathcal{L}^1(V \setminus B).$$

- (c) Finally, $x \mapsto \mathcal{L}^1((x+E) \cap E)$ is positive at 0 and if $(x+E) \cap E \neq \emptyset$, then $x \in E - E$.

Solution:

- (a) We prove the case when U is an open interval I , note for all subset A, B of \mathbb{R} ,

$$((x+A) \cap B) \setminus ((y+A) \cap B) = (x+A) \setminus (y+A) \cap B.$$

Therefore

$$|\mathcal{L}^1((x+I) \cap V) - \mathcal{L}^1((y+I) \cap V)| \leq \mathcal{L}^1((x+I) \setminus (y+I)) + \mathcal{L}^1((y+I) \setminus (x+I)) \leq 4|x-y|.$$

the function is Lipschitz and continuous. In general U can be written as countable union of disjoint open intervals $\{I_i\}$, as $\sum_{i=1}^{\infty} \ell(I_i) < \infty, \exists N$ s.t. for all $k \geq N$,

$$\sum_{i=k}^{\infty} \ell(I_i) < \varepsilon.$$

We have

$$\sum_{i=1}^{\infty} \mathcal{L}^1((x+I_i) \cap V) - \mathcal{L}^1((y+I_i) \cap V) \leq \sum_{i=1}^k \mathcal{L}^1((x+I_i) \cap V) - \mathcal{L}^1((y+I_i) \cap V) + 2\varepsilon < 3\varepsilon$$

for x sufficiently close to y . Similarly

$$\sum_{i=1}^{\infty} \mathcal{L}^1((y+I_i) \cap V) - \mathcal{L}^1((x+I_i) \cap V) \leq 3\varepsilon.$$

We have the function $\mathcal{L}^1((x+U) \cap V)$ is continuous.

(b) Obviously, $((x+U) \cap V) \setminus ((x+A) \cap B) \subseteq U \setminus A \cup V \setminus B$. Therefore, we have

$$0 \leq \mathcal{L}^1((x+U) \cap V) - \mathcal{L}^1((x+A) \cap B) \leq \mathcal{L}^1(U \setminus A) + \mathcal{L}^1(V \setminus B).$$

Note RHS is independent on x, y , so the result follow from outer regularity of Lebesgue measure.

(c) the function $\mathcal{L}^1((x+E) \cap E)$ is continuous and positive at 0, $\exists a > 0$ s.t the function remain positive on $(-a, a)$, i.e

$$(x+E) \cap E \neq \emptyset$$

and $\forall x \in (-a, a), \exists e_1 e_2 \in E$ s.t

$$x = e_1 - e_2 \in E - E.$$

Alternate proof. The following is a simple proof due to Karl Stromberg.

By the regularity of \mathcal{L}^1 , for every $\varepsilon > 0$ there are a compact set $K \subset E$ and an open set $U \supset E$ such that

$$\mathcal{L}^1(K) + \varepsilon > \mathcal{L}^1(E) > \mathcal{L}^1(U) - \varepsilon.$$

For our purpose it is enough to choose K and U such that

$$2\mathcal{L}^1(K) > \mathcal{L}^1(U).$$

Since $K \subset U$, there is an open cover of K that is contained in U . Since K is compact, one can choose a small neighborhood V of 0 such that

$$K + V \subset U.$$

Let $v \in V$, and suppose

$$(K + v) \cap K = \emptyset.$$

Then,

$$2\mathcal{L}^1(K) = \mathcal{L}^1(K + v) + \mathcal{L}^1(K) < \mathcal{L}^1(U),$$

contradicting our choice of K and U . Hence for all $v \in V$ there exists $k_1, k_2 \in K \subset E$ such that

$$k_1 + v = k_2,$$

which means that $V \subset E - E$.

- (9) Give an example of a continuous map ϕ and a measurable f such that $f \circ \phi$ is not measurable. Hint: May use the function $h = x + g(x)$ where g is the Cantor function as ϕ .

Solution: Let $h = x + g(x)$ where g is the Cantor function. Then $h : [0, 1] \rightarrow [0, 2]$ is a strictly monotonic and continuous map, so its inverse $\phi = h^{-1}$ is continuous too. Since g is constant on every interval in the complement of C , one has that h maps such an interval to an interval of the same length. Hence $\mu(h(C)) = 1$, where C is the cantor set. Then $h(C)$ contains a non-measurable set A due to Proposition 3.3. Let $B = \phi(A)$. Set $f = \chi_B$. Then $f \circ \phi$ is not measurable since its inverse image of 1 is A .