

MATH 5061 Lecture 14 (Apr 21)

Reminder: Take-home final Apr 28 2:30 PM - May 5 2:30 PM
(available and submit via Blackboard.)

Last time:

Rauch comparison theorem


Let (M^n, g) , $(\tilde{M}^{n+k}, \tilde{g})$ be complete Riem. manifolds

• $\gamma: [0, a] \rightarrow M$, $\tilde{\gamma}: [0, a] \rightarrow \tilde{M}$ be geodesic in M and \tilde{M}
st. $|\gamma'| = |\tilde{\gamma}'|$ (ie. same speed)


• J, \tilde{J} are normal Jacobi fields along $\gamma, \tilde{\gamma}$ respectively

st. $J(0) = 0 = \tilde{J}(0)$, $|J'(0)| = |\tilde{J}'(0)|$

(M^n, g)



$(\tilde{M}^{n+k}, \tilde{g})$



ASSUME: ① \nexists conjugate pt along $\tilde{\gamma}$ on $[0, a]$

* ② $\forall t \in [0, a]$, $\forall x \in T_{\gamma(t)}M$, $\forall \tilde{x} \in T_{\tilde{\gamma}(t)}\tilde{M}$.

we have $K_{\gamma(t)}(\text{span}\{\gamma'(t), x\}) \leq \tilde{K}_{\tilde{\gamma}(t)}(\text{span}\{\tilde{\gamma}'(t), \tilde{x}\})$

THEN: $|J(t)| \geq |\tilde{J}(t)| \quad \forall t \in [0, a]$

Last week, we proved the

Index Lemma: On a geodesic segment without conjugate points,

$$I(J, J) \leq I(V, V) \quad \text{where } J \text{ and } V \text{ agree at the end points and } J \text{ is a Jacobi field.}$$

Proof of Rauch comparison Thm:

WLOG, assume $|J'(0)| = |\tilde{J}'(0)| > 0$.

Denote: $v(t) := |J(t)|^2$ and $\tilde{v}(t) := |\tilde{J}(t)|^2$

Goal: $v(t) \geq \tilde{v}(t) \quad \forall t \in [0, a]$

Observe, by assumption \odot , $\tilde{v}(t) > 0 \quad \forall t \in (0, a]$. So we define

$$h(t) := \frac{v(t)}{\tilde{v}(t)} \quad t \in (0, a] \quad \text{Goal: } h(t) \geq 1 \quad \forall t \in (0, a]$$

By L'Hospital's Rule,


$$v'(t) = 2 \langle J(t), J'(t) \rangle$$

$$\tilde{v}''(t) = 2 \langle J'(t), J'(t) \rangle + 2 \langle J(t), J''(t) \rangle$$

$$\lim_{t \rightarrow 0} h(t) = \lim_{t \rightarrow 0} \frac{v'(t)}{\tilde{v}''(t)} = \lim_{t \rightarrow 0} \frac{v''(t)}{\tilde{v}''(t)} = \frac{|J'(0)|^2}{|\tilde{J}'(0)|^2} = 1$$

Claim: $h'(t) \geq 0 \quad \forall t \in (0, a)$

i.e. $v' \tilde{v} \geq v \tilde{v}' \quad \forall t \in (0, a)$ (*)



Pf of Claim: Fix any $t_0 \in (0, a)$.

Case 1: $v(t_0) = 0 \quad (\Rightarrow J(t_0) = 0)$

Then, $v'(t_0) = 2 \langle J(t_0), J'(t_0) \rangle = 0$

Clearly, (*) holds.

Case 2: $v(t_0) \neq 0$ ($\Rightarrow v(t_0) = |J(t_0)|^2 > 0$)

By rescaling, we obtain two new Jacobi fields

$$U(t) := \frac{1}{\sqrt{v(t_0)}} J(t) \quad ; \quad \tilde{U}(t) := \frac{1}{\sqrt{\tilde{v}(t_0)}} \tilde{J}(t)$$

Observe: $\frac{v'(t_0)}{v(t_0)} = 2 I_{t_0}(U, U)$ AND $\frac{\tilde{v}'(t_0)}{\tilde{v}(t_0)} = 2 I_{t_0}(\tilde{U}, \tilde{U})$

Why? $2 I_{t_0}(U, U) := 2 \int_0^{t_0} \langle U', U' \rangle - R(\gamma', U, \gamma', U) dt$
 $= \int_0^{t_0} (\langle U, U \rangle)' dt \stackrel{\text{by part}}{=} \langle U, U \rangle'(t_0)$
 $= 2 \langle U(t_0), U'(t_0) \rangle = 2 \frac{\langle J(t_0), J'(t_0) \rangle}{v(t_0)} = \frac{v'(t_0)}{v(t_0)}$

So, it suffices to show

$$I_{t_0}(\tilde{U}, \tilde{U}) \leq I_{t_0}(U, U) \quad (*)$$

Fix parallel O.N.B. along $\gamma, \tilde{\gamma}$:

$$\{ e_1 = \frac{\gamma'}{|\gamma'|}, e_2, \dots, e_n \}_{(t)} \text{ for } T_{\gamma(t)} M^n \text{ st. } e_2(t_0) = U(t_0)$$

$$\{ \tilde{e}_1 = \frac{\tilde{\gamma}'}{|\tilde{\gamma}'|}, \tilde{e}_2, \dots, \tilde{e}_{n+k} \}_{(t)} \text{ for } T_{\tilde{\gamma}(t)} \tilde{M}^{n+k} \text{ st. } \tilde{e}_2(t_0) = \tilde{U}(t_0)$$

Define a map $\phi : \left\{ \begin{array}{l} \text{vector field} \\ \text{along } \gamma(t) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{vector field} \\ \text{along } \tilde{\gamma}(t) \end{array} \right\}$

$$V(t) = \sum_{i=1}^n g_i(t) e_i(t) \xrightarrow{\phi} (\phi V)(t) := \sum_{i=1}^n g_i(t) \tilde{e}_i(t)$$

← same coefficients →

Properties of ϕ :

- $\langle \phi V, \phi V \rangle = \langle V, V \rangle \quad \forall t \in [0, a]$
- $(\phi V)' = \phi(V')$ $\forall t \in [0, a]$
- * $I_{t_0}(\phi V, \phi V) \leq I_{t_0}(V, V)$

Reason: $I_{t_0}(V, V) := \int_0^{t_0} |V'(t)|^2 - R(\gamma', V, \gamma', V) dt$

\wedge

$I_{t_0}(\phi V, \phi V) := \int_0^{t_0} |(\phi V)'|^2 - \tilde{R}(\tilde{\gamma}', \phi V, \tilde{\gamma}', \phi V) dt$

|| by first 2 properties || by 1st property and $K \leq \tilde{K}$

Recall by construction, we have

$$(\phi U)(t_0) = \phi(e_2(t_0)) = \tilde{e}_2(t_0) = \tilde{U}(t_0)$$

i.e. ϕU and \tilde{U} have the same boundary values along $\tilde{\gamma}$ on $[0, t_0]$

By (*) and Index Lemma,

$$\frac{1}{2} \frac{\tilde{U}'(t_0)}{\tilde{U}(t_0)} = I_{t_0}(\tilde{U}, \tilde{U}) \leq I_{t_0}(\phi U, \phi U) \stackrel{(*)}{\leq} I_{t_0}(U, U) = \frac{1}{2} \frac{U'(t_0)}{U(t_0)}$$

□

Volume comparison via Jacobi field estimates

Thm: (Bishop-Gromov & Gunther)

Let (M^n, g) be a complete Riem. mfd.

Fix $p \in M$, let $a, b \in \mathbb{R}$ be constants.

(i) If $\text{Ric}_x^M(u, u) \geq (n-1)a \quad \forall x \in M, \forall u \in T_x M, |u| = 1,$

then $\text{Vol}(B_p^M(r)) \leq \text{Vol}(B^{\text{ll}_a}(r)) \quad \forall r \in (0, \text{inj}_p(M)).$

Here, $B^{\text{ll}_a}(r)$ = volume of ball of radius r
inside a space form ll_a of const. curvature a

(ii) If $K^M \leq b$, then

$\text{Vol}(B_p^M(r)) \geq \text{Vol}(B^{\text{ll}_b}(r)) \quad \forall r \in (0, \text{inj}_p(M)).$

Digestion: Jacobi field and Volume estimates

Volume form of (M^n, g) : $dV_g = \sqrt{\det(g_{ij})} dx^1 \wedge \dots \wedge dx^n$
in local coord. (x^1, \dots, x^n) .


and $\text{Vol}(U^M) = \int_U \sqrt{\det(g_{ij})} dx^1 \dots dx^n$

In particular, in geodesic normal coordinates (based at $p \in M$)


$$\text{Vol}(B_p^M(r)) = \int_{S^{n-1}} \int_0^r J(u, t) \cdot t^{n-1} dt du$$

where $J(u, t) := t^{-(n-1)} \sqrt{\det \langle \underline{J}_i(t), \underline{J}_j(t) \rangle}$
some Jacobi fields

Picture:



\exp_p



Fix $u \in T_p M$, $|u| = 1$, and a geodesic $\gamma(t) = \exp_p(tu)$

Fix an O.N.B. $\{u = e_1, e_2, \dots, e_n\}$ of $T_p M$

Let $J_i(t)$ be the Jacobi field along $\gamma(t)$ s.t.

$$J_i(0) = 0 \quad \text{and} \quad J_i'(0) = e_i$$

i.e. $J_i(t) = d(\exp_p)_{tu}(te_i)$

$$\text{So, } dV_g = \sqrt{\det(g_{ij})} dx^1 \dots dx^n$$

$$= \sqrt{\det \left(\underbrace{d \exp(e_i)}_{\frac{1}{t} J_i(t)}, \underbrace{d \exp(e_j)}_{\frac{1}{t} J_j(t)} \right)} t^{n-1} dt du$$

$$= J(u, t) t^{n-1} dt du$$

On the other hand, we have explicit formulae for the

Jacobi fields in space forms \mathcal{M}_a


parallel v.f.

Jacobi eqⁿ: $J'' + a J = 0 \Rightarrow J(t) = S_a(t) E(t)$
and $J(0) = 0$

where $S_a(t) := \begin{cases} \sin \sqrt{a} t & \text{if } a > 0 \\ t & \text{if } a = 0 \\ \sinh \sqrt{-a} t & \text{if } a < 0 \end{cases}$

Proof of Volume Comparison Thm (i) : Ric $\geq (n-1) a$

Picture:



Denote $Y_i^r(t)$ to be the unique Jacobi field st.

$$Y_i^r(0) = 0 \quad \text{and} \quad Y_i^r(r) = E_i(r)$$

and $v_i \in T_p M$ st. $d(\exp_p)_{tu}(t v_i) = Y_i^r(t)$

Note: $J(u, t) = C_r t^{1-n} \det(Y_2^r(t), \dots, Y_n^r(t))$

(Ex)

where $C_r = [\det(Y_2^r(0), \dots, Y_n^r(0))]$

Denote: For each $u \in T_p M$, $|u| = 1$,

$$f(t) := J(u, t)$$

Lemma: $\frac{f'(r)}{f(r)} = \sum_{i=2}^n I_r(Y_i^r, Y_i^r) - \frac{n-1}{r}$

Proof of Lemma:

Let $S(t) := \det(\langle Y_i^r(t), Y_j^r(t) \rangle)$. Then

$$f(t) := J(u, t) := C_r t^{1-n} \det(Y_2^r(t), \dots, Y_n^r(t)) = C_r t^{1-n} \sqrt{S(t)}$$

$$\Rightarrow f'(t) = C_r t^{1-n} \cdot \frac{1}{2} \frac{g'(t)}{\sqrt{g(t)}} - C_r (n-1) t^{-n} \sqrt{g(t)}$$

$$\Rightarrow \frac{f'(t)}{f(t)} = \frac{g'(t)}{2g(t)} - \frac{n-1}{t}$$

On the other hand, we want $\frac{g'(r)}{2g(r)} = \sum_{i=1}^n \text{I}_r(Y_i^r, Y_i^r)$.

Note: $g(r) := \det(\langle Y_i^r(r), Y_j^r(r) \rangle) = \det \langle E_i(r), E_j(r) \rangle = 1$

and $g'(r) = 2 \sum_{i=1}^n \langle (Y_i^r)'(r), Y_i^r(r) \rangle$

By the proof of Claim of Index Lemma, $\text{I}_r(Y_i^r, Y_i^r) = \langle Y_i^r(r), Y_i^r(r) \rangle$.

Lemma \square

To finish the proof of Volume Comparison Thm (i),

Consider the vector field along γ

$$X_i^r(t) := \frac{S_a(t)}{S_a(r)} E_i(t)$$

s.t. $X_i^r(0) = 0 = Y_i^r(0)$ and $X_i^r(r) = E_i(r) = Y_i^r(r)$.

By Index Lemma, $\text{I}_r(Y_i^r, Y_i^r) \leq \text{I}_r(X_i^r, X_i^r)$.

Note: $\text{I}_r(X_i^r, X_i^r) = \int_0^r \left(\frac{S_a'(t)}{S_a(r)} \right)^2 - \left(\frac{S_a(t)}{S_a(r)} \right)^2 R(\gamma', E_i, \gamma', E_i) dt$

$$= \int_0^r \left(\frac{S_a'(t)}{S_a(r)} \right)^2 \underbrace{[a - R(\gamma', E_i, \gamma', E_i)]}_{\sum_{i=1}^n (\cdot) \leq 0 \text{ by assumption}} dt + \frac{S_a'(r)}{S_a(r)}$$

$$\Rightarrow \frac{f'(r)}{f(r)} = \sum_{i=1}^n \text{I}_r(Y_i^r, Y_i^r) - \frac{n-1}{r} \leq \sum_{i=1}^n \text{I}_r(X_i^r, X_i^r) - \frac{n-1}{r}$$

Integrate in $r \Rightarrow f(r) \leq S_a(r)$. \Rightarrow DONE!
integrate again $\frac{1}{(n-1)} \left(\frac{S_a'(r)}{S_a(r)} - \frac{1}{r} \right)$ \square