


## Chapter 4: Differentiation I

**Learning Objectives:**

- (1) Define the derivatives, and study its basic properties.
- (2) Study the relationship between differentiability and continuity.
- (3) Use the constant multiple rule, sum rule, power rule, product rule, quotient rule and chain rule to find derivatives.
- (4) Explore logarithmic differentiation.

**4.1 Motivation & Definition**

**Motivation from physics: velocity** Suppose an object is moving along  $x$ -axis from the origin to right. Let  $S = S(t)$  be the position of the object at time  $t$ . What is the average velocity of this object from  $t = 1$  to  $t = 2$ ?



$$\begin{aligned}
 \text{Average velocity from } t = 1 \text{ to } t = 2 &= \frac{\text{Change of position}}{\text{Change of time}} \\
 &= \frac{\Delta S}{\Delta t} \\
 &= \frac{S(2) - S(1)}{2 - 1} \\
 &= \text{slope of secant line passing through } (1, S(1)) \text{ and } (2, S(2))
 \end{aligned}$$


**Question:** What is the instantaneous velocity at  $t = 1$ ?

Idea: Average velocity from  $t = 1$  to  $t = 1 + \Delta t$  is  $\frac{S(1 + \Delta t) - S(1)}{\Delta t}$ , where  $\Delta t$  is small.

Let  $\Delta t \rightarrow 0$ , the instantaneous velocity at  $t = 1$  is defined to be

$$S'(1) = \lim_{\Delta t \rightarrow 0} \frac{S(1 + \Delta t) - S(1)}{\Delta t},$$

which is called the **derivative** of  $S$  at  $t = 1$ .  $S'(1)$  describes the **rate of change** of  $S(t)$  at  $t = 1$ .



*Remark. Terminology:* The term “velocity” takes the direction of motion into account; it can be positive or negative. The term “speed” only takes into account the rate of change, disregarding the direction. It is the absolute value of the velocity.

**Definition 4.1.1.** The **derivative** of  $f(x)$  is the function

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}. \quad (4.1)$$

The process of computing the derivative is called **differentiation**, and we say that  $f(x)$  is **differentiable** at  $x = x_0$  if  $f'(x_0)$  exists; that is,  $\lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x}$  exists.

*Remark.* 1. By definition, if  $f(x_0)$  is not well-defined, we cannot define  $f'(x_0)$ . So  $f(x)$  must not be differentiable at  $x = x_0$ .

2. Another equivalent formula:

$$f'(x_0) = \lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x} = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}.$$

3.

$$\frac{\Delta f}{\Delta x} = \frac{f(x) - f(x_0)}{x - x_0}$$

is called **difference quotient**.

4.  $f'(x_0)$  describes the rate of change of  $f(x)$  at  $x = x_0$ .

5. When we say that we use **the first principle** to find derivatives, we mean that we use the definition (4.1) to find the derivative. However, later we will learn faster techniques to find derivatives.

**Geometrical interpretation of differentiation:**  $f'(x_0)$  is the slope of tangent line to the curve of  $f(x)$  at  $x = x_0$ .

**Example 4.1.1.** Let  $f(x) = x^2$ . Then (i) prove that  $f(x)$  is differentiable at  $x = 1$ ; (ii) find  $f'(1)$  and the equation of the tangent line to the graph of  $f$  at  $x = 1$ .

*Solution.* (i) By the definition, at  $x = 1$

$$\begin{aligned} \lim_{\Delta x \rightarrow 0} \frac{f(1 + \Delta x) - f(1)}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{(1 + \Delta x)^2 - 1^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2 + \Delta x) \\ &= 2, \end{aligned}$$

So,  $f$  is differentiable at  $1$ , and  $f'(1) = 2$ .

(ii) The tangent line passes through  $(1, f(1)) = (1, 1)$  with slope  $f'(1) = 2$ . So, the equation of the tangent line is

$$\frac{y - f(1)}{x - (1)} = 2.$$

Thus

$$y = 2x - 1.$$

■

**Definition 4.1.2.** If  $f(x) : A \rightarrow \mathbb{R}$  is differentiable at every point  $x \in A$ , then  $f(x)$  is said to be a differentiable function in  $A$ , and the derivative function  $f'(x) : A \rightarrow \mathbb{R}$  is well-defined.

**Example 4.1.2.** Let  $f(x) = x^2$ . Prove that  $f(x)$  is differentiable on  $\mathbb{R}$ , and find  $f'(x)$ .

*Solution.* For any  $x \in \mathbb{R}$ ,

$$\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 - x^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x.$$

So,  $f$  is differentiable at  $x$ , and  $f'(x) = 2x$ . ■

**Notation:** For  $y = f(x) = x^2$ ,

$$f'(x) = \frac{dy}{dx} = \frac{df}{dx} = 2x; \quad f'(4) = \left. \frac{dy}{dx} \right|_{x=4} = \left. \frac{df}{dx} \right|_{x=4} = 2 \cdot 4 = 8.$$

*Question* Where does the minimum of  $x^2$  occur? (Hint: what is the slope of the tangent line at the minimum?)

**Example 4.1.3.** Let  $f(x) = \frac{x+1}{x-1}$ . Using the definition of derivatives, compute  $f'(x)$  for  $x \neq 1$ .

*Solution.*

$$\begin{aligned} f(x + \Delta x) - f(x) &= \frac{x + \Delta x + 1}{x + \Delta x - 1} - \frac{x + 1}{x - 1} \\ &= \frac{(x - 1)(x + \Delta x + 1) - (x + 1)(x + \Delta x - 1)}{(x - 1)(x + \Delta x - 1)} \\ &= \frac{-2\Delta x}{(x - 1)(x + \Delta x - 1)}. \end{aligned}$$

Therefore

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{-2}{(x - 1)(x + \Delta x - 1)} \\ &= \frac{\lim_{\Delta x \rightarrow 0} (-2)}{\lim_{\Delta x \rightarrow 0} (x - 1)(x + \Delta x - 1)} = \frac{-2}{(x - 1)^2}. \end{aligned}$$
■

**Example 4.1.4.** Find the derivative of  $f(x) = \sqrt{x}$  for  $x > 0$ .

*Solution.*

$$\begin{aligned} \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x} - \sqrt{x}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(\sqrt{x + \Delta x} - \sqrt{x})(\sqrt{x + \Delta x} + \sqrt{x})}{\Delta x(\sqrt{x + \Delta x} + \sqrt{x})} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x} + \sqrt{x}} \\ &= \frac{1}{2\sqrt{x}}. \end{aligned}$$

So,  $(x^{\frac{1}{2}})' = \frac{1}{2}x^{-\frac{1}{2}}, x > 0$ . ■

**Example 4.1.5.** Find the derivative of  $f(x) = \sqrt[3]{x}$ .

**Hint:**  $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$ .

*Solution.* For any  $x \neq 0$ ,

$$\begin{aligned} \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{\sqrt[3]{x + \Delta x} - \sqrt[3]{x}}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{(\sqrt[3]{x + \Delta x} - \sqrt[3]{x})((\sqrt[3]{x + \Delta x})^2 + \sqrt[3]{x + \Delta x} \cdot \sqrt[3]{x} + (\sqrt[3]{x})^2)}{\Delta x((\sqrt[3]{x + \Delta x})^2 + \sqrt[3]{x + \Delta x} \cdot \sqrt[3]{x} + (\sqrt[3]{x})^2)} \\ &= \lim_{h \rightarrow 0} \frac{x + \Delta x - x}{\Delta x((\sqrt[3]{x + \Delta x})^2 + \sqrt[3]{x + \Delta x} \cdot \sqrt[3]{x} + (\sqrt[3]{x})^2)} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{(\sqrt[3]{x + \Delta x})^2 + \sqrt[3]{x + \Delta x} \cdot \sqrt[3]{x} + (\sqrt[3]{x})^2} \\ &= \frac{1}{3(\sqrt[3]{x})^2} = \frac{1}{3}x^{-\frac{2}{3}}. \end{aligned}$$

For  $x = 0$ ,

$$\lim_{\Delta x \rightarrow 0} \frac{f(0 + \Delta x) - f(0)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\sqrt[3]{\Delta x} - \sqrt[3]{0}}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{1}{(\Delta x)^{\frac{2}{3}}} \text{ does not exist.}$$

So,

$$(x^{1/3})' = \begin{cases} \frac{1}{3}x^{-\frac{2}{3}}, & x \neq 0 \\ \text{Not exist at } x = 0, \text{ i.e. } x^{\frac{1}{3}} \text{ not differentiable at } 0 \end{cases}$$
■

**Example 4.1.6.** Discuss the differentiability of  $f(x) = |x|$ .

*Solution.* For  $x_0 > 0$ ,

$$\lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{(x_0 + \Delta x) - x_0}{\Delta x} = 1.$$

For  $x_0 < 0$ ,

$$\lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x) - f(x_0)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{-(x_0 + \Delta x) - (-x_0)}{\Delta x} = -1.$$

For  $x_0 = 0$ .

$$\lim_{\Delta x \rightarrow 0^+} \frac{f(0 + \Delta x) - f(0)}{\Delta x} = \lim_{\Delta x \rightarrow 0^+} \frac{\Delta x}{\Delta x} = 1.$$

$$\lim_{\Delta x \rightarrow 0^-} \frac{f(0 + \Delta x) - f(0)}{\Delta x} = \lim_{\Delta x \rightarrow 0^-} \frac{-\Delta x}{\Delta x} = -1.$$

$1 \neq -1$ , so  $f$  is not differentiable at  $x = 0$ . So,

$$(|x|)' = \begin{cases} 1 & \text{if } x > 0, \\ \text{undefined} & \text{if } x = 0. \\ -1 & \text{if } x < 0, \end{cases}$$

■

## 4.2 Properties of derivatives

### 4.2.1 Differentiation and Continuity

**Proposition 1.**  $f(x)$  is differentiable at  $x = x_0 \implies f(x)$  is continuous at  $x = x_0$ .

*Proof.* Suppose  $f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$  exists, then

$$\begin{aligned} \lim_{x \rightarrow x_0} (f(x) - f(x_0)) &= \lim_{x \rightarrow x_0} \left( \frac{f(x) - f(x_0)}{x - x_0} \cdot (x - x_0) \right) \\ &= \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \cdot \lim_{x \rightarrow x_0} (x - x_0) \\ &= f'(x_0) \cdot 0 = 0. \end{aligned}$$

So,  $\lim_{x \rightarrow x_0} f(x) = \lim_{x \rightarrow x_0} (f(x) - f(x_0)) + \lim_{x \rightarrow x_0} f(x_0) = 0 + f(x_0) = f(x_0)$ , that is,  $f(x)$  is continuous at  $x_0$ . □

**The converse is not true.** For example, let  $f(x) = |x|$ . It is not differentiable at  $x = 0$  but is continuous at  $x = 0$ .

**Exercise 4.2.1.** Let

$$f(x) = \begin{cases} x^2 - 1, & \text{if } x \geq 1 \\ 1 - x, & \text{if } x < 1 \end{cases}$$

- (a) Show that  $f(x)$  is continuous at  $x = 1$ .  
 (b) Show that  $f(x)$  is differentiable everywhere except  $x = 1$ , and

$$f'(x) = \begin{cases} 2x, & \text{if } x > 1 \\ \text{undefined}, & \text{if } x = 1 \\ -1, & \text{if } x < 1 \end{cases}$$

## 4.2.2 Differentiation and Arithmetic Operations

**Theorem 2.** Let  $f(x)$  and  $g(x)$  be differentiable functions. Then

(1) *Sum rule:*  $(f + g)'(x) = f'(x) + g'(x)$ .

(2) *Difference rule:*  $(f - g)'(x) = f'(x) - g'(x)$ .

(3) *Product rule:*  $(fg)'(x) = f'(x)g(x) + f(x)g'(x)$ .

(4) *Quotient rule:*  $\left(\frac{f}{g}\right)'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$ .

*Proof.* (1)

$$\begin{aligned} (f + g)'(x) &= \lim_{\Delta x \rightarrow 0} \frac{(f + g)(x + \Delta x) - (f + g)(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) + g(x + \Delta x) - (f(x) + g(x))}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} + \lim_{\Delta x \rightarrow 0} \frac{g(x + \Delta x) - g(x)}{\Delta x} \\ &= f'(x) + g'(x). \end{aligned}$$

(3)

$$\begin{aligned}
(fg)'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x)g(x + \Delta x) - f(x)g(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x)g(x + \Delta x) - f(x + \Delta x)g(x) + f(x + \Delta x)g(x) - f(x)g(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x)g(x + \Delta x) - f(x + \Delta x)g(x)}{\Delta x} + \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x)g(x) - f(x)g(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} f(x + \Delta x) \cdot \frac{g(x + \Delta x) - g(x)}{\Delta x} + \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \cdot g(x) \\
&= \lim_{\Delta x \rightarrow 0} f(x + \Delta x) \cdot \lim_{\Delta x \rightarrow 0} \frac{g(x + \Delta x) - g(x)}{\Delta x} + \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \cdot \lim_{\Delta x \rightarrow 0} g(x) \\
&= f(x)g'(x) + f'(x)g(x).
\end{aligned}$$

*Remark.* Here we used:

$$g(x) \text{ is differentiable at } x \quad \Rightarrow \quad g(x) \text{ is continuous at } x$$

$$\text{so, } \lim_{\Delta x \rightarrow 0} f(x + \Delta x) = f(x).$$

□

**Exercise 4.2.2.** Prove other rules using the first principle.

*Remark.* 1. The product rule is more commonly referred to as the *Leibniz rule*.

Caveat:  $(f \cdot g)' \neq f' \cdot g'$ !

2. The quotient rule (4) can be derived from the Leibniz rule together with the chain rule (Section 4.3).

### 4.2.3 Derivatives of Elementary Functions

**Theorem 3** (Constant functions).

$$\boxed{f(x) = k \quad \Rightarrow \quad f'(x) = 0}$$

*Proof.*

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{k - k}{\Delta x} = 0.$$

□



As a consequence, we have

$$\boxed{(kf(x))' = (k)'f(x) + kf'(x) = kf'(x)}, \quad \text{for any constant } k.$$

*Remark.* It can also be proved by the first principle.

**Theorem 4** (The Power Rule).

$$\boxed{(x^a)' = ax^{a-1}}, \quad \text{whenever it is well-defined, } a \in \mathbb{R}.$$

*Proof.* We will only prove the special case when  $n$  is an integer.

Recall

$$x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + \cdots + xy^{n-2} + y^{n-1}).$$

So

$$(x + \Delta x)^n - x^n = (x + \Delta x - x)((x + \Delta x)^{n-1} + (x + \Delta x)^{n-2}x + \cdots + (x + \Delta x)x^{n-2} + x^{n-1}).$$

We have

$$\begin{aligned} \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^n - x^n}{\Delta x} &= \lim_{\Delta x \rightarrow 0} ((x + \Delta x)^{n-1} + (x + \Delta x)^{n-2}x + \cdots + (x + \Delta x)x^{n-2} + x^{n-1}) \\ &= x^{n-1} + x^{n-2}x + \cdots + xx^{n-2} + x^{n-1} = nx^{n-1}. \end{aligned}$$

□

*Remark.* Alternatively, combine the fact that  $x' = 1$  and the Leibniz rule.

**Example 4.2.1.**

$$\begin{aligned} (x^3)' &= 3x^2, & x \in \mathbb{R} \\ (\sqrt{x})' &= \frac{1}{2}x^{-\frac{1}{2}}, & x > 0. \quad \text{Caution: } x \text{ can not be 0.} \\ (\sqrt[3]{x})' &= \frac{1}{3}x^{-\frac{2}{3}}, & x \neq 0. \quad \text{Caution: } x \text{ can be negative.} \\ (x^{\frac{3}{2}})' &= \frac{3}{2}x^{\frac{1}{2}}, & x > 0. \end{aligned}$$

**Theorem 5** (Exponential functions and Logarithmic functions).

$$\boxed{(e^x)' = e^x; \quad (a^x)' = a^x \ln a,} \quad a > 0, a \neq 1, x \in \mathbb{R}.$$

$$\boxed{(\ln x)' = \frac{1}{x}; \quad (\log_a x)' = \frac{1}{x \ln a},} \quad a > 0, a \neq 1, x > 0.$$

*Proof.* (Optional!)

$$\begin{aligned}
 (\ln x)' = \frac{1}{x} &\iff \lim_{\Delta x \rightarrow 0} \frac{\ln(x + \Delta x) - \ln x}{\Delta x} = \frac{1}{x} \\
 &\iff \lim_{\Delta x \rightarrow 0} \frac{\ln(1 + \frac{\Delta x}{x})}{\frac{\Delta x}{x}} = 1 \\
 &\iff \lim_{y \rightarrow 0} \ln(1 + y)^{\frac{1}{y}} = 1, \quad (\text{change variable: } y := \frac{\Delta x}{x}) \\
 &\iff \lim_{y \rightarrow 0} (1 + y)^{\frac{1}{y}} = e \\
 &\iff \lim_{z \rightarrow +\infty} \left(1 + \frac{1}{z}\right)^z = \lim_{y \rightarrow 0^+} (1 + y)^{\frac{1}{y}} = e \quad (\text{change variable: } z = \frac{1}{y}) \\
 &\quad \text{and } \lim_{z \rightarrow -\infty} \left(1 + \frac{1}{z}\right)^z = \lim_{y \rightarrow 0^-} (1 + y)^{\frac{1}{y}} = e \quad (\text{definition of } e).
 \end{aligned}$$

$$\begin{aligned}
 (e^x)' = e^x &\iff \lim_{\Delta x \rightarrow 0} \frac{e^{x+\Delta x} - e^x}{\Delta x} = e^x \\
 &\iff \lim_{\Delta x \rightarrow 0} \frac{e^{\Delta x} - 1}{\Delta x} = 1 \\
 &\iff \lim_{y \rightarrow 0} \frac{y}{\ln(1 + y)} = 1, \quad (\text{let } y = e^{\Delta x} - 1) \\
 &\iff \lim_{y \rightarrow 0} \frac{\ln(1 + y)}{y} = \left. \frac{d \ln x}{dx} \right|_{x=1} = 1.
 \end{aligned}$$

For general  $a$ : The formulae can be deduced from the preceding special case of  $a = e$  using the chain rule (Section 4.3).  $\square$

*Remark.* 1. Instead of the definition given in Section 2.5, some books use  $\lim_{y \rightarrow 0} (1 + y)^{\frac{1}{y}}$  as the definition of  $e$ .

2. The formula for  $(e^x)'$  and the formula for  $(\ln x)'$  imply each other, as  $e^x$  and  $\ln x$  are “inverse functions” of each other. (Cf. Chapter 5.)

**Example 4.2.2.**

$$1. (\sqrt{x} + 2^x - 3 \log_2 x)' = (\sqrt{x})' + (2^x)' - 3(\log_2 x)' = \frac{1}{2}x^{-\frac{1}{2}} + 2^x \ln 2 - \frac{3}{x \ln 2}$$

$$2. \frac{d}{dx}(x^2 e^x) = \frac{d}{dx}(x^2) \cdot e^x + x^2 \cdot \frac{d}{dx}(e^x) = (2x + x^2)e^x$$

$$3. \left(\frac{\sqrt{x}}{3^x}\right)' = ?$$

$$\text{by the quotient rule: } \frac{(\sqrt{x})'3^x - \sqrt{x}(3^x)'}{(3^x)^2} = \frac{\frac{1}{2}x^{-\frac{1}{2}} \cdot 3^x - x^{\frac{1}{2}} \cdot 3^x \ln 3}{(3^x)^2} = \frac{\frac{1}{2}x^{-\frac{1}{2}} - x^{\frac{1}{2}} \ln 3}{3^x}$$

$$\text{or, by Leibniz's rule: } \left(\sqrt{x} \cdot \left(\frac{1}{3}\right)^x\right)' = \frac{1}{2}x^{-\frac{1}{2}} \left(\frac{1}{3}\right)^x + x^{\frac{1}{2}} \left(\frac{1}{3}\right)^x \ln \frac{1}{3} = \frac{\frac{1}{2}x^{-\frac{1}{2}} - x^{\frac{1}{2}} \ln 3}{3^x}.$$

**Exercise 4.2.3.** Use two different methods to compute  $\left(\frac{1-x^2}{\sqrt{x}}\right)'$ .

**Example 4.2.3.** Suppose  $f(x)$  and  $g(x)$  are differentiable. Given  $f(1) = 1$ ,  $f'(1) = 2$ ,  $g(1) = 3$ ,  $g'(1) = 4$ . Find the value of

$$\frac{d}{dx}(f(x)g(x))$$

at  $x = 1$ .

*Solution.* By the product rule

$$\frac{d}{dx}(f(x)g(x)) = f'(x)g(x) + f(x)g'(x).$$

At  $x = 1$ , the above is

$$f'(1)g(1) + f(1)g'(1) = 2 \times 3 + 1 \times 4 = 10.$$

■

**Example 4.2.4.** Suppose  $f(x)$ ,  $g(x)$ ,  $h(x)$  are differentiable. Compute

$$\frac{d}{dx}(f(x)g(x)h(x)).$$

*Solution.*

$$\begin{aligned} \frac{d}{dx}(f(x)g(x)h(x)) &= (f(x)g(x)) \frac{d}{dx}h(x) + h(x) \frac{d}{dx}(f(x)g(x)) \\ &= f(x)g(x)h'(x) + h(x)\left(f(x) \frac{d}{dx}g(x) + g(x) \frac{d}{dx}f(x)\right) \\ &= f(x)g(x)h'(x) + f(x)g'(x)h(x) + f'(x)g(x)h(x). \end{aligned}$$

■

### 4.3 The Chain Rule (for composite functions)

**Theorem 6** (The Chain Rule).

If  $y = f(u)$  is a differentiable function of  $u$ ,

$u = g(x)$  is a differentiable function of  $x$ ,

then the composite function  $y = f(g(x))$  is a differentiable function of  $x$ , and

$$\boxed{\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}}$$

or equivalently

$$\boxed{\frac{dy}{dx} = f'(g(x))g'(x)}.$$

**A heuristic explanation:** Rewrite the difference quotient as a product:  $\frac{\Delta y}{\Delta x} = \frac{\Delta y}{\Delta u} \cdot \frac{\Delta u}{\Delta x}$ , then take  $\Delta x \rightarrow 0$ . (The notation “ $dx$ ” is conventionally used to represent an “infinitesimal  $\Delta x$ ”.)

**Example 4.3.1.** Find

$$\frac{d}{dx}(1 + 2x)^5.$$

*Solution.* Set  $y = f(u) = u^5$  and  $u = g(x) = 1 + 2x$ . Then  $f(g(x)) = (1 + 2x)^5$ .

By the chain rule,

$$f'(u) = \frac{dy}{du} = 5u^4 \quad \text{and} \quad g'(x) = \frac{du}{dx} = 2.$$

Hence,

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = (5u^4)(2) = 10(1 + 2x)^4,$$

or, alternatively written:

$$\frac{dy}{dx} = f'(g(x))g'(x) = 10(1 + 2x)^4.$$

■

**Example 4.3.2.** Find

$$\frac{d}{dx} \sqrt{1 + \sqrt{x}}.$$

*Solution.* Let  $y = f(u) = \sqrt{u}$ ,  $u = g(x) = 1 + \sqrt{x}$ . Then  $f(g(x)) = \sqrt{1 + \sqrt{x}}$ .

$$\frac{dy}{du} = \frac{1}{2} u^{-1/2} = \frac{1}{2\sqrt{u}} \quad \text{and} \quad \frac{du}{dx} = \frac{1}{2\sqrt{x}}.$$

Therefore

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{1}{2\sqrt{u}} \frac{1}{2\sqrt{x}} = \frac{1}{4\sqrt{x}\sqrt{1 + \sqrt{x}}}.$$

■

**Example 4.3.3.** Using  $(e^x)' = e^x$  and the chain rule, one may prove that  $(a^x)' = a^x \ln a$  ( $a > 0$ ).

*Proof.* Note that

$$\boxed{a^x = e^{\ln a^x}} \quad \text{(Very useful technique!)}$$

Then,

$$\begin{aligned} (a^x)' &= (e^{\ln a^x})' \\ &= (e^{x \ln a})' \\ &= e^{x \ln a} \cdot \ln a \\ &= a^x \cdot \ln a. \end{aligned}$$

□

**Example 4.3.4.** Use the Leibniz rule and the chain rule to prove the quotient rule.

*Proof.* By the Leibniz rule, we have

$$\left(\frac{f}{g}\right)' = \left(f \cdot \frac{1}{g}\right)' = f' \cdot \frac{1}{g} + f \cdot \left(\frac{1}{g}\right)'.$$

For  $\left(\frac{1}{g}\right)'$ , let  $y = \frac{1}{u}$ , where  $u = g(x)$ . Then, by the chain rule,

$$\left(\frac{1}{g}\right)' = \frac{dy}{du} \cdot \frac{du}{dx} = -\frac{1}{g^2(x)} g'(x).$$

Therefore,

$$\left(\frac{f}{g}\right)' = f' \frac{1}{g} - f \frac{g'}{g^2} = \frac{f'g - fg'}{g^2}.$$

□

**Example 4.3.5.** Find

$$\frac{d}{dx} e^{\sqrt{x^2+x}}.$$

*Solution.*

$$\begin{aligned} \frac{dy}{dx} &= e^{\sqrt{x^2+x}} \cdot (\sqrt{x^2+x})' && \text{(using the chain rule; write } y = e^u, u = \sqrt{x^2+x} \text{)} \\ &= e^{\sqrt{x^2+x}} \cdot \frac{1}{2}(x^2+x)^{-\frac{1}{2}} \cdot (2x+1) && \text{(using the chain rule again: let } u = \sqrt{w}, w = x^2+x \text{)} \end{aligned}$$

■

**Exercise 4.3.1.** Prove that

1.

$$\frac{d}{dx}(g(x))^n = n(g(x))^{n-1}g'(x).$$

2.

$$\frac{d}{dx} e^{\sqrt{\frac{x-1}{x+1}}} = e^{\sqrt{\frac{x-1}{x+1}}} \cdot (x-1)^{-\frac{1}{2}} \cdot (x+1)^{-\frac{3}{2}}.$$

### 4.3.1 Some tricks involving the log function and its derivative

**Example 4.3.6.** Show that

$$\boxed{\frac{d}{dx} \ln |x| = \frac{1}{x}, \quad x \neq 0.}$$

*Proof.* Let

$$y = \ln |x| = \begin{cases} \ln x, & \text{if } x > 0 \\ \ln(-x), & \text{if } x < 0 \end{cases}$$

For  $x > 0$ ,  $\frac{dy}{dx} = \frac{1}{x}$ ;

For  $x < 0$ ,  $\frac{dy}{dx} = \frac{1}{-x} \cdot (-1) = \frac{1}{x}$ . (by the chain rule)

□

**Example 4.3.7.** Let  $y = \sqrt[3]{\frac{(x-2)(x-3)^2}{x-5}}$ . Find  $\frac{dy}{dx}$ .

*Solution.*

$$\begin{aligned} y^3 &= \frac{(x-2)(x-3)^2}{x-5} \\ \ln y^3 &= \ln \frac{(x-2)(x-3)^2}{x-5} \\ 3 \ln y &= \ln(x-2) + 2 \ln(x-3) - \ln(x-5) \\ \frac{3}{y} \frac{dy}{dx} &= \frac{1}{x-2} + \frac{2}{x-3} - \frac{1}{x-5} \\ \frac{dy}{dx} &= \frac{y}{3} \left( \frac{1}{x-2} + \frac{2}{x-3} - \frac{1}{x-5} \right) \\ \frac{dy}{dx} &= \frac{1}{3} \sqrt[3]{\frac{(x-2)(x-3)^2}{x-5}} \left( \frac{1}{x-2} + \frac{2}{x-3} - \frac{1}{x-5} \right) \end{aligned}$$

■

*Remark.* Alternatively, one may regard  $y$  as a function of  $x$  defined “implicitly” via the relation  $(x-5)y^3 = (x-2)(x-3)^2$ . (Cf. Chapter 5.)

**Example 4.3.8.** Compute the derivative of  $x^x$ ,  $x > 0$ .

*Solution.* Write  $x^x = e^{x \ln x}$ . Let  $y = e^u$ , where  $u = x \ln x$ . Then

$$\begin{aligned}\frac{d}{dx}x^x &= \frac{dy}{du} \frac{du}{dx} \\ &= e^u \left( \ln x \frac{dx}{dx} + x \frac{d \ln x}{dx} \right) \\ &= e^u \left( \ln x + x \frac{1}{x} \right) \\ &= x^x (\ln x + 1).\end{aligned}$$

■

**Exercise 4.3.2.** Let  $y = f(x)^{g(x)}$ . Prove that  $y' = f(x)^{g(x)} \left( g'(x) \ln f(x) + \frac{f'(x)}{f(x)} g(x) \right)$ .