

MMAT 5340 Assignment #10
Please submit your assignment online on Blackboard
Due at 12 p.m. on Wednesday, Dec. 8, 2021

1. Consider a Markov chain $X = (X_n)_{n \geq 0}$ with a state space $S = \{1, 2, 3\}$ and the transition matrix

$$P = \begin{bmatrix} 0.9 & 0.1 & 0 \\ 0.1 & 0.8 & 0.1 \\ 0 & 0.1 & 0.9 \end{bmatrix}.$$

- (a) Show that the Markov chain is irreducible and recurrent.
 (b) Find the stationary distribution π such that $\pi^\top P = \pi^\top$.
 (c) Let $f(x) = x$, compute

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n f(X_k)}{n}.$$

2. Consider a Markov chain $X = (X_n)_{n \geq 0}$ with a state space $S = \mathbb{N}_0 = \{0, 1, 2, \dots\}$ and

$$P(x, x+1) = p, P(x, 0) = 1-p$$

for some $p \in (0, 1)$

- (a) Find the transition matrix P of X .
 (b) Is this Markov chain irreducible or reducible?

Recall that $\tau_x^1 := \inf\{n \geq 1 : X_n = x\}$.

- (c) Show that $P_x[\tau_0^1 = n] := P[\tau_0^1 = n | X_0 = x] = p^{n-1}(1-p)$. Compute $\mathbb{E}_x[\tau_0^1]$.
 (d) Prove that for $x \geq 2$,

$$\mathbb{E}_0[\tau_x^1] = \mathbb{E}_0[\tau_{x-1}^1] + \mathbb{E}_{x-1}[\tau_x^1]$$

and

$$\mathbb{E}_{x-1}[\tau_x^1] = 1 + (1-p)\mathbb{E}_0[\tau_x^1].$$

Hint: For the first equality, first show that for $\forall m, n \geq 1$,

$$\mathbb{P}_0[\tau_{x-1}^1 = m, \tau_x^1 - \tau_{x-1}^1 = n] = \mathbb{P}_0[\tau_{x-1}^1 = m] \cdot \mathbb{P}_{x-1}[\tau_x^1 = n].$$

Then use the fact that

$$\mathbb{P}_0[\tau_x^1 - \tau_{x-1}^1 = n] = \sum_{m=1}^{\infty} \mathbb{P}_0[\tau_{x-1}^1 = m] \cdot \mathbb{P}_{x-1}[\tau_x^1 = n].$$

For the second equality, you may accept that

$$\mathbb{P}_{x-1}[\tau_x^1 = n] = p \cdot \mathbf{1}_{\{n=1\}} + (1-p) \cdot \mathbf{1}_{\{n>1\}} \cdot \mathbb{P}_0[\tau_x^1 = n-1].$$

- (e) Optional. Define $f(x) := \mathbb{E}_0[\tau_x^1]$, $g(x) := \mathbb{E}_x[\tau_x^1]$. Deduce that

$$f(x) = \left(\frac{1}{p}\right)^x \frac{1}{1-p} - \frac{1}{1-p} \text{ for } x \geq 1,$$

and

$$g(x) = \left(\frac{1}{p}\right)^x \frac{1}{1-p}.$$

Hint: Compute $f(1)$. Define $u(x) := f(x) + \frac{1}{1-p}$, then use the result from (d) to deduce $u(x) = \frac{u(x-1)}{p}$. Note that $g(x) = \mathbb{E}_0[\tau_x^1] + \mathbb{E}_x[\tau_0^1]$.

(f) Check that

2

$$\pi(x) = \frac{1}{g(x)} = \frac{1}{\mathbb{E}_x[\tau_x^1]}$$

is a stationary distribution. Find the stationary probability for $x \in S$.