

1. Recall the definition for the notions of *orthonormal set* and *orthonormal basis* from the handout *Orthonormal basis and orthogonal projections*.

Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in \mathbb{R}^n$.

(a) We say that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute an orthonormal set in \mathbb{R}^n if and only if

$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ are pairwise orthogonal and $\|\mathbf{u}_j\| = 1$ for each $j = 1, 2, \dots, k$.

(b) Suppose V is a subspace of \mathbb{R}^n .

Then we say $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute an orthonormal basis for V if and only if

$\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute a basis for V and also constitute an orthonormal set.

Also recall the result (\star), which is a part of Theorem (C), as stated below:

(\star) Let W be a subspace of \mathbb{R}^n .


Suppose $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute an orthonormal basis for W .

Suppose $\mathbf{z} \in \mathbb{R}^n$.

Define $\mathbf{v} \in W$ by $\mathbf{v} = \langle \mathbf{z}, \mathbf{u}_1 \rangle \mathbf{u}_1 + \langle \mathbf{z}, \mathbf{u}_2 \rangle \mathbf{u}_2 + \dots + \langle \mathbf{z}, \mathbf{u}_k \rangle \mathbf{u}_k$.

Define $\mathbf{y} \in \mathbb{R}^n$ by $\mathbf{y} = \mathbf{z} - \mathbf{v}$.

Then $\mathbf{z} = \mathbf{v} + \mathbf{y}$, and $\mathbf{y} \perp \mathbf{s}$ for any $\mathbf{s} \in W$.



2. Lemma (G).

Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{z}$ be vectors in \mathbb{R}^n .

Suppose $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute an orthonormal set in \mathbb{R}^n .

Further suppose \mathbf{z} is not a linear combination of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$.

Define


$$\mathbf{y} = \mathbf{z} - \langle \mathbf{z}, \mathbf{u}_1 \rangle \mathbf{u}_1 - \langle \mathbf{z}, \mathbf{u}_2 \rangle \mathbf{u}_2 - \dots - \langle \mathbf{z}, \mathbf{u}_k \rangle \mathbf{u}_k.$$

Then the statements below hold:

(a) $\|\mathbf{y}\| \neq 0$.

(b) $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \frac{1}{\|\mathbf{y}\|}\mathbf{y}$ constitute an orthonormal set in \mathbb{R}^n .

(c) $\text{Span} (\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{z}\}) = \text{Span} \left(\left\{ \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \frac{1}{\|\mathbf{y}\|}\mathbf{y} \right\} \right)$.



Assumed to not belong to
 $W = \text{Span}(\{u_1, u_2, \dots, u_k\})$.
 (So $z - v \neq 0$.)

$W = \text{Span}(\{u_1, u_2, \dots, u_k\})$, with only
 u_1, u_2
 being
 displayed.

$$v = \langle z, u_1 \rangle u_1 + \langle z, u_2 \rangle u_2 + \dots + \langle z, u_k \rangle u_k$$

Some vector of
 norm 1 which is
 orthogonal to each
 of u_1, u_2, \dots, u_k
 and hence every vector in W .

3. Proof of Lemma (G).

Let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{z}$ be vectors in \mathbb{R}^n .

Suppose $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute an orthonormal set in \mathbb{R}^n .

Further suppose \mathbf{z} is not a linear combination of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$.

Define

$$W = \text{Span} (\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}).$$

By definition, $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitutes an orthonormal basis for W .

Define

$$\mathbf{y} = \mathbf{z} - \langle \mathbf{z}, \mathbf{u}_1 \rangle \mathbf{u}_1 - \langle \mathbf{z}, \mathbf{u}_2 \rangle \mathbf{u}_2 - \dots - \langle \mathbf{z}, \mathbf{u}_k \rangle \mathbf{u}_k.$$

Define

$$\mathbf{v} = \langle \mathbf{z}, \mathbf{u}_1 \rangle \mathbf{u}_1 + \langle \mathbf{z}, \mathbf{u}_2 \rangle \mathbf{u}_2 + \dots + \langle \mathbf{z}, \mathbf{u}_k \rangle \mathbf{u}_k.$$

Then $\mathbf{y} = \mathbf{z} - \mathbf{v}$ by definition.

(a) Since \mathbf{z} is not a linear combination of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$, we have $\mathbf{z} \neq \mathbf{v}$.

Then $\mathbf{y} = \mathbf{z} - \mathbf{v} \neq \mathbf{0}$. Therefore $\|\mathbf{y}\| \neq 0$.

(b) By the result (\star), $\mathbf{y} \perp \mathbf{s}$ for any $s \in W$.

Note that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in W$.

Then for each $j = 1, 2, \dots, n$, we have $\left\langle \frac{1}{\|\mathbf{y}\|} \mathbf{y}, \mathbf{u}_j \right\rangle = \frac{1}{\|\mathbf{y}\|} \langle \mathbf{y}, \mathbf{u}_j \rangle = 0$.

Hence $\frac{1}{\|\mathbf{y}\|} \mathbf{y} \perp \mathbf{u}_j$.

Also note that $\left\| \frac{1}{\|\mathbf{y}\|} \mathbf{y} \right\| = \frac{1}{\|\mathbf{y}\|} \cdot \|\mathbf{y}\| = 1$.

It follows that $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \frac{1}{\|\mathbf{y}\|} \mathbf{y}$ constitute an orthonormal set in \mathbb{R}^n .

(c) By definition, we have $\frac{1}{\|\mathbf{y}\|} \mathbf{y} = \frac{1}{\|\mathbf{y}\|} \mathbf{z} - \frac{\langle \mathbf{z}, \mathbf{u}_1 \rangle}{\|\mathbf{y}\|} \mathbf{u}_1 - \frac{\langle \mathbf{z}, \mathbf{u}_2 \rangle}{\|\mathbf{y}\|} \mathbf{u}_2 - \dots - \frac{\langle \mathbf{z}, \mathbf{u}_k \rangle}{\|\mathbf{y}\|} \mathbf{u}_k$.

Then each of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \frac{1}{\|\mathbf{y}\|} \mathbf{y}$ is a linear combination of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{z}$.

We also have $\mathbf{z} = \|\mathbf{y}\| \cdot \left(\frac{1}{\|\mathbf{y}\|} \mathbf{y} \right) + \langle \mathbf{z}, \mathbf{u}_1 \rangle \mathbf{u}_1 + \langle \mathbf{z}, \mathbf{u}_2 \rangle \mathbf{u}_2 + \dots + \langle \mathbf{z}, \mathbf{u}_k \rangle \mathbf{u}_k$.

Then each of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{z}$ is a linear combination of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \frac{1}{\|\mathbf{y}\|} \mathbf{y}$.

It follows that $\text{Span} (\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \mathbf{z}\}) = \text{Span} \left(\left\{ \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k, \frac{1}{\|\mathbf{y}\|} \mathbf{y} \right\} \right)$.

4. **Theorem (H). (Existence of orthonormal basis.)**

Suppose W is a non-zero subspace of \mathbb{R}^n . Then W has an orthonormal basis.

Remark.

The constructive argument in the proof below, generating an orthonormal basis for W from an (arbitrary) basis for W , is referred to as the Gram-Schmidt orthogonalization process.

5. **Proof of Theorem (H).**

Suppose W is a non-zero subspace of \mathbb{R}^n . Write $\dim(W) = k$.

By assumption, k is between 1 and n .

Pick some basis for W , which is a collection of k vectors, denoted by $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_k$.

For each $j = 1, 2, \dots, k$, define

$$W_j = \text{Span} (\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_j\}).$$

Note that $\dim(W_j) = j$, and by definition, \mathbf{z}_{j+1} does not belong to W_j .

[So now we have a sequence of subspaces of W , namely,

$$W_1, W_2, W_3, \dots, W_{k-2}, W_{k-1}, W_k,$$

in which $W_{\ell-1}$ is a ‘proper’ subspace of W_ℓ , in view of $\mathbf{z}_\ell \notin W_{\ell-1}$ and $\mathbf{z}_\ell \in W_\ell$.]

(a) Note that $\mathbf{z}_1 \neq \mathbf{0}$. Then $\|\mathbf{z}_1\| \neq 0$. (Take $\mathbf{y}_1 = \mathbf{z}_1$.)

Define $\mathbf{u}_1 = \frac{1}{\|\mathbf{z}_1\|}\mathbf{z}_1$. We have $\|\mathbf{u}_1\| = 1$.

\mathbf{u}_1 and \mathbf{z}_1 are non-zero scalar multiples of each other.

Then $W_1 = \text{Span}(\{\mathbf{z}_1\}) = \text{Span}(\{\mathbf{u}_1\})$.

Therefore \mathbf{u}_1 constitutes an orthonormal basis for W_1 .

(b) \mathbf{z}_2 does not belong to W_1 . Then \mathbf{z}_2 is not a linear combination of \mathbf{u}_1 .

Define $\mathbf{y}_2 = \mathbf{z}_2 - \langle \mathbf{z}_2, \mathbf{u}_1 \rangle \mathbf{u}_1$. By Lemma (G), $\|\mathbf{y}_2\| \neq 0$.

Define $\mathbf{u}_2 = \frac{1}{\|\mathbf{y}_2\|}\mathbf{y}_2$. By Lemma (G), $\mathbf{u}_1, \mathbf{u}_2$ constitute an orthonormal set in \mathbb{R}^n .

Since $\text{Span}(\{\mathbf{z}_1\}) = \text{Span}(\{\mathbf{u}_1\})$, we have $W_2 = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{z}_2\})$.


Then, again by Lemma (G),

$$W_2 = \text{Span}(\{\mathbf{u}_1, \mathbf{z}_2\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\}).$$

Therefore $\mathbf{u}_1, \mathbf{u}_2$ constitutes an orthonormal basis for W_2 .

$$\begin{aligned}
 W_2 &= \text{Span}(\{z_1, z_2\}) \\
 &= \text{Span}(\{u_1, z_2\}) \\
 &= \text{Span}(\{u_1, u_2\})
 \end{aligned}$$

$$y_2 = z_2 - \langle z_2, u_1 \rangle u_1$$



$$\begin{aligned}
 W &= \text{Span}(\{z_1\}) \\
 &= \text{Span}(\{u_1\})
 \end{aligned}$$

$$u_1 = \frac{z_1}{\|z_1\|}$$

$$u_2 = \frac{y_2}{\|y_2\|}$$

(c) \mathbf{z}_3 does not belong to W_2 . Then \mathbf{z}_3 is not a linear combination of $\mathbf{u}_1, \mathbf{u}_2$.

Define $\mathbf{y}_3 = \mathbf{z}_3 - \langle \mathbf{z}_3, \mathbf{u}_1 \rangle \mathbf{u}_1 - \langle \mathbf{z}_3, \mathbf{u}_2 \rangle \mathbf{u}_2$. By Lemma (G), $\|\mathbf{y}_3\| \neq 0$.

Define $\mathbf{u}_3 = \frac{1}{\|\mathbf{y}_3\|} \mathbf{y}_3$.

By Lemma (G), $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ constitute an orthonormal set in \mathbb{R}^n .


Since $\text{Span}(\{\mathbf{z}_1, \mathbf{z}_2\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$, we have

$$W_3 = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{z}_3\}).$$



Then, again by Lemma (G),

$$W_3 = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{z}_3\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}).$$

Therefore $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ constitutes an orthonormal basis for W_3 .



Done already



$W_3 = \text{Span}(\{z_1, z_2, z_3\})$
 $= \text{Span}(\{u_1, u_2, z_3\})$
 $= \text{Span}(\{u_1, u_2, u_3\})$

(d) Let ℓ be any one of $2, 3, \dots, k$.

Suppose that the vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_{\ell-1}$ and $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}$ are successively defined



by $\mathbf{y}_1 = \mathbf{z}_1$, $\mathbf{u}_1 = \frac{1}{\|\mathbf{z}_1\|} \mathbf{z}_1$ and

$$\left\{ \begin{array}{l} \mathbf{y}_2 = \mathbf{z}_2 - \langle \mathbf{z}_2, \mathbf{u}_1 \rangle \mathbf{u}_1, \\ \mathbf{u}_2 = \frac{1}{\|\mathbf{y}_2\|} \mathbf{y}_2, \\ \mathbf{y}_3 = \mathbf{z}_3 - \langle \mathbf{z}_3, \mathbf{u}_1 \rangle \mathbf{u}_1 - \langle \mathbf{z}_3, \mathbf{u}_2 \rangle \mathbf{u}_2, \\ \mathbf{u}_3 = \frac{1}{\|\mathbf{y}_3\|} \mathbf{y}_3, \\ \vdots \\ \mathbf{y}_{\ell-1} = \mathbf{z}_{\ell-1} - \langle \mathbf{z}_{\ell-1}, \mathbf{u}_1 \rangle \mathbf{u}_1 - \langle \mathbf{z}_{\ell-1}, \mathbf{u}_2 \rangle \mathbf{u}_2 - \dots - \langle \mathbf{z}_{\ell-1}, \mathbf{u}_{\ell-2} \rangle \mathbf{u}_{\ell-2}, \\ \mathbf{u}_{\ell-1} = \frac{1}{\|\mathbf{y}_{\ell-1}\|} \mathbf{y}_{\ell-1}, \end{array} \right.$$


and satisfies:

- $\|\mathbf{y}_2\| \neq 0$, $\|\mathbf{y}_3\| \neq 0$, ..., $\|\mathbf{y}_{\ell-1}\| \neq 0$, and
- for each $j = 2, 3, \dots, \ell - 1$,

the vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_j$ constitute an orthonormal basis for W_j .



$W_{l-1} = \text{Span}(\{z_1, z_2, \dots, z_{l-1}\})$
 in which only z_1 and z_{l-1} are displayed.



$$W_l = \text{Span}(\{z_1, \dots, z_{l-1}, z_l\})$$

$$= \text{Span}(\{u_1, \dots, u_{l-1}, z_l\})$$

$$= \text{Span}(\{u_1, \dots, u_{l-1}, u_l\})$$

We now note that \mathbf{z}_ℓ does not belong to $W_{\ell-1}$.

Then \mathbf{z}_ℓ is not a linear combination of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}$.

Define

$$\mathbf{y}_\ell = \mathbf{z}_\ell - \langle \mathbf{z}_\ell, \mathbf{u}_1 \rangle \mathbf{u}_1 - \langle \mathbf{z}_\ell, \mathbf{u}_2 \rangle \mathbf{u}_2 - \dots - \langle \mathbf{z}_\ell, \mathbf{u}_{\ell-1} \rangle \mathbf{u}_{\ell-1} - \langle \mathbf{z}_\ell, \mathbf{u}_\ell \rangle \mathbf{u}_\ell.$$

By Lemma (G), $\|\mathbf{y}_\ell\| \neq 0$.

Define $\mathbf{u}_\ell = \frac{1}{\|\mathbf{y}_\ell\|} \mathbf{y}_\ell$.

By Lemma (G), $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}, \mathbf{u}_\ell$ constitute an orthonormal set in \mathbb{R}^n .

Since $\text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_{\ell-1}\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}\})$, we have

$$W_\ell = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_{\ell-1}, \mathbf{z}_\ell\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}, \mathbf{z}_\ell\}).$$

Then, again by Lemma (G),

$$W_\ell = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_{\ell-1}, \mathbf{z}_\ell\}) = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}, \mathbf{u}_\ell\}).$$

Therefore $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell-1}, \mathbf{u}_\ell$ constitutes an orthonormal basis for W_ℓ .

Hence W has an orthonormal basis, namely $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$.

6. Gram-Schmidt orthogonalization process.

Suppose W is a subspace of \mathbb{R}^n , and

$\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \dots, \mathbf{z}_k$ constitute a basis for W .

The argument in the proof of Theorem (H) provides an algorithm for

obtaining an orthonormal basis $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ for W ,

for which the equality

$$\text{Span} (\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_j\}) = \text{Span} (\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_j\})$$

holds for each $j = 1, 2, \dots, k$:

- **Step (1).** We define $\mathbf{y}_1 = \mathbf{z}_1$.
- **Step (2).** We define $\mathbf{y}_2, \mathbf{y}_3, \dots, \mathbf{y}_k$ inductively by

$$\mathbf{y}_j = \mathbf{z}_j - \frac{\langle \mathbf{z}_j, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_j, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 - \dots - \frac{\langle \mathbf{z}_j, \mathbf{y}_{j-1} \rangle}{\|\mathbf{y}_{j-1}\|^2} \mathbf{y}_{j-1} \quad \text{for each } j = 2, 3, \dots, k.$$

When written out explicitly, $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k$ are given recursively by:

$$\left\{ \begin{array}{l} \mathbf{y}_1 = \mathbf{z}_1 \\ \mathbf{y}_2 = \mathbf{z}_2 - \frac{\langle \mathbf{z}_2, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 \\ \mathbf{y}_3 = \mathbf{z}_3 - \frac{\langle \mathbf{z}_3, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_3, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 \\ \mathbf{y}_4 = \mathbf{z}_4 - \frac{\langle \mathbf{z}_4, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_4, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 - \frac{\langle \mathbf{z}_4, \mathbf{y}_3 \rangle}{\|\mathbf{y}_3\|^2} \mathbf{y}_3 \\ \vdots \\ \mathbf{y}_k = \mathbf{z}_k - \frac{\langle \mathbf{z}_k, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_k, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 - \frac{\langle \mathbf{z}_k, \mathbf{y}_3 \rangle}{\|\mathbf{y}_3\|^2} \mathbf{y}_3 - \dots - \frac{\langle \mathbf{z}_k, \mathbf{y}_{k-1} \rangle}{\|\mathbf{y}_{k-1}\|^2} \mathbf{y}_{k-1} \end{array} \right.$$

- **Step (3).** For each $j = 1, 2, \dots, k$, define $\mathbf{u}_j = \frac{1}{\|\mathbf{y}_j\|} \mathbf{y}_j$.

For each $\ell = 1, 2, \dots, k$,

the vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_\ell$ constitute an orthonormal basis for $\text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_\ell\})$.

In particular, the vectors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ constitute an orthonormal basis for W .

7. Illustrations on the Gram-Schmidt orthogonalization process.

(a) Let $\mathbf{z}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$, $\mathbf{z}_2 = \begin{bmatrix} -1 \\ 1 \\ 4 \end{bmatrix}$, $\mathbf{z}_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix}$.

Take for granted that $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3$ are linearly independent.

We proceed to find an orthonormal basis for $W = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3\}) = \mathbb{R}^3$.

• Take $\mathbf{y}_1 = \mathbf{z}_1$. Then $\mathbf{y}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$, and $\|\mathbf{y}_1\|^2 = 9$.

Take $\mathbf{u}_1 = \frac{1}{\|\mathbf{y}_1\|} \mathbf{y}_1$. Then $\mathbf{u}_1 = \begin{bmatrix} 1/3 \\ 2/3 \\ 2/3 \end{bmatrix}$.

- Take $\mathbf{y}_2 = \mathbf{z}_2 - \frac{\langle \mathbf{z}_2, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1$. We have $\langle \mathbf{z}_2, \mathbf{y}_1 \rangle = 9$.

$$\text{Then } \mathbf{y}_2 = \begin{bmatrix} -1 \\ 1 \\ 4 \end{bmatrix} - \frac{9}{9} \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} -2 \\ -1 \\ 2 \end{bmatrix}, \text{ and } \|\mathbf{y}_2\|^2 = 9.$$

$$\text{Take } \mathbf{u}_2 = \frac{1}{\|\mathbf{y}_2\|} \mathbf{y}_2. \text{ Then } \mathbf{u}_2 = \begin{bmatrix} -2/3 \\ -1/3 \\ 2/3 \end{bmatrix}.$$

- Take $\mathbf{y}_3 = \mathbf{z}_3 - \frac{\langle \mathbf{z}_3, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_3, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2$. We have $\langle \mathbf{z}_3, \mathbf{y}_1 \rangle = -3$, $\langle \mathbf{z}_3, \mathbf{y}_2 \rangle = 6$.

$$\text{Then } \mathbf{y}_3 = \begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix} - \frac{-3}{9} \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} - \frac{6}{9} \begin{bmatrix} -2 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2/3 \\ -2/3 \\ 1/3 \end{bmatrix}, \text{ and } \|\mathbf{y}_3\|^2 = 1.$$

$$\text{Take } \mathbf{u}_3 = \frac{1}{\|\mathbf{y}_3\|} \mathbf{y}_3. \text{ Then } \mathbf{u}_3 = \begin{bmatrix} 2/3 \\ -2/3 \\ 1/3 \end{bmatrix}.$$

$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ constitute an orthonormal basis for W .

Also note that, by construction, $\text{Span}(\{\mathbf{u}_1\}) = \text{Span}(\{\mathbf{z}_1\})$ and $\text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\}) = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2\})$.

(b) Let $\mathbf{z}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{z}_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$, $\mathbf{z}_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}$.

Take for granted that $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3$ are linearly independent.

We proceed to find an orthonormal basis for $W = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3\})$.

- Take $\mathbf{y}_1 = \mathbf{z}_1$.

Then $\mathbf{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$, and $\|\mathbf{y}_1\|^2 = 2$.

Take $\mathbf{u}_1 = \frac{1}{\|\mathbf{y}_1\|}\mathbf{y}_1$. Then $\mathbf{u}_1 = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$.

- Take $\mathbf{y}_2 = \mathbf{z}_2 - \frac{\langle \mathbf{z}_2, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1$.

We have $\langle \mathbf{z}_2, \mathbf{y}_1 \rangle = 2$.

$$\text{Then } \mathbf{y}_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{2}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}, \text{ and } \|\mathbf{y}_2\|^2 = 2.$$

$$\text{Take } \mathbf{u}_2 = \frac{1}{\|\mathbf{y}_2\|} \mathbf{y}_2. \text{ Then } \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{bmatrix}.$$

- Take $\mathbf{y}_3 = \mathbf{z}_3 - \frac{\langle \mathbf{z}_3, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_3, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2$.

We have $\langle \mathbf{z}_3, \mathbf{y}_1 \rangle = 1$, $\langle \mathbf{z}_3, \mathbf{y}_2 \rangle = 2$.

$$\text{Then } \mathbf{y}_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} - \frac{2}{2} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1/2 \\ 0 \\ 1/2 \\ 0 \end{bmatrix}, \text{ and } \|\mathbf{y}_3\|^2 = \frac{1}{2}.$$

$$\text{Take } \mathbf{u}_3 = \frac{1}{\|\mathbf{y}_3\|} \mathbf{y}_3. \text{ Then } \mathbf{u}_3 = \begin{bmatrix} -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \\ 0 \end{bmatrix}.$$

$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ constitute an orthonormal basis for W .

Also note that, by construction,

$$\text{Span} (\{\mathbf{u}_1\}) = \text{Span} (\{\mathbf{z}_1\}) \text{ and } \text{Span} (\{\mathbf{u}_1, \mathbf{u}_2\}) = \text{Span} (\{\mathbf{z}_1, \mathbf{z}_2\}).$$

(c) Let $\mathbf{z}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{z}_2 = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 1 \end{bmatrix}$, $\mathbf{z}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$.

Take for granted that $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3$ are linearly independent.

We proceed to find an orthonormal basis for $W = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3\})$.

- Take $\mathbf{y}_1 = \mathbf{z}_1$.

Then $\mathbf{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$, and $\|\mathbf{y}_1\|^2 = 2$.

Take $\mathbf{u}_1 = \frac{1}{\|\mathbf{y}_1\|}\mathbf{y}_1$. Then $\mathbf{u}_1 = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$.

- Take $\mathbf{y}_2 = \mathbf{z}_2 - \frac{\langle \mathbf{z}_2, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1$.

We have $\langle \mathbf{z}_2, \mathbf{y}_1 \rangle = 2$.

$$\text{Then } \mathbf{y}_2 = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 1 \end{bmatrix} - \frac{2}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \text{ and } \|\mathbf{y}_2\|^2 = 4.$$

$$\text{Take } \mathbf{u}_2 = \frac{1}{\|\mathbf{y}_2\|} \mathbf{y}_2. \text{ Then } \mathbf{u}_2 = \begin{bmatrix} -1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{bmatrix}.$$

- Take $\mathbf{y}_3 = \mathbf{z}_3 - \frac{\langle \mathbf{z}_3, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_3, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2$.

We have $\langle \mathbf{z}_3, \mathbf{y}_1 \rangle = 2$, $\langle \mathbf{z}_3, \mathbf{y}_2 \rangle = 2$.

$$\text{Then } \mathbf{y}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{2}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} - \frac{2}{4} \begin{bmatrix} -1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1/2 \\ 1/2 \\ -1/2 \\ 1/2 \end{bmatrix}, \text{ and } \|\mathbf{y}_3\|^2 = 1.$$

$$\text{Take } \mathbf{u}_3 = \frac{1}{\|\mathbf{y}_3\|} \mathbf{y}_3. \text{ Then } \mathbf{u}_3 = \begin{bmatrix} 1/2 \\ 1/2 \\ -1/2 \\ 1/2 \end{bmatrix}.$$

$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ constitute an orthonormal basis for W .

Also note that, by construction,

$$\text{Span} (\{\mathbf{u}_1\}) = \text{Span} (\{\mathbf{z}_1\}) \text{ and } \text{Span} (\{\mathbf{u}_1, \mathbf{u}_2\}) = \text{Span} (\{\mathbf{z}_1, \mathbf{z}_2\}).$$

$$(d) \text{ Let } \mathbf{z}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \\ 4 \end{bmatrix}, \mathbf{z}_2 = \begin{bmatrix} -2 \\ 6 \\ 2 \\ 9 \end{bmatrix}, \mathbf{z}_3 = \begin{bmatrix} 9 \\ -2 \\ -4 \\ 7 \end{bmatrix}, \mathbf{z}_4 = \begin{bmatrix} -3 \\ -1 \\ -3 \\ 9 \end{bmatrix}.$$

Take for granted that $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \mathbf{z}_4$ are linearly independent.

We proceed to find an orthonormal basis for $W = \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \mathbf{z}_4\}) = \mathbb{R}^4$.

- Take $\mathbf{y}_1 = \mathbf{z}_1$.

$$\text{Then } \mathbf{y}_1 = \begin{bmatrix} 1 \\ 2 \\ 2 \\ 4 \end{bmatrix}, \text{ and } \|\mathbf{y}_1\|^2 = 25.$$

$$\text{Take } \mathbf{u}_1 = \frac{1}{\|\mathbf{y}_1\|} \mathbf{y}_1. \text{ Then } \mathbf{u}_1 = \begin{bmatrix} 1/5 \\ 2/5 \\ 2/5 \\ 4/5 \end{bmatrix}.$$

- Take $\mathbf{y}_2 = \mathbf{z}_2 - \frac{\langle \mathbf{z}_2, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1$.

We have $\langle \mathbf{z}_2, \mathbf{y}_1 \rangle = 50$.

$$\text{Then } \mathbf{y}_2 = \begin{bmatrix} -2 \\ 6 \\ 2 \\ 9 \end{bmatrix} - \frac{50}{25} \begin{bmatrix} 1 \\ 2 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} -4 \\ 2 \\ -2 \\ 1 \end{bmatrix}, \text{ and } \|\mathbf{y}_2\|^2 = 25.$$

$$\text{Take } \mathbf{u}_2 = \frac{1}{\|\mathbf{y}_2\|} \mathbf{y}_2. \text{ Then } \mathbf{u}_2 = \begin{bmatrix} -4/5 \\ 2/5 \\ -2/5 \\ 1/5 \end{bmatrix}.$$

- Take $\mathbf{y}_3 = \mathbf{z}_3 - \frac{\langle \mathbf{z}_3, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_3, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2$.

We have $\langle \mathbf{z}_3, \mathbf{y}_1 \rangle = 25$, $\langle \mathbf{z}_3, \mathbf{y}_2 \rangle = -25$.

$$\text{Then } \mathbf{y}_3 = \begin{bmatrix} 9 \\ -2 \\ -4 \\ 7 \end{bmatrix} - \frac{25}{25} \begin{bmatrix} 1 \\ 2 \\ 2 \\ 4 \end{bmatrix} - \frac{-25}{25} \begin{bmatrix} -4 \\ 2 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -2 \\ -8 \\ 4 \end{bmatrix}, \text{ and } \|\mathbf{y}_3\|^2 = 100.$$

$$\text{Take } \mathbf{u}_3 = \frac{1}{\|\mathbf{y}_3\|} \mathbf{y}_3. \text{ Then } \mathbf{u}_3 = \begin{bmatrix} 2/5 \\ -1/5 \\ -4/5 \\ 2/5 \end{bmatrix}.$$

- Take $\mathbf{y}_4 = \mathbf{z}_4 - \frac{\langle \mathbf{z}_4, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_4, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 - \frac{\langle \mathbf{z}_4, \mathbf{y}_3 \rangle}{\|\mathbf{y}_3\|^2} \mathbf{y}_3$.

We have $\langle \mathbf{z}_4, \mathbf{y}_1 \rangle = 25$, $\langle \mathbf{z}_4, \mathbf{y}_2 \rangle = 25$, $\langle \mathbf{z}_4, \mathbf{y}_3 \rangle = 50$.

$$\text{Then } \mathbf{y}_4 = \begin{bmatrix} -3 \\ -1 \\ -3 \\ 9 \end{bmatrix} - \frac{25}{25} \begin{bmatrix} 1 \\ 2 \\ 2 \\ 4 \end{bmatrix} - \frac{25}{25} \begin{bmatrix} -4 \\ 2 \\ -2 \\ 1 \end{bmatrix} - \frac{50}{100} \begin{bmatrix} 4 \\ -2 \\ -8 \\ 4 \end{bmatrix} = \begin{bmatrix} -2 \\ -4 \\ 1 \\ 2 \end{bmatrix}, \text{ and } \|\mathbf{y}_4\|^2 = 25.$$

$$\text{Take } \mathbf{u}_4 = \frac{1}{\|\mathbf{y}_4\|} \mathbf{y}_4. \text{ Then } \mathbf{u}_4 = \begin{bmatrix} -2/5 \\ -4/5 \\ 1/5 \\ 2/5 \end{bmatrix}.$$

$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4$ constitute an orthonormal basis for W .

Also note that, by construction,

$$\begin{aligned} \text{Span}(\{\mathbf{u}_1\}) &= \text{Span}(\{\mathbf{z}_1\}), & \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\}) &= \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2\}) \\ \text{and } \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}) &= \text{Span}(\{\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3\}). \end{aligned}$$

8. Gram-Schmidt orthogonalization, presented as QR-decomposition.

Suppose Z is an $(n \times k)$ -matrix, with $n \geq k$. For each $j = 1, 2, \dots, k$, the j -th column of Z by \mathbf{z}_j for.

Suppose $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \dots, \mathbf{z}_k$ are linearly independent.

We define

$$\mathbf{y}_1 = \mathbf{z}_1, \quad \mathbf{u}_1 = \frac{1}{\|\mathbf{y}_1\|} \mathbf{y}_1,$$

and define

$$\mathbf{y}_2, \mathbf{y}_3, \dots, \mathbf{y}_k$$

inductively by

$$\begin{cases} \mathbf{y}_j = \mathbf{z}_j - \frac{\langle \mathbf{z}_j, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 - \frac{\langle \mathbf{z}_j, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 - \dots - \frac{\langle \mathbf{z}_j, \mathbf{y}_{j-1} \rangle}{\|\mathbf{y}_{j-1}\|^2} \mathbf{y}_{j-1}, \\ \mathbf{u}_j = \frac{1}{\|\mathbf{y}_j\|} \mathbf{y}_j \end{cases}$$

for each $j = 2, 3, \dots, k$.

According to the argument for Theorem (H), these vectors $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3, \dots, \mathbf{y}_k$ are well-defined

Define the $(n \times k)$ -matrix $Q = \left[\mathbf{u}_1 \mid \mathbf{u}_2 \mid \dots \mid \mathbf{u}_k \right]$.

For each $j = 1, 2, \dots, k$, we have

$$\begin{aligned} \mathbf{z}_j &= \frac{\langle \mathbf{z}_j, \mathbf{y}_1 \rangle}{\|\mathbf{y}_1\|^2} \mathbf{y}_1 + \frac{\langle \mathbf{z}_j, \mathbf{y}_2 \rangle}{\|\mathbf{y}_2\|^2} \mathbf{y}_2 + \dots + \frac{\langle \mathbf{z}_j, \mathbf{y}_{j-1} \rangle}{\|\mathbf{y}_{j-1}\|^2} \mathbf{y}_{j-1} + \mathbf{y}_j \\ &= \langle \mathbf{z}_j, \mathbf{u}_1 \rangle \mathbf{u}_1 + \langle \mathbf{z}_j, \mathbf{u}_2 \rangle \mathbf{u}_2 + \dots + \langle \mathbf{z}_j, \mathbf{u}_{j-1} \rangle \mathbf{u}_{j-1} + \|\mathbf{y}_j\| \cdot \mathbf{u}_j + 0 \cdot \mathbf{u}_{j+1} + \dots + 0 \cdot \mathbf{u}_k \\ &= Q \begin{bmatrix} \langle \mathbf{u}_1, \mathbf{z}_j \rangle \\ \langle \mathbf{u}_2, \mathbf{z}_j \rangle \\ \vdots \\ \langle \mathbf{u}_{j-1}, \mathbf{z}_j \rangle \\ \|\mathbf{y}_j\| \\ 0 \\ \vdots \\ 0 \end{bmatrix}. \end{aligned}$$

Define the $(n \times n)$ -square matrix R , whose (i, j) -th entry is denoted by r_{ij} and given by

$$r_{ij} = \begin{cases} \langle \mathbf{u}_i, \mathbf{z}_j \rangle & \text{if } i < j \\ \|\mathbf{y}_j\| & \text{if } i = j \\ 0 & \text{if } i > j \end{cases}$$

(So, for each $j = 1, 2, \dots, n$, the j -th column of R is $\begin{bmatrix} \langle \mathbf{u}_1, \mathbf{z}_j \rangle \\ \langle \mathbf{u}_2, \mathbf{z}_j \rangle \\ \vdots \\ \langle \mathbf{u}_{j-1}, \mathbf{z}_j \rangle \\ \|\mathbf{y}_j\| \\ 0 \\ \vdots \\ 0 \end{bmatrix}$.)

Then $Z = QR$.

This ‘factorization’ of Z into the product QR is called the ‘QR-decomposition’ for Z .

Note that $\mathcal{C}(Z) = \mathcal{C}(Q)$ and the columns of Q is an orthonormal basis for $\mathcal{C}(Z)$.

The matrix R encodes the Gram-Schmidt orthogonalization process from which we obtain the orthonormal set $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ from the linearly independent set $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_k$.

9. Illustrations of QR-decomposition.

Refer to *Illustrations on the Gram-Schmidt orthogonalization process* above.

The respective constructions can be displayed as the ‘factorizations’ below:

$$(a) \begin{bmatrix} 1 & -1 & -1 \\ 2 & 1 & -2 \\ 2 & 4 & 1 \end{bmatrix} = \begin{bmatrix} 1/3 & -2/3 & 2/3 \\ 2/3 & -1/3 & -2/3 \\ 2/3 & 2/3 & 1/3 \end{bmatrix} \begin{bmatrix} 3 & 3 & -1 \\ 0 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$(b) \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1/\sqrt{2} & 0 \\ 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1/\sqrt{2} & 0 \end{bmatrix} \begin{bmatrix} \sqrt{2} & \sqrt{2} & 1/\sqrt{2} \\ 0 & \sqrt{2} & \sqrt{2} \\ 0 & 0 & 1/\sqrt{2} \end{bmatrix}.$$

$$(c) \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 2 & 1 \\ 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} & -1/2 & -1/2 \\ 0 & 1/2 & 1/2 \\ 1/\sqrt{2} & 1/2 & -1/2 \\ 0 & 1/2 & 1/2 \end{bmatrix} \begin{bmatrix} \sqrt{2} & \sqrt{2} & \sqrt{2} \\ 0 & 2 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$(d) \begin{bmatrix} 1 & -2 & 9 & -3 \\ 2 & 6 & -2 & -1 \\ 2 & 2 & -4 & -3 \\ 4 & 9 & 7 & 9 \end{bmatrix} = \begin{bmatrix} 1/5 & -4/5 & 2/5 & -2/5 \\ 2/5 & 2/5 & -1/5 & -4/5 \\ 2/5 & -2/5 & -4/5 & 1/5 \\ 4/5 & 1/5 & 2/5 & 2/5 \end{bmatrix} \begin{bmatrix} 5 & 10 & 5 & 5 \\ 0 & 5 & -5 & 5 \\ 0 & 0 & 10 & 5 \\ 0 & 0 & 0 & 5 \end{bmatrix}.$$