

e.g. Prove that  $e^x \geq 1+x \quad \forall x \in \mathbb{R}$ .

(i.e.  $e^x - x - 1 \geq 0$ )

Let  $f(x) = e^x - x - 1$

(Want to find the global minimum of  $f(x)$  and see if it is  $\geq 0$ .)

$$\text{Let } f(x) = e^x - x - 1$$

(Want to find the global minimum of  $f(x)$  and see if it is  $\geq 0$ .)

$$f'(x) = e^x - 1$$

$f'(x) > 0$  if  $x > 0$  and  $f'(x) < 0$  if  $x < 0$

$f$  is strictly increasing when  $x > 0$  and strictly decreasing when  $x < 0$

(and  $f$  is continuous at  $x=0$ .)

$f$  attains minimum when  $x=0$  (By 1st derivative check)


(In fact, global minimum, why?)

$$\therefore f(x) \geq f(0) \quad \forall x \in \mathbb{R} \quad \text{--- (*)}$$

$$= e^0 - 0 - 1$$

$$= 0$$

Note: The equality holds iff  $x=0$



Ex: By considering  $f(x) = \sin x - x \cos x$  for  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ ,

show that  $\cos x < \frac{\sin x}{x}$  for  $-\frac{\pi}{2} < x < 0$  or  $0 < x < \frac{\pi}{2}$ .

## Stationary Points :

If  $f'(a) = 0$ , then  $(a, f(a))$  is called a stationary point.


But even  $f'(a) = 0$ , it's still hard to say !

e.g. If  $f(x) = x^3$ , then  $f'(x) = 3x^2$ .

Note: 1)  $f'(0) = 0$

2)  $f'(x) = 3x^2 > 0$  for  $x \neq 0$

i.e. No change of sign of  $f'(x)$  at  $x = 0$ .



Note: a stationary is NOT necessary to be a max./min. point !

Theorem: Let  $f: (a, b) \rightarrow \mathbb{R}$  be a function and  $c \in (a, b)$  such that

1)  $f'(c)$  exists

2)  $f$  attains maximum (or minimum) at  $x=c$ .

Then, we have  $f'(c) = 0$ .

proof: Assume  $f$  attains maximum at  $x=c$ .

$$f'(c) \text{ exist} \Rightarrow \lim_{\Delta x \rightarrow 0^+} \frac{f(c+\Delta x) - f(c)}{\Delta x} = \lim_{\Delta x \rightarrow 0^-} \frac{f(c+\Delta x) - f(c)}{\Delta x} = f'(c)$$

$$\text{Note: } \frac{f(c+\Delta x) - f(c)}{\Delta x} \leq 0 \text{ for all } \Delta x > 0 \Rightarrow f'(c) = \lim_{\Delta x \rightarrow 0^+} \frac{f(c+\Delta x) - f(c)}{\Delta x} \leq 0$$

$$\frac{f(c+\Delta x) - f(c)}{\Delta x} \geq 0 \text{ for all } \Delta x < 0 \Rightarrow f'(c) = \lim_{\Delta x \rightarrow 0^-} \frac{f(c+\Delta x) - f(c)}{\Delta x} \geq 0$$

$$\therefore f'(c) = 0.$$

Remark: If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a differentiable function,


solving the equation  $f'(x) = 0$  is sufficient to capture all maximum and minimum.

e.g. If  $f(x) = x^3 - 3x^2 - 9x + 5$


then  $f'(x) = 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) = 3(x-3)(x-1)$

$f'(x) > 0$  if  $x > 3$  or  $x < -1$

$f'(x) < 0$  if  $-1 < x < 3$



Furthermore,



## Higher Derivatives :

$s(t)$  : distance function (depends on time  $t$ )

(instantaneous) speed = rate of change of distance travelled  
with respect to  $t$ .

$$v(t) = \frac{ds}{dt} \quad (\text{still a function of } t)$$

Question : What is  $\frac{dv}{dt}$  ?


Answer : Acceleration !

= rate of change of speed with respect to  $t$ .




We write  $a(t) = \frac{dv}{dt} = \frac{d^2s}{dt^2}$


e.g.  $s(t) = t^2$



$v(t) = \frac{ds}{dt} = 2t$



$a(t) = \frac{dv}{dt} = \frac{d^2s}{dt^2} = 2$



speed is increasing  
i.e. accelerating

In general, let  $y = f(x)$

We have : (1st derivative)  $\frac{dy}{dx} = \frac{df}{dx} = f'(x)$

(2nd derivative)  $\frac{d^2y}{dx^2} = \frac{d^2f}{dx^2} = f''(x)$


(n<sup>th</sup> derivative)  $\frac{d^ny}{dx^n} = \frac{d^nf}{dx^n} = f^{(n)}(x)$

## 2nd Derivative and Concavity:

Think: If  $f''(x) > 0$  for  $a < x < b$

then  $f'(x)$  is strictly increasing on  $(a, b)$

Picture:




Slope of the tangent line at  $(x, f(x))$  increases as  $x$  increases!

(NOT  $f(x)$  is increasing!)

If  $f''(x) > 0$  for  $a < x < b$ ,

then  $f(x)$  is a **concave** function on  $(a, b)$ .

Picture :



Slope of the tangent line at  $(x, f(x))$  increases as  $x$  increases!  
(NOT  $f(x)$  is increasing!)

If  $f''(x) > 0$  for  $a < x < b$ ,

then  $f(x)$  is a **concave** function on  $(a, b)$ .

Similarly : If  $f''(x) < 0$  for  $a < x < b$ ,

then  $f(x)$  is a **convex** function on  $(a, b)$ .

## 2nd Derivative Check:

Suppose  $f(x)$  is twice differentiable at  $x=a$ . (i.e.  $f'(a)$  and  $f''(a)$  exist)

If (1)  $f'(a) = 0$  (i.e.  $(a, f(a))$  is a stationary point.)

(2)  $f''(a) < 0$  (Roughly speaking:  $f(x)$  is convex near  $x=a$ ,

if  $f''(x)$  is continuous at  $x=a$ .)

then  $(a, f(a))$  is a relative maximum.

We have similar result for relative minimum.

Caution: If  $f''(a) = 0$ , then NO conclusion!

Consider  $f(x) = x^4, x^3, -x^4$

We have  $f'(0) = f''(0) = 0$  in each case, but  $(0,0)$  is

- min. for the 1st case.
- saddle point for the 2nd case.
- max. for the 3rd case.

e.g. If  $f(x) = x^3 - 3x^2 - 9x + 5$

then  $f'(x) = 3x^2 - 6x - 9 = 3(x^2 - 2x - 3) = 3(x-3)(x+1)$

$$f'(x) > 0 \text{ if } x > 3 \text{ or } x < -1$$

$$f'(x) < 0 \text{ if } -1 < x < 3$$



$$f''(x) = 6x - 6$$

$$f''(x) > 0 \text{ if } x > 1$$

$$f''(-1) = 12 < 0$$

$$f''(x) < 0 \text{ if } x < 1$$

$$f''(3) = 12 > 0$$



Note: The curve changes from being convex to concave at  $(1, 6)$ .

This point is called a **point of inflection**.

Point of inflection:

Suppose  $f(x)$  is continuous at  $x=a$  and differentiable on some open interval  $I$  containing  $x=a$ , except possibly at  $x=a$  itself.

If  $f''(x) > 0$  (resp.  $f''(x) < 0$ ) for all  $x$  in  $I$  with  $x < a$ , and

$f''(x) < 0$  (resp.  $f''(x) > 0$ ) for all  $x$  in  $I$  with  $x > a$ ,

then  $(a, f(a))$  is a point of inflection.

(Remember the slogan: Change sign of  $f''(x)$  at  $x=a$ .)



e.g.  $f(x) = 12x^5 - 105x^4 + 340x^3 - 510x^2 + 360x - 120$

Find the range of  $x$  such that

(1)  $f'(x) > 0$  ,  $f'(x) < 0$

(2)  $f''(x) > 0$  ,  $f''(x) < 0$

Step 1 : Find  $f'(x)$  and factorize it.

$$f'(x) = 60x^4 - 420x^3 + 1020x^2 - 1020x + 360$$

$$= 60(x^4 - 7x^3 + 17x^2 - 17x + 6)$$

$$= 60(x-1)^2(x-2)(x-3)$$

(Using factor theorem)

Step 2:



↓ gives intervals

$$x < 1 \quad 1 < x < 2 \quad 2 < x < 3 \quad x > 3$$

(Reason: those factors may change sign at the boundaries of the intervals.)

Step 3:

	$x < 1$	$x = 1$	$1 < x < 2$	$x = 2$	$2 < x < 3$	$x = 3$	$x > 3$
$(x-1)^2$	+	0	+	+	+	+	+
$(x-2)$	-	-	-	0	+	+	+
$(x-3)$	-	-	-	-	-	0	+
<hr/>							
$f'(x)$	+	0	+	0	-	0	+

$f(x)$       inc      saddle pt.      inc.      max.      dec.      min      inc.


saddle point =  $(1, -23)$

max =  $(2, -16)$

min =  $(3, -39)$

Similarly,

$$\begin{aligned} f''(x) &= 240x^3 - 1260x^2 + 2040x - 1020 \\ &= 60(x-1)(4x^2 - 17x + 17) \\ &= 240(x-1) \left[ x - \left( \frac{17+\sqrt{17}}{8} \right) \right] \left[ x - \left( \frac{17-\sqrt{17}}{8} \right) \right] \end{aligned}$$



points of inflection :  $(1, -23)$ ,  $(\frac{17 \pm \sqrt{17}}{8}, f(\frac{17 \pm \sqrt{17}}{8}))$

$$y = f(x) = 12x^5 - 105x^4 + 340x^3 - 510x^2 + 360x - 120$$

(1.00, -23.0)




It is a saddle  
point as well as  
a point of inflection

(1.61, -19.1)

(2.00, -16.0)


(2.64, -29.7)

(3.00, -39.0)





eg.  $f(x) = \frac{x}{(x+1)^2} \quad x \neq -1$

$$f'(x) = \frac{1-x}{(x+1)^3}$$





max. =  $(1, \frac{1}{4})$

$$f''(x) = \frac{2(x-2)}{(x+1)^4}$$



point of inflection :  $(2, \frac{2}{9})$





Note : The graph of  $y=f(x)$  behaves like

- the vertical line  $x=-1$ , when  $x$  is "near"  $-1$ .
- the horizontal line  $y=0$ , when  $x$  is "near"  $+\infty$  or  $-\infty$ .

In fact,  $x=-1$  is called a vertical asymptote,

$y=0$  is called a horizontal asymptote.



Finding vertical asymptote:

If  $\lim_{x \rightarrow a^+} f(x)$  or  $\lim_{x \rightarrow a^-} f(x) = +\infty$  or  $-\infty$ , then  $x=a$  is called a vertical asymptote.

Finding horizontal asymptote:

If  $\lim_{x \rightarrow +\infty} f(x) = L$ , where  $L$  is a real number, then  $y=L$  is a horizontal asymptote.

(Similar for  $\lim_{x \rightarrow -\infty} f(x)$ )

Note: It may happen that both  $\lim_{x \rightarrow +\infty} f(x)$  and  $\lim_{x \rightarrow -\infty} f(x)$  exist but they are NOT the same.