

THE CHINESE UNIVERSITY OF HONG KONG
Department of Mathematics
MATH 2050B Mathematical Analysis I
Tutorial 3 (September 25, 27)

The following were discussed in the tutorial this week:

1 More on Limit of a Sequence

Example 1. Let (y_n) be a sequence of positive numbers such that $\lim_n y_n = 2$. By virtue of ε - N terminology, show that

$$\lim_n \frac{y_n}{y_n^2 - 3} = 2.$$

Solution. Let $\varepsilon > 0$ be given. For $n \in \mathbb{N}$,

$$\begin{aligned} \left| \frac{y_n}{y_n^2 - 3} - 2 \right| &= \left| \frac{y_n - 2y_n^2 + 6}{y_n^2 - 3} \right| = \left| \frac{(2y_n + 3)(y_n - 2)}{y_n^2 - 3} \right| \\ &= \frac{|2y_n + 3|}{|y_n^2 - 3|} \cdot |y_n - 2|. \end{aligned}$$

Want: a positive lower bound of $|y_n^2 - 3|$ when n is large.

To archive this, we choose a neighbourhood of 2 that avoids the zeros of $y^2 - 3$, that is $\pm\sqrt{3} \approx \pm 1.73$. For example, $(2 - \frac{1}{4}, 2 + \frac{1}{4})$.

If $|y_n - 2| < \frac{1}{4}$, then

$$\frac{7}{4} < y_n < \frac{9}{4} \implies \frac{1}{16} < y_n^2 - 3 < \frac{33}{16},$$

and

$$|2y_n + 3| = |2(y_n - 2) + 7| \leq 2|y_n - 2| + 7 \leq 2(1) + 7 = 9.$$

Combining the two bounds, we have

$$|y_n - 2| < \frac{1}{4} \implies \left| \frac{y_n}{y_n^2 - 3} - 2 \right| \leq \frac{9}{\frac{1}{16}} |y_n - 2| = 144|y_n - 2|.$$

Take $\varepsilon' := \min \left\{ \frac{1}{4}, \frac{\varepsilon}{144} \right\}$. Since $\lim_n y_n = 2$, there exists $N \in \mathbb{N}$ such that

$$|y_n - 2| < \varepsilon' \quad \text{for all } n \geq N.$$

Now, for $n \geq N$, we have

$$\left| \frac{y_n}{y_n^2 - 3} - 2 \right| \leq 144|y_n - 2| < 144\varepsilon' \leq \varepsilon.$$



Example 2. Let (x_n) be a sequence of real numbers. Define

$$s_n = \frac{x_1 + x_2 + \cdots + x_n}{n} \quad \text{for all } n \in \mathbb{N}.$$

(a) If $\lim(x_n) = \ell$, where $\ell \in \mathbb{R}$, show that $\lim(s_n) = \ell$.

(b) Is the converse of (a) true?

Solution. (a) Without loss of generality, we assume that $\ell = 0$. (This can be done by letting $y_n = x_n - \ell$.)

For $1 \leq m < n$, we separate s_n into two parts:

$$s_n = \frac{x_1 + \cdots + x_m}{n} + \frac{x_{m+1} + \cdots + x_n}{n}.$$

In order to show that $|s_n|$ is small when n is large, we will use different approaches to estimate the size of the first and second part.

Since (x_n) is convergent, it is bounded, so we can find $M > 0$ such that

$$|x_n| \leq M \quad \text{for all } n \in \mathbb{N}.$$

Let $\varepsilon > 0$ be given. Since $\lim(x_n) = 0$, there exists $m \in \mathbb{N}$ such that

$$|x_n| < \varepsilon/2 \quad \text{for all } n \geq m.$$

By Archimedean Property, choose $N \in \mathbb{N}$ such that $N > \max\left\{\frac{mM}{\varepsilon/2}, m\right\}$.

Now, for $n \geq N$, we have

$$\begin{aligned} |s_n| &\leq \frac{|x_1| + \cdots + |x_m|}{n} + \frac{|x_{m+1}| + \cdots + |x_n|}{n} \\ &< \frac{mM}{n} + \frac{(n-m)\varepsilon/2}{n} \\ &< \varepsilon/2 + \varepsilon/2 \\ &= \varepsilon. \end{aligned}$$

Hence $\lim(s_n) = 0$.

(b) No. Consider $x_n := (-1)^n$. Then $s_n = \begin{cases} -\frac{1}{n} & n \text{ odd,} \\ 0 & n \text{ even.} \end{cases}$

Hence $\lim(s_n) = 0$ while (x_n) diverges.



2 Monotone Sequences

Monotone Convergence Theorem. *A monotone sequence of real number is convergent if and only if it is bounded. Furthermore,*

(a) *If (x_n) is a bounded increasing sequence, then $\lim(x_n) = \sup\{x_n : n \in \mathbb{N}\}$.*

(b) *If (y_n) is a bounded decreasing sequence, then $\lim(y_n) = \inf\{y_n : n \in \mathbb{N}\}$.*

Example 3. Establish the convergence or the divergence of the sequence (y_n) , where

$$y_n := \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \quad \text{for } n \in \mathbb{N}.$$

Example 4. Let (x_n) be the sequence defined by

$$x_1 := 1, \quad x_{n+1} := \frac{3 + 2x_n}{3 + x_n} \quad \text{for } n \geq 1.$$

Show that (x_n) is convergent and find its limit.