

MATH1050 Examples: Miscellanies on inequalities.

1. Let n be a positive integer.
 - (a) Prove that $k(n - k + 1) \geq n$ for each integer k amongst $1, 2, \dots, n$.
 - (b) Hence, or otherwise, prove that $(n!)^2 \geq n^n$.

2. (a) Let $n \in \mathbb{N} \setminus \{0\}$. Prove that $\frac{2n}{2n+1} < \frac{2n+1}{2n+2}$.

Remark. There is no need for mathematical induction.

- (b) Prove that $\prod_{k=1}^{5000} \frac{2k-1}{2k} < \frac{1}{100}$

3. (a) Prove that $\frac{2m-1}{2m} \leq \sqrt{\frac{3m-2}{3m+1}}$ for any $m \in \mathbb{N} \setminus \{0\}$.

Remark. There is no need to use mathematical induction.

- (b) Hence, or otherwise, prove that

$$\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot \dots \cdot \frac{2n-1}{2n} \leq \frac{1}{\sqrt{3n+1}} \quad \text{for any } n \in \mathbb{N} \setminus \{0\}$$

4. Here we take for granted the validity of the statement (#):

(#) Suppose $x, y \in \mathbb{R}$. Then $x^2 + y^2 \geq 2xy$. Moreover, equality holds iff $x = y$.

- (a) Prove the statement (b) below:

(b) Suppose $u, v, w \in \mathbb{R}$. Then $u^2 + v^2 + w^2 \geq uv + vw + wu$. Moreover equality holds iff $u = v = w$.

- (b) By applying the result described by (#), or otherwise, prove the statements below:

i. Suppose r, s, t be positive real numbers. Then $r + s + t \geq \sqrt{rs} + \sqrt{st} + \sqrt{tr}$.

ii. Suppose $x, y, z \in \mathbb{R}$. Then $x^2y^2 + y^2z^2 + z^2x^2 \geq xyz(x + y + z)$.

iii. Suppose a, b, c, d are positive real numbers. Then $(a+b)(a+c)(a+d)(b+c)(b+d)(c+d) \geq 64(abcd)^{3/2}$.

iv. Let p, q, r, s, t be positive real numbers. Suppose $pqrst = 1$. Then $(1+p)(1+q)(1+r)(1+s)(1+t) \geq 32$.

5. (a) Prove the statement below:

(#) Suppose x, y are positive real numbers. Then $\frac{x}{y} + \frac{y}{x} \geq 2$. Moreover, equality holds iff $x = y$.

- (b) By applying the result described by (#), or otherwise, prove the statements below:

i. Suppose $a > 1$ and $b > 1$. Then $\log_a(b) + \log_b(a) \geq 2$. Equality holds iff $a = b$.

ii. Suppose $u \in \mathbb{R}$. Then $\frac{u^2+2}{\sqrt{u^2+1}} \geq 2$. Equality holds iff $u = 0$.

iii. Suppose $v \in \mathbb{R}$. Then $\frac{v^2}{1+v^4} \leq \frac{1}{2}$. Equality holds iff ($v = 1$ or $v = -1$).

6. We introduce this definition below:

Let a, b, c be three positive real numbers (not necessarily distinct from each other). The numbers a, b, c are said to **constitute the three sides of a triangle** if the three inequalities $a + b > c$, $b + c > a$, $c + a > b$ hold simultaneously.

- (a) Let a, b be positive real numbers. Suppose $a \geq b$. Prove that there exists some positive real number c such that a, b, c constitute the three sides of a triangle.

Remark. For the geometric interpretation, see Proposition 22, Book I of *Euclid's Elements*.

- (b) Let a, b, c be positive real numbers. Suppose a, b, c constitute the three sides of a triangle. Prove that $\sqrt{a}, \sqrt{b}, \sqrt{c}$ constitute the three sides of a triangle.

(c) Let a, b, c be positive real numbers. Suppose a, b, c constitute the three sides of a triangle. Prove the statements below:

- i. $a^2 + b^2 + c^2 < 2(ab + bc + ca)$.
- ii. $3(ab + bc + ca) \leq (a + b + c)^2 < 4(ab + bc + ca)$.
- iii. $(a + b + c)(a + b - c) < 4ab$.

7. In this question, you may assume without proof the validity of the statement (#):

(#) For any real numbers μ, ν , if $0 < \mu < \nu < \frac{\pi}{2}$ then $0 < \sin(\mu) < \sin(\nu) < 1$.

Let the angles at vertices A, B, C in $\triangle ABC$ be α, β, γ respectively. Suppose each angle in $\triangle ABC$ is an acute angle. Prove the statements below:

- (a) $\cos(\frac{\gamma}{2}) > \sin(\frac{\gamma}{2})$.
- (b) $\sin(\alpha) + \sin(\beta) > \cos(\alpha) + \cos(\beta)$.
- (c) $\sin(\alpha) + \sin(\beta) + \sin(\gamma) > \cos(\alpha) + \cos(\beta) + \cos(\gamma)$.

8. (a) Prove the statement (#). (It may be easier to use the method of proof-by-contradiction.)

(#) Let a, b be real numbers. Suppose $|a| \leq 1$ and $|b| \leq 1$. Then $\sqrt{1 - a^2} + \sqrt{1 - b^2} \leq 2\sqrt{1 - \frac{(a + b)^2}{4}}$.

(b) Applying the statement (#), or otherwise, prove the statement (##).

(##) Let a, b, c, d be real numbers. Suppose $|a| \leq 1$ and $|b| \leq 1$ and $|c| \leq 1$ and $|d| \leq 1$.

Then $\sqrt{1 - a^2} + \sqrt{1 - b^2} + \sqrt{1 - c^2} + \sqrt{1 - d^2} \leq 4\sqrt{1 - \frac{(a + b + c + d)^2}{16}}$.

(c) Applying the statement (##), or otherwise, prove the statement (‡).

(‡) Let a, b, c be real numbers. Suppose $|a| \leq 1$ and $|b| \leq 1$ and $|c| \leq 1$.

Then $\sqrt{1 - a^2} + \sqrt{1 - b^2} + \sqrt{1 - c^2} \leq 3\sqrt{1 - \frac{(a + b + c)^2}{9}}$.

9. Prove the statement (#). (It may be easier to use the method of proof-by-contradiction.)

(#) Let a, b be real numbers. Suppose $ab \neq 0$. Then $\left| \frac{a + \sqrt{a^2 + 2b^2}}{2b} \right| < 1$ or $\left| \frac{a - \sqrt{a^2 + 2b^2}}{2b} \right| < 1$.

10. Prove the statement (#).

(#) Let a, n be positive integers. Suppose $n \geq a$. Then $(2a - 1)^n + (2a)^n < (2a + 1)^n$.

Remark. There is no need to apply mathematical induction.

11. Let α be a complex number. Suppose $0 < |\alpha| < 1$. Define the number β by $\beta = \frac{1}{|\alpha|} - 1$. Note that $\beta > 0$.

Let n be a positive integer.

(a) Suppose $n \geq 3$. By applying the Binomial Theorem, or otherwise, prove that $(1 + \beta)^n \geq \frac{n(n - 1)}{2}\beta^2$.

Hence deduce that $n|\alpha|^n \leq \frac{4}{n\beta^2}$.

(b) Suppose $n \geq 4$. By applying the Binomial Theorem prove that $(1 + \beta)^n \geq \frac{n(n - 1)(n - 2)}{6}\beta^3$.

Hence deduce that $n^2|\alpha|^n \leq \frac{36}{n\beta^3}$.

(c) Let k be a non-negative integer. Suppose $n \geq k + 2$.

By applying the Binomial Theorem, or otherwise, prove that $n^k|\alpha|^n \leq \frac{[(k + 1)!]^2}{n\beta^{k+1}}$.

Remark. The inequalities described here constitute the key step in the argument for the statement ‘ $\lim_{n \rightarrow \infty} n^k \alpha^n = 0$ ’.

12. For each $n \in \mathbb{N} \setminus \{0, 1\}$, define $a_n = \sqrt[n]{n} - 1$.

(a) Prove that $a_n \geq 0$ for any $n \in \mathbb{N} \setminus \{0, 1\}$.

(b) By applying the Binomial Theorem to the expression $(1 + a_n)^n$, prove that $a_n \leq \sqrt{\frac{2}{n-1}}$ for any $n \in \mathbb{N} \setminus \{0, 1\}$.

Remark. The inequalities described here constitute the key step in the argument for the statement ‘ $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$ ’.

13. Let $\alpha > 1$. For any $n \in \mathbb{N} \setminus \{0\}$, define $a_n = \alpha^{\frac{1}{n}} - 1$.

(a) Prove that $a_n > 0$ for any $n \in \mathbb{N} \setminus \{0\}$.

(b) By applying Bernoulli's Inequality, or otherwise, prove that $a_n \leq \frac{\alpha}{n}$ for any $n \in \mathbb{N} \setminus \{0\}$.

Remark. The inequalities described here constitute the key step in the argument for the statement ‘ $\lim_{n \rightarrow \infty} \alpha^{1/n} = 1$ ’.