

1. Definition.

Let A, B be sets, and $f : A \longrightarrow B, g : B \longrightarrow A$ be functions.

g is said to be an **inverse function** of f if both of the following statements hold:

- (a) For any $x \in A, (g \circ f)(x) = x.$ (b) For any $y \in B, (f \circ g)(y) = y.$

Definition.

Let C be a set.

Define the function $\text{id}_C : C \longrightarrow C$ by $\text{id}_C(z) = z$ for any $z \in C.$

id_C is called the **identity function on the set $C.$**

Remark 1 on the definition for the notion of inverse function.

By the respective definitions for the notions of inverse function, composition, and identity function:

$g : B \longrightarrow A$ is an inverse function of $f : A \longrightarrow B$
iff

$(g \circ f = \text{id}_A \text{ and } f \circ g = \text{id}_B \text{ as functions}).$

Definition.

Let A, B be sets, and $f : A \longrightarrow B, g : B \longrightarrow A$ be functions.

g is said to be an **inverse function** of f if both of the following statements hold:

- (a) For any $x \in A, (g \circ f)(x) = x.$ (b) For any $y \in B, (f \circ g)(y) = y.$

Remark 2 on the definition for the notion of inverse function.

See the ‘symmetry’ in the definition: simultaneously interchange

- ‘ A ’ and ‘ B ’, • ‘ f ’ and ‘ g ’, • ‘ x ’ and ‘ y ’.

What do we get?

Consequence of this ‘symmetry’ in the definition:

$g : B \longrightarrow A$ is an inverse function of $f : A \longrightarrow B$
iff

$f : A \longrightarrow B$ is an inverse function of $g : B \longrightarrow A.$

Definition.

Let A, B be sets, and $f : A \longrightarrow B, g : B \longrightarrow A$ be functions.


g is said to be an **inverse function** of f if both of the following statements hold:

- (a) For any $x \in A, (g \circ f)(x) = x.$ (b) For any $y \in B, (f \circ g)(y) = y.$


Remark 3 on the definition for the notion of inverse function.

How does such a function g ‘interact’ with f ?


- (a) Pick any $x \in A.$ x is ‘assigned’ by f to $f(x).$




- (b) Pick any $y \in B.$ y is ‘assigned’ by g to $g(y).$



But then $f(x)$ is ‘assigned’ by g to $x.$



But then $g(y)$ is ‘assigned’ by f to $y.$



So g ‘cancels’ what f does to $x.$

So f ‘cancels’ what g does to $y.$

Formal formulation: For any $x \in A,$ for any $y \in B, (y = f(x) \text{ iff } x = g(y)).$

2. Theorem (1). (Re-formulation of the definition of inverse function.)

Let A, B be sets, and $f : A \longrightarrow B, g : B \longrightarrow A$ be functions.

The statements below are logically equivalent:

- (\star_0) g is an inverse function of f .
- (\star_1) $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$ as functions.
- (\star_2) f is an inverse function of g .
- (\star_3) For any $x \in A$, for any $y \in B$, ($y = f(x)$ iff $x = g(y)$).

Proof of Theorem (1).

Let A, B be sets, and $f : A \longrightarrow B, g : B \longrightarrow A$ be functions.

Logical equivalence amongst (\star_0), (\star_1), (\star_2): see Remark 1, Remark 2 above.

- [$(\star_0) \implies (\star_3)$?] Exercise: carefully formulate the idea in Remark 3 above.
- [$(\star_3) \implies (\star_0)$?] Suppose that for any $x \in A, y \in B$, ($y = f(x)$ iff $x = g(y)$).

* Pick any $s \in A$. [Try to deduce: $(g \circ f)(s) = s$.]

Define $u = f(s)$. We have $u \in B$. By assumption, $s = g(u)$. Then $(g \circ f)(s) = g(f(s)) = g(u) = s$.

* Pick any $v \in B$. [Try to deduce: $(f \circ g)(v) = v$.]

Define $t = g(v)$. We have $t \in A$. By assumption, $v = f(t)$. Then $(f \circ g)(v) = f(g(v)) = f(t) = v$.

It follows that g is an inverse function of f .

3. Theorem (2). (Uniqueness of inverse function.)

Let A, B be sets, and $f : A \rightarrow B$ be a function.

f has at most one inverse function.

Proof of Theorem (2).

Let A, B be sets, and $f : A \rightarrow B$ be a function.

Suppose $g, h : B \rightarrow A$ are inverse functions of f .

[We want to deduce that $g(y) = h(y)$ for any $y \in B$.]

Pick any $y \in B$.

Define $x = g(y)$. We have $x \in A$.

$$\begin{aligned} \text{Then } y &\stackrel{\circledast}{=} (f \circ g)(y) \\ &= f(g(y)) = f(x). \end{aligned}$$

$$\begin{aligned} \text{Therefore } h(y) &= h(f(x)) = (h \circ f)(x) \\ &\stackrel{\circledast}{=} x \end{aligned}$$

$= g(y)$.
It follows that $g = h$ as functions. \square

Ask: What do we want to prove?
What is 'uniqueness'?

Answer. We want to prove:

- If there are two functions from B to A , both serving as inverse functions of f , then they are the same as each other as functions.

4. Definition.

Let D, R be sets and $h : D \longrightarrow R$ be a function.

h is said to be **bijective** if h is both surjective and injective.

Remark.

Hence h is bijective iff both of the statements (S) , (I) below hold:

(S) : For any $v \in R$, there exists some $u \in D$ such that $v = h(u)$.

(I) : For any $u, t \in D$, if $h(u) = h(t)$ then $u = t$.

5. Theorem (3). (Necessary condition for existence of inverse function.)

Let A, B be sets, $f : A \rightarrow B$ be a function.

Suppose f has an inverse function, say, $g : B \rightarrow A$.

Then each of f, g is bijective.

Proof of Theorem (3).

Let A, B be sets, $f : A \rightarrow B$ be a function.

Suppose f has an inverse function, say, $g : B \rightarrow A$.

- ['Is f surjective?'] [Want to deduce: For any $y \in B$, there exists some $x \in A$ such that $y = f(x)$.]
Pick any $y \in B$. Define $x = g(y)$. By definition, $x \in A$.
For the same $x \in A, y \in B$, we have $f(x) = f(g(y)) = (f \circ g)(y) = y$. Hence f is surjective.
- ['Is f injective?'] [Want to deduce: For any $x, w \in A$, if $f(x) = f(w)$ then $x = w$.]
Pick any $x, w \in A$. Suppose $f(x) = f(w)$. Then
$$x = (g \circ f)(x) = g(f(x)) = g(f(w)) = (g \circ f)(w) = w.$$
Hence f is injective.

By definition, g is an inverse function of f . Then by Theorem (1), g has an inverse function, namely, f .

It follows from the argument above that g is both surjective and injective. \square

Remark. The natural question to ask is: *Is the necessary condition sufficient?*

Answer. Yes, but...