

1. Recall:

(a) **Definition.**

Let  $A, B$  be sets. The set  $\text{Map}(A, B)$  is defined to be the set of all functions from  $A$  to  $B$ .

**Remark.**  $\text{Map}(\mathbb{N}, B)$  is the set of all infinite sequences in  $B$ : each  $\varphi \in \text{Map}(\mathbb{N}, B)$  is the infinite sequence  $(\varphi(0), \varphi(1), \varphi(2), \dots, \varphi(n), \varphi(n+1), \dots)$ .

(b) **Example ( $\epsilon$ ).** Let  $A$  be a set.  $\mathfrak{P}(A) \sim \text{Map}(A, \{0, 1\})$ .

(c) **Theorem (VI).** There is no surjective function from  $\mathbb{N}$  to  $\text{Map}(\mathbb{N}, \{0, 1\})$ .

(d) **Corollary (VII).** There is no bijective function from  $\mathbb{N}$  to  $\text{Map}(\mathbb{N}, \{0, 1\})$ . (Hence  $\mathbb{N} \not\sim \text{Map}(\mathbb{N}, \{0, 1\})$ .)

(e) **Theorem (VIII).** Let  $A$  be a set.  $A \not\sim \text{Map}(A, \{0, 1\})$ .  $A \not\sim \mathfrak{P}(A)$ .

2. **Theorem (XIII). (Baby version of Cantor's Theorem.)**

$\mathbb{N} < \text{Map}(\mathbb{N}, \{0, 1\})$ .

**Proof.**

By Corollary (VII),  $\mathbb{N} \not\sim \text{Map}(\mathbb{N}, \{0, 1\})$ . We now prove that  $\mathbb{N} \lesssim \text{Map}(\mathbb{N}, \{0, 1\})$ :

- For any  $n \in \mathbb{N}$ , define  $\delta_n : \mathbb{N} \rightarrow \{0, 1\}$  by

$$\delta_n(k) = \begin{cases} 1 & \text{if } k = n \\ 0 & \text{if } k \neq n \end{cases}$$

Define  $\Delta : \mathbb{N} \rightarrow \text{Map}(\mathbb{N}, \{0, 1\})$  by  $\Delta(n) = \delta_n$  for any  $n \in \mathbb{N}$ .

$\Delta$  is an injective function. (Why?)

Hence  $\mathbb{N} \lesssim \text{Map}(\mathbb{N}, \{0, 1\})$ .

We now have  $\mathbb{N} \lesssim \text{Map}(\mathbb{N}, \{0, 1\})$  and  $\mathbb{N} \not\sim \text{Map}(\mathbb{N}, \{0, 1\})$ . It follows that  $\mathbb{N} < \text{Map}(\mathbb{N}, \{0, 1\})$ .

3. **Theorem (XIV). (Cantor's Theorem.)**

Suppose  $A$  is a set. Then  $A < \text{Map}(A, \{0, 1\})$ , and  $A < \mathfrak{P}(A)$ .

**Proof.**

Let  $A$  be a set. By Theorem (VIII),  $A \not\sim \text{Map}(A, \{0, 1\})$ . We generalize the argument for Theorem (XIII) to prove that  $A \lesssim \text{Map}(A, \{0, 1\})$ :

- Recall that for any  $x \in A$ , the function  $\chi_{\{x\}}^A : A \rightarrow \{0, 1\}$  is given by

$$\chi_{\{x\}}^A(y) = \begin{cases} 1 & \text{if } y = x \\ 0 & \text{if } y \neq x \end{cases}$$

( $\chi_{\{x\}}^A$  is the characteristic function of  $\{x\}$  in  $A$ .)

Define the function  $\Delta : A \rightarrow \text{Map}(A, \{0, 1\})$  by  $\Delta(x) = \chi_{\{x\}}^A$  for any  $x \in A$ .  $\Delta$  is an injective function from  $A$  to  $\text{Map}(A, \{0, 1\})$ . (Why?) Hence  $A \lesssim \text{Map}(A, \{0, 1\})$ .

We now have  $A \lesssim \text{Map}(A, \{0, 1\})$  and  $A \not\sim \text{Map}(A, \{0, 1\})$ .

It follows that  $A < \text{Map}(A, \{0, 1\})$ . Since  $\mathfrak{P}(A) \sim \text{Map}(A, \{0, 1\})$ , we have  $A < \mathfrak{P}(A)$ . (Why?)

4. **Question.** Note that  $\mathbb{Q} \lesssim \mathbb{R}$ . Is it true that  $\mathbb{Q} \sim \mathbb{R}$ , or that  $\mathbb{Q} < \mathbb{R}$ ?

**Lemma (XV).**

Let  $A, B, C$  be sets. Suppose  $A \lesssim B$  and  $B \lesssim C$ . Also suppose  $A < B$  or  $B < C$ . Then  $A < C$ .

**Proof.**

Let  $A, B, C$  be sets. Suppose  $A \lesssim B$  and  $B \lesssim C$ . Also suppose  $A < B$  or  $B < C$ .

Since  $A \lesssim B$  and  $B \lesssim C$ , we have  $A \lesssim C$ .

Since  $A < B$  or  $B < C$ , we have  $A \not\sim B$  or  $B \not\sim C$ . We verify that  $A \not\sim C$ :

- Suppose it were true that  $A \sim C$ . Then  $C \lesssim A$ .

Since  $B \lesssim C$  and  $C \lesssim A$ , we would have  $B \lesssim A$ . Then, since  $A \lesssim B$  and  $B \lesssim A$ , we would have  $A \sim B$  according to the Schröder-Bernstein Theorem.

Since  $C \lesssim A$  and  $A \lesssim B$ , we would have  $C \lesssim B$ . Then, since  $B \lesssim C$  and  $C \lesssim B$ , we would have  $B \sim C$  according to the Schröder-Bernstein Theorem.

Hence  $A \sim B$  and  $B \sim C$ . But by assumption,  $A \not\sim B$  or  $B \not\sim C$ . Contradiction arises. Hence  $A \not\sim C$  in the first place.

Then, since  $A \lesssim C$  and  $A \not\sim C$ , we have  $A < C$ .

**Theorem (XVI).**

$\mathbb{N} < [0, 1]$ , and  $\mathbb{N} < \mathbb{R}$ , and  $\mathbb{Q} < \mathbb{R}$ .

**Proof.**

$\mathbb{N} \lesssim \text{Map}(\mathbb{N}, \{0, 1\}) \lesssim \text{Map}(\mathbb{N}, [0, 9]) \sim [0, 1] \sim \mathbb{R}$ .

Also,  $\mathbb{N} < \text{Map}(\mathbb{N}, \{0, 1\})$ .

Then, by Lemma (XV),  $\mathbb{N} < [0, 1]$  and  $\mathbb{N} < \mathbb{R}$ .

Since  $\mathbb{Q} \sim \mathbb{N}$ , we also have  $\mathbb{Q} < \mathbb{R}$ .

**Remark.** Hence there are much much more real numbers than there are rational numbers.

5. **Question.** Why are ‘Venn diagram arguments’ not good enough?

**Theorem (XVII.)**

There exists some set  $T$  such that  $S < T$  for any subset  $S$  of  $\mathbb{R}^2$ .

**Proof.**

Define  $T = \mathfrak{P}(\mathbb{R})$ .

Pick any subset  $S$  of  $\mathbb{R}^2$ . We have  $S \lesssim \mathbb{R}^2 \sim \mathbb{R}$ . By Cantor’s Theorem,  $\mathbb{R} < \mathfrak{P}(\mathbb{R}) = T$ . Then by Lemma (XV), we have  $S < T$ .

**Remark.**

When we draw a Venn diagram for a set, say,  $A$ , we are ‘identifying’ the set  $A$  with some subset, say,  $B$ , of  $\mathbb{R}^2$ , in the sense that the elements of  $A$  are ‘identified’ as the points in  $B$ , via some bijective function from  $A$  to  $B$ . This bijective function guarantees that distinct elements of  $A$  are identified as distinct points of  $B$ . So we are implicitly assuming that there is an injective function from  $A$  to  $\mathbb{R}^2$ .

But now we know that there are sets which are too ‘large’ to be drawn in a Venn diagram.

6. **Question.** Is there any ‘universal set’, which contains every conceivable object as its element?

**Theorem (XVIII.)**

Denote  $\{x \mid x = x\}$  by  $U$ . The mathematical object  $U$  is not a set.

**Proof.**

Suppose  $U$  were a set. Then, by Cantor’s Theorem,  $U < \text{Map}(U, \{0, 1\})$ .

For any  $\varphi \in \text{Map}(U, \{0, 1\})$ , we would have  $\varphi = \varphi$ , and hence  $\varphi \in U$ .

It would follow that  $\text{Map}(U, \{0, 1\}) \subset U$ .

Then  $\text{Map}(U, \{0, 1\}) \lesssim U$ . Therefore  $U < \text{Map}(U, \{0, 1\}) \lesssim U$ .

By Lemma (XV),  $U < U$ . In particular,  $U \not\sim U$ . There would be no bijective function from  $U$  to  $U$ . But  $\text{id}_U$  is a bijective function from  $U$  to  $U$ . Contradiction arises.

Hence  $U$  is not a set in the first place.

**Remark.** Hence if we insist Cantor’s Theorem to be a true statement, then there is no such thing as a ‘universal set’. This is known as **Cantor’s Paradox**.