

CMSC5724: Quiz 3

Problem 1 (50%). Consider the kernel function $K(p, q) = (2(p \cdot q) + 1)^2$, where $p = (p[1], p[2])$ and $q = (q[1], q[2])$ are 2D vectors. Recall that there is a mapping function ϕ from \mathbb{R}^2 to \mathbb{R}^d for some integer d , such that $K(p, q)$ equals the dot product of $\phi(p)$ and $\phi(q)$. Give the details of ϕ .

Answer: Rewrite K as dot product form.

$$\begin{aligned} K(p, q) &= (2p[1]q[1] + 2p[2]q[2] + 1)^2 \\ &= 4p[1]^2q[1]^2 + 4p[2]^2q[2]^2 + 8p[1]p[2]q[1]q[2] + 4p[1]q[1] + 4p[2]q[2] + 1. \end{aligned}$$

Hence, $\phi(p) = (2p[1]^2, 2p[2]^2, 2\sqrt{2}p[1]p[2], 2p[1], 2p[2], 1)$.

Problem 2 (50%). Consider a 3-class linear classifier in 2D space that is defined by vectors $\mathbf{w}_1 = (3, 5)$, $\mathbf{w}_2 = (-2, 9)$, and $\mathbf{w}_3 = (0, 7)$. Given a point $\mathbf{p} = (-5, 1)$, explain what is the label assigned to \mathbf{p} and why.

Answer: Computing the dot product between each \mathbf{w}_i and \mathbf{p} where $i \in [1, 3]$, we have:

- $\mathbf{w}_1 \cdot \mathbf{p} = -10$;
- $\mathbf{w}_2 \cdot \mathbf{p} = 19$;
- $\mathbf{w}_3 \cdot \mathbf{p} = 7$.

Since $\mathbf{w}_2 \cdot \mathbf{p}$ is largest, the label assigned to \mathbf{p} is 2.