

Case (b) $L = \pm\infty$. (of Thm 6.3.5)

subcase $L = +\infty$

$\lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)} = +\infty$ implies that

$$\forall M > 1, \exists \delta > 0 \text{ s.t. } \frac{f'(u)}{g'(u)} > M$$

and hence $\frac{f(\beta) - f(\alpha)}{g(\beta) - g(\alpha)} > M, \forall a < \alpha < \beta < a + \delta$

As in case (a), $\lim_{x \rightarrow a^+} g(x) = +\infty$ implies

$\exists c \ \& \ c_1 \in (a, a + \delta)$ such that

- $a < c_1 < c < a + \delta$
- $g(\alpha) > 0, \forall \alpha \in (a, c]$
- $0 < \frac{g(c)}{g(\alpha)} < \frac{1}{2}, \forall \alpha \in (a, c_1)$
- $0 \leq \frac{|f(c)|}{g(\alpha)} < \frac{1}{2}, \forall \alpha \in (a, c_1)$

Letting $\beta = c$, we have

$$\frac{f(c) - f(\alpha)}{g(c) - g(\alpha)} > M, \forall \alpha \in (a, c)$$

(\leftarrow both terms < 0)

And hence for $\alpha \in (a, c_1)$

$$\frac{f(\alpha) - f(c)}{g(\alpha)} > M \left(1 - \frac{g(c)}{g(\alpha)}\right) > \frac{1}{2} M, \forall \alpha \in (a, c_1)$$

($\leftarrow (1 - \frac{g(c)}{g(\alpha)}) > \frac{1}{2}$)

$$\Rightarrow \frac{f(x)}{g(x)} > \frac{1}{2}M + \frac{f(c)}{g(x)} > \frac{1}{2}M - \frac{1}{2}$$

$$= \frac{1}{2}(M-1), \quad \forall x \in (a, c_1)$$

Since $M > 1$ is arbitrary, this shows that

$$\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = +\infty.$$

Subcase of " $L = -\infty$ " is similar. ~~✗~~

eg 6.3.6

$$(a) \lim_{x \rightarrow \infty} \frac{\ln x}{x}$$

- $f(x) = \ln x$ has derivative $f'(x) = \frac{1}{x}$ on $(0, \infty)$
- $g(x) = x$ has derivative $g'(x) = 1 \neq 0$ on $(0, \infty)$
- $\lim_{x \rightarrow \infty} g(x) = +\infty$
- $\lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow \infty} \frac{1/x}{1} = 0$

$$\therefore \text{L'Hospital Rule II} \Rightarrow \lim_{x \rightarrow \infty} \frac{\ln x}{x} = 0.$$

(usually, one simply write $\lim_{x \rightarrow \infty} \frac{\ln x}{x} = \lim_{x \rightarrow \infty} \frac{1/x}{1} = 0$)

$$(b) \quad \lim_{x \rightarrow \infty} e^{-x} x^2 = \lim_{x \rightarrow \infty} \frac{x^2}{e^x}$$

$$\bullet (x^2)' = 2x, \quad \forall x$$

$$\bullet (e^x)' = e^x \neq 0, \quad \forall x$$

$$\bullet e^x \rightarrow +\infty \text{ as } x \rightarrow +\infty$$

But $\lim_{x \rightarrow \infty} \frac{2x}{e^x}$ still indeterminate.

So we need to start with $\lim_{x \rightarrow \infty} \frac{2x}{e^x}$ first:

$$\bullet (2x)' = 2, \quad \forall x$$

$$\bullet (e^x)' = e^x \neq 0, \quad \forall x$$

$$\bullet e^x \rightarrow +\infty \text{ as } x \rightarrow +\infty$$

$$\bullet \lim_{x \rightarrow \infty} \frac{2}{e^x} = 0 \text{ (exists)}$$

$$\therefore \text{L'Hospital Rule} \Rightarrow \lim_{x \rightarrow \infty} \frac{2x}{e^x} = 0 \text{ (exists)}$$

And applying L'Hospital Rule again, $\lim_{x \rightarrow \infty} \frac{x^2}{e^x} = 0$.

(We usually just write

$$\lim_{x \rightarrow \infty} e^{-x} x^2 = \lim_{x \rightarrow \infty} \frac{x^2}{e^x} = \lim_{x \rightarrow \infty} \frac{2x}{e^x} = \lim_{x \rightarrow \infty} \frac{2}{e^x} = 0.)$$

$$\begin{aligned}
 (c) \quad \lim_{x \rightarrow 0^+} \frac{\ln \sin x}{\ln x} &= \lim_{x \rightarrow 0^+} \frac{(\ln \sin x)'}{(\ln x)'} \\
 &= \lim_{x \rightarrow 0^+} \frac{\frac{\cos x}{\sin x}}{\frac{1}{x}} = \lim_{x \rightarrow 0^+} \cos x \cdot \frac{x}{\sin x} \\
 &= 1 \quad \left(\text{as } \lim_{x \rightarrow 0^+} \frac{x}{\sin x} = 1 = \lim_{x \rightarrow 0^+} \cos x \right)
 \end{aligned}$$

(d) It is easy to see $\lim_{x \rightarrow \infty} \frac{x - \sin x}{x + \sin x} = \lim_{x \rightarrow \infty} \frac{1 - \frac{\sin x}{x}}{1 + \frac{\sin x}{x}} = 1$.

However, $\lim_{x \rightarrow \infty} \frac{(x - \sin x)'}{(x + \sin x)'} = \lim_{x \rightarrow \infty} \frac{1 - \cos x}{1 + \cos x}$ doesn't exist.

\therefore The condition " $\lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)}$ exists" is necessary

for L'Hospital Rule.

Further examples (other indeterminate forms)

eg 6.3.7

(a) ($\infty - \infty$ form)

$$\lim_{x \rightarrow 0^+} \left(\frac{1}{x} - \frac{1}{\sin x} \right) \quad \left(x \in \left(0, \frac{\pi}{2} \right) \right)$$

$$= \lim_{x \rightarrow 0^+} \frac{\sin x - x}{x \sin x} \quad \left(\text{transform to } \frac{0}{0} \text{ form} \right)$$

$$= \lim_{x \rightarrow 0^+} \frac{\cos x - 1}{\sin x + x \cos x} \quad (\text{L'Hospital}) \quad (\text{still } \frac{0}{0} \text{ form})$$

$$= \lim_{x \rightarrow 0^+} \frac{-\sin x}{2 \cos x - x \sin x} \quad (\text{L'Hospital})$$

$$= 0 \quad (\text{limit exists, calculation justified})$$

(b) $(0 \cdot (-\infty))$ form

$$\lim_{x \rightarrow 0^+} x \ln x \quad (x \in (0, \infty))$$

$$= \lim_{x \rightarrow 0^+} \frac{\ln x}{\frac{1}{x}} \quad (\text{transforms to } \frac{-\infty}{\infty} \text{ form})$$

$$= \lim_{x \rightarrow 0^+} \frac{\frac{1}{x}}{-\frac{1}{x^2}} \quad (\text{L'Hospital})$$

$$= \lim_{x \rightarrow 0^+} (-x) = 0 \quad (\text{limit exists, calculation justified})$$

(c) (0^0) form

$$\lim_{x \rightarrow 0^+} x^x$$

$$= \lim_{x \rightarrow 0^+} e^{x \ln x}$$

$$= e^{\lim_{x \rightarrow 0^+} x \ln x}$$

$$= e^0 = 1$$

(transforms to $0 \cdot (-\infty)$ form which can be calculated using L'Hospital as in (b))

(d) (1^∞ form)

$$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x \quad x \in (1, \infty)$$

$$= \lim_{x \rightarrow \infty} e^{x \ln\left(1 + \frac{1}{x}\right)}$$

Transforms to the calculation of

$$\lim_{x \rightarrow \infty} x \ln\left(1 + \frac{1}{x}\right) \quad (\infty \cdot 0 \text{ form})$$

$$= \lim_{x \rightarrow \infty} \frac{\ln\left(1 + \frac{1}{x}\right)}{\frac{1}{x}} \quad (\text{transform to } \frac{0}{0} \text{ form})$$

$$= \lim_{x \rightarrow \infty} \frac{\left(\frac{1}{1 + \frac{1}{x}}\right) \cdot \left(-\frac{1}{x^2}\right)}{\left(-\frac{1}{x^2}\right)} \quad (\text{L'Hospital})$$

$$= \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{1}{x}} = 1 \quad (\text{limit exists, calculation justified})$$

$$\text{And hence } \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e^{\lim_{x \rightarrow \infty} x \ln\left(1 + \frac{1}{x}\right)} = e$$

(e) (∞^0 form)

$$\lim_{x \rightarrow 0^+} \left(1 + \frac{1}{x}\right)^x \quad (x \in (0, \infty)) \quad (\text{limit of the other end})$$

$$= e^{\lim_{x \rightarrow 0^+} x \ln\left(1 + \frac{1}{x}\right)} \quad (\text{transform to } 0 \cdot \infty \text{ form})$$

$$= e^{\lim_{x \rightarrow 0^+} \frac{1}{1 + \frac{1}{x}}} \quad (\text{L'Hospital as before})$$

$$= e^0 = 1 \quad (\text{limit exists, calculation justified})$$