

**MATH 2028 Honours Advanced Calculus II**  
**2022-23 Term 1**  
**Problem Set 11**

*due on Dec 7, 2022 (Wednesday) at 11:59PM*

**Instructions:** You are allowed to discuss with your classmates or seek help from the TAs but you are required to write/type up your own solutions. You can either type up your assignment or scan a copy of your written assignment into ONE PDF file and submit through Blackboard on/before the due date. Please remember to write down your name and student ID. **No late homework will be accepted.**

**Problems to hand in**

1. Compute the area of the surface in  $\mathbb{R}^4$  parametrized by

$$g(u, v) = (u, v, u^2 - v^2, 2uv)$$

with  $(u, v) \in \mathbb{R}^2$  satisfying  $u^2 + v^2 \leq 1$ .

2. Let  $\Omega \subset \mathbb{R}^3$  be the region bounded above by the sphere  $x^2 + y^2 + z^2 = a^2$  and below by the plane  $z = 0$ . Compute

$$\int_{\partial\Omega} xz \, dy \wedge dz + yz \, dz \wedge dx + (x^2 + y^2 + z^2) \, dx \wedge dy$$

directly and by applying Stokes' Theorem.

3. (a) Suppose  $M$  and  $M'$  are two compact oriented  $k$ -dimensional submanifolds of  $\mathbb{R}^n$  with boundary, and suppose  $\partial M = \partial M'$ . Prove that for any  $(k-1)$  form  $\omega$ , we have

$$\int_M d\omega = \int_{M'} d\omega.$$

- (b) Use (a) to compute  $\int_M d\omega$  where  $M$  is the upper hemisphere  $x^2 + y^2 + z^2 = a^2$ ,  $z \geq 0$ , oriented with outward-pointing normal having positive  $z$ -component and

$$\omega = (x^3 + 3x^2y - y) \, dx + (y^3z + x + x^3) \, dy + (x^2 + y^2 + z) \, dz.$$

**Suggested Exercises**

1. Check that the boundary orientation on  $\partial\mathbb{R}_+^k$  is  $(-1)^k$  times the usual orientation on  $\mathbb{R}^{k-1}$ .
2. Let  $C$  be the intersection of the cylinder  $x^2 + y^2 = 1$  and the plane  $2x + 3y - z = 1$ , oriented counterclockwise as viewed from high above the  $xy$ -plane. Evaluate

$$\int_C y \, dx - 2z \, dy + x \, dz$$

directly and by applying Stokes' Theorem.

3. Compute  $\int_C (y - z) \, dx + (z - x) \, dy + (x - y) \, dz$  where  $C$  is the intersection of the cylinder  $x^2 + y^2 = a^2$  and the plane  $\frac{x}{a} + \frac{z}{b} = 1$ , oriented clockwise as viewed from high above the  $xy$ -plane.

4. Let  $C$  be the intersection of the sphere  $x^2 + y^2 + z^2 = a^2$  and the plane  $x + y + z = 0$ , oriented counterclockwise as viewed from high above the  $xy$ -plane. Evaluate

$$\int_C 2z \, dx + 3x \, dy - dz.$$

5. Let  $\omega = y^2 \, dy \wedge dz + x^2 \, dz \wedge dx + z^2 \, dx \wedge dy$ , and  $M$  be the solid paraboloid  $0 \leq z \leq 1 - x^2 - y^2$ . Evaluate  $\int_{\partial M} \omega$  directly and by applying Stokes' Theorem.

6. Let  $M$  be the surface of the paraboloid  $z = 1 - x^2 - y^2 \geq 0$ , oriented so that the outward-pointing normal has positive  $z$ -component. Let  $F(x, y, z) = (x^2z, y^2z, x^2 + y^2)$ . Compute  $\int_M F \cdot \vec{n} \, d\sigma$  directly and by applying Stokes' Theorem.

7. Compute  $\int_M d\omega$  where  $M$  is the portion of the paraboloid  $z = x^2 + y^2$  lying beneath  $z = 4$ , oriented with outward-pointing normal having positive  $z$ -component, and  $\omega = y \, dx + z \, dy + x \, dz$ .

8. Let  $M = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : x_1^2 + x_2^2 + x_3^2 \leq x_4 \leq 1\}$ , with the standard orientation inherited from  $\mathbb{R}^4$ . Evaluate

$$\int_{\partial M} (x_1^3 x_2^4 + x_4) \, dx_1 \wedge dx_2 \wedge dx_3.$$

9. Let  $S$  be the portion of the cylinder  $x^2 + y^2 = a^2$  lying above the  $xy$ -plane and below the sphere  $x^2 + (y - a)^2 + z^2 = 4a^2$ . Let  $C$  be the intersection of the cylinder and sphere, oriented clockwise as viewed from high above the  $xy$ -plane.

(a) Evaluate  $\int_S z \, d\sigma$ .

(b) Use (a) to compute  $\int_C y(z^2 - 1) \, dx + x(1 - z^2) \, dy + z^2 \, dz$ .

10. Let  $C$  be the intersection of the sphere  $x^2 + y^2 + z^2 = 1$  and the plane  $x + y + z = 0$ , oriented counterclockwise as viewed from high above the  $xy$ -plane. Evaluate  $\int_C z^3 \, ds$ .

## Challenging Exercises

1. Let  $f : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  be a smooth function whose graph is the surface  $S$ .

(a) Consider the area 2-form  $\sigma$  on  $S$  given by

$$\sigma = \frac{1}{\sqrt{1 + |\nabla f|^2}} \left( -\frac{\partial f}{\partial x} \, dy \wedge dz - \frac{\partial f}{\partial y} \, dz \wedge dx + dx \wedge dy \right).$$

Show that  $d\sigma = 0$  if and only if  $f$  satisfies the minimal surface equation:

$$\left( 1 + \left( \frac{\partial f}{\partial y} \right)^2 \right) \frac{\partial^2 f}{\partial x^2} - 2 \frac{\partial f}{\partial x} \frac{\partial f}{\partial y} \frac{\partial^2 f}{\partial x \partial y} + \left( 1 + \left( \frac{\partial f}{\partial x} \right)^2 \right) \frac{\partial^2 f}{\partial y^2} = 0.$$

(b) Show that for any compact oriented surface  $N \subset \mathbb{R}^3$ , we have

$$\int_N \sigma \leq \text{area}(N)$$

and equality holds if and only if  $N$  is parallel to  $S$ .

- (c) Suppose further that  $\partial N = \partial S$ . Prove that  $\text{area}(S) \leq \text{area}(N)$ .
2. (a) Prove that a  $k$ -dimensional submanifold with boundary  $M \subset \mathbb{R}^n$  is orientable if and only if there is a nowhere-zero  $k$ -form on  $M$ .
- (b) Show that  $M$  is orientable if and only if there is a volume form globally defined on  $M$ .