

MATH 2050 C Lecture 11 (Feb 21)

Q: Can we determine the limit of (x_n) exist without knowing the value of the limit?

Last time: limit thms, squeeze thm, ratio test

Today: "Monotone Convergence Thm"


Recall: (x_n) convergent \Rightarrow (x_n) bdd
~~↗~~
false

Cor: (x_n) unbdd \Rightarrow (x_n) divergent. "Divergence Test"

Counterexample: $(x_n) = (-1)^n$ is bdd BUT divergent

Pf: Suppose (x_n) is convergent, say $\lim(x_n) = a \in \mathbb{R}$.

Take $\varepsilon = 1$, $\exists K \in \mathbb{N}$ s.t. $|x_n - a| < \varepsilon = 1 \quad \forall n \geq K$



For $n \geq K$ is odd, we have

$$|x_n - a| = |-1 - a| < 1$$

$$\Rightarrow -2 < a < 0$$

For $n \geq K$ is even, we have

$$|x_n - a| = |1 - a| < 1$$

$$\Rightarrow 0 < a < 2$$

Contradiction!

Q: Under what condition(s) does

(x_n) bdd $\Rightarrow (x_n)$ convergent?

Monotone Convergence Theorem (MCT)

(x_n) bdd + monotone $\Rightarrow (x_n)$ convergent


Defⁿ: (x_n) is monotone if it is

either (i) increasing, i.e. $x_1 \leq x_2 \leq x_3 \leq \dots$ ($x_n \leq x_{n+1} \forall n \in \mathbb{N}$)

or (ii) decreasing, i.e. $x_1 \geq x_2 \geq x_3 \geq \dots$ ($x_n \geq x_{n+1} \forall n \in \mathbb{N}$)

Note: If inequalities are strict, then we say it is strictly monotone / increasing / decreasing.

Picture:



Proof of MCT: Idea: $\lim (x_n) = \sup \{x_n \mid n \in \mathbb{N}\}$

Suppose (x_n) is bdd and increasing. Consider

$$\emptyset \neq S := \{x_n \mid n \in \mathbb{N}\} \subseteq \mathbb{R}$$

Note (x_n) is bdd $\Rightarrow S$ is bdd above & below

By completeness of \mathbb{R} , $x := \sup S$ exists.

Claim: $\lim (x_n) = x$

Pf of Claim: We show this using ϵ - K defⁿ of limit.

Let $\epsilon > 0$ be fixed but arbitrary.

Since $x = \sup S$, $x - \epsilon$ CANNOT be an upper bd for S

i.e. $\exists K \in \mathbb{N}$ st. $x - \epsilon < x_K$

Since (x_n) is increasing (i.e. $x_n \leq x_{n+1} \forall n \in \mathbb{N}$)

\Rightarrow ①: $x - \epsilon < x_K \leq x_{K+1} \leq x_{K+2} \leq \dots \leq x_n \quad \forall n \geq K$

On the other hand, $x = \sup S$ is an upper bd for S

\Rightarrow ②: $x_n \leq x < x + \epsilon \quad \forall n \in \mathbb{N}$

Combining ① & ②.

$$x - \epsilon < x_n < x + \epsilon \quad \forall n \geq K$$

Example 1 "Harmonic series"

Let $h_n := 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} \quad , \quad n \in \mathbb{N}$.

i.e. $h_1 = 1$, $h_2 = 1 + \frac{1}{2} = \frac{3}{2}$, ...

Show that (h_n) is divergent.

Pf: Note $h_{n+1} = h_n + \frac{1}{n+1} > h_n \quad \forall n \in \mathbb{N}$

i.e. (h_n) is strictly increasing!

By MCT, (h_n) divergent \Leftrightarrow (h_n) unbdd

Claim: (h_n) is unbdd!

Consider $n = 2^m$, $m \in \mathbb{N}$.

$$\begin{aligned} h_{2^m} &= 1 + \underbrace{\frac{1}{2}}_{1 \text{ term}} + \underbrace{\left(\frac{1}{3} + \frac{1}{4}\right)}_{2 \text{ terms}} + \dots + \underbrace{\left(\frac{1}{2^{m-1}+1} + \dots + \frac{1}{2^m}\right)}_{2^{m-1} \text{ terms}} \\ &> 1 + \frac{1}{2} + \underbrace{\left(\frac{1}{4} + \frac{1}{4}\right)}_{2 \text{ terms}} + \dots + \underbrace{\left(\frac{1}{2^m} + \dots + \frac{1}{2^m}\right)}_{2^{m-1} \text{ terms}} \\ &= \underbrace{1 + \frac{1}{2} + \frac{1}{2} + \dots + \frac{1}{2}}_{m \text{ terms}} = 1 + \frac{m}{2} \end{aligned}$$

... ∞

$$\begin{aligned} h_1 &= 1 \\ h_4 &= 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} \\ &> 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{4} \\ &= 1 + \frac{1}{2} \end{aligned}$$

Unbdd as $m \rightarrow \infty$.

$\Rightarrow (h_n)$ is unbdd.

Remark: MCT works well for recursive sequence.

Example 2: Let (y_n) be defined "recursively" by:

$$y_1 := 1 ; \quad y_{n+1} := \frac{1}{4}(2y_n + 3) \quad \forall n \in \mathbb{N}$$

Show that $\lim (y_n) = \frac{3}{2}$.

Proof: General Strategy

- Step 1: Apply MCT to show the limit first
- Step 2: Take limit in the recursive relation (*) to compute the limit of the seq.

We first show that (y_n) is bdd & monotone.

Claim: (y_n) is bdd above by 2.

Pf of claim: Use M.I. Note $y_1 := 1 < 2$.

Suppose $y_k < 2$. Then, $y_{k+1} = \frac{1}{4}(2y_k + 3) < \frac{7}{4} < 2$.

... ∞

$$\begin{aligned} y_1 &= 1 \\ y_2 &= \frac{1}{4}(2+3) = \frac{5}{4} \\ y_3 &= \frac{1}{4}\left(2 \cdot \frac{5}{4} + 3\right) = \frac{11}{8} \end{aligned}$$

Claim: (y_n) is increasing, i.e. $y_n \leq y_{n+1} \quad \forall n \in \mathbb{N}$.

Pf of Claim: Use M.I. Note $y_1 := 1 < \frac{5}{4} = y_2$.

Assume $y_k \leq y_{k+1}$. Then

$$y_{k+1} = \frac{1}{4}(2y_k + 3) \leq \frac{1}{4}(2y_{k+1} + 3) = y_{k+2}.$$

So (y_n) is bdd & monotone, by MCT, $\lim(y_n) = y$ exists.

Since (y_n) is convergent, we have $\lim(y_{n+1}) = \lim(y_n) = y$

Take $n \rightarrow \infty$ on both sides of $(*)$:

$$\lim(y_{n+1}) = \lim \frac{1}{4}(2y_n + 3) \stackrel{\substack{\uparrow \\ \text{Limit} \\ \text{Thm}}}{=} \frac{1}{4}(2 \lim(y_n) + 3)$$

$$\Rightarrow y = \frac{1}{4}(2y + 3)$$

Solving for y , get $y = \frac{3}{2}$.

Example 3: Fix $a > 0$. Define inductively

$$S_1 := 1; \quad \underline{S_{n+1} := \frac{1}{2} \left(S_n + \frac{a}{S_n} \right)} \quad \forall n \in \mathbb{N} \quad (**)$$

Show that $\lim(S_n) = \sqrt{a} > 0$.

Proof: Claim 1: (S_n) is bdd below by \sqrt{a} (for $n \geq 2$)

Pf of Claim: Note $S_n > 0 \quad \forall n \in \mathbb{N}$. Rewrite $(**)$ as

$$S_n^2 - 2S_{n+1}S_n + a = 0$$

So, $x^2 - 2S_{n+1}x + a = 0$ has at least a real root S_n

$$\Rightarrow 4S_{n+1}^2 - 4a \geq 0 \Rightarrow S_{n+1} \geq \sqrt{a} \quad \forall n \in \mathbb{N}$$

Claim 2: (S_n) is decreasing "eventually", i.e. $S_n \geq S_{n+1} \quad \forall n \geq 2$.

Pf of Claim: $\forall n \geq 2$,

use Claim 1

$$S_n - S_{n+1} = S_n - \frac{1}{2} \left(S_n + \frac{a}{S_n} \right) = \frac{1}{2} \left(\frac{S_n^2 - a}{S_n} \right) \geq 0$$

By MCT, $\lim (S_n) =: S$ exists.

Take $n \rightarrow \infty$ on both sides of (**), then we obtain

$$S = \frac{1}{2} \left(S + \frac{a}{S} \right) \quad \left(\begin{array}{l} \text{Note: } S_n \geq \sqrt{a} \quad \forall n \geq 2 \\ \Rightarrow S \geq \sqrt{a} > 0. \end{array} \right)$$

Solve for S

$$\Rightarrow S = \sqrt{a} > 0.$$

MIDTERM UP TO HERE

Subsequences (§ 3.4 in textbook)

Defⁿ: Let $(x_n)_{n \in \mathbb{N}}$ be a seq. of real numbers.

Suppose $n_1 < n_2 < n_3 < \dots$ be a strictly increasing seq. of natural no.

THEN.

$$(x_{n_k})_{k \in \mathbb{N}} := (x_{n_1}, x_{n_2}, x_{n_3}, \dots, x_{n_k}, \dots)$$

is called a **subsequence** of $(x_n)_{n \in \mathbb{N}}$.

k^{th} term of (x_{n_k})

"
 n_k^{th} term of (x_n)

Intuitively:

$$(x_n) = (x_1, x_2, x_3, x_4, x_5, x_6, \dots)$$

$$(x_{n_k}) = (x_1, x_2, x_4, x_6, \dots)$$

$k=1$

$k=2$

$k=3$

$k=4$

$n_1=1$

$n_2=2$

$n_3=4$

$n_4=6$

E.g.) (Tail of a seq.) For each fixed $l \in \mathbb{N}$, then

the l -tail $(x_{k+l})_{k \in \mathbb{N}}$ is a subsequence of $(x_n)_{n \in \mathbb{N}}$

(Here, $n_k = k + l$)

E.g.) $(x_n) = (-1)^n$

Then $(1, 1, 1, \dots, 1, \dots)$ is a subseq.