

## MATH 2050C Lecture 4 (Jan 19)

- Midterm date fixed Mar 2, 2023 in-class 4:30-6:15pm
- Quiz 1 on Feb 2

Goal:  $\mathbb{R}$  is a complete ordered field.

Completeness Property: Every  $\emptyset \neq S \subseteq \mathbb{R}$  which is bounded above must have a supremum in  $\mathbb{R}$ . [Note:  $\mathbb{Q}$  fails this!]

Last time, we proved:

Prop:  $u = \sup S$  iff

①  $s \leq u$ ,  $\forall s \in S$  (i.e.  $u$  is an upper bd.)

②  $\forall \varepsilon > 0$ ,  $\exists s' \in S$  s.t.  $u - \varepsilon < s'$  (i.e.  $u$  is the smallest upper bd.)

Similarly, for infimum, we have:

Prop:  $u = \inf S$  iff

①  $s \geq u$ ,  $\forall s \in S$  (i.e.  $u$  is a lower bd.)

②  $\forall \varepsilon > 0$ ,  $\exists s' \in S$  s.t.  $u + \varepsilon > s'$  (i.e.  $u$  is the greatest lower bd.)


Q: What about the existence of infimum?

A: It follows from the completeness property.

Prop: Every  $\emptyset \neq S \subseteq \mathbb{R}$  that is bounded below must have an infimum in  $\mathbb{R}$ .

Proof: Given  $\emptyset \neq S \subseteq \mathbb{R}$ , consider the subset:

$$\emptyset \neq \bar{S} := \{-s \mid s \in S\} \in \mathbb{R}$$



Claim:  $\bar{S}$  is bdd above.

Pf of Claim:

Since  $S$  is bdd below, i.e.

$\exists$  some lower bd.  $u$  of  $S$

$$\Leftrightarrow u \leq s \quad \forall s \in S$$

$$\Rightarrow -u \geq -s \quad \forall s \in S$$

$\Rightarrow -u$  is an upper bd for  $\bar{S}$  i.e.  $\bar{S}$  is bdd above.

By Completeness Property,  $\sup \bar{S}$  exists in  $\mathbb{R}$ .

Claim:  $\inf S$  exists.  $\inf S = -\sup \bar{S}$ .

Pf of Claim:

Check:  $-\sup \bar{S}$  is a lower bd for  $S$

(Ex:)

This is the same by reversing the arguments of the claim above.

Check:  $-\sup \bar{S}$  is the greatest lower bd. for  $S$

Let  $\varepsilon > 0$  be fixed but arbitrary.

(Want to show:  $\exists s' \in S$  s.t.  $-\sup \bar{S} + \varepsilon > s'$ ) — (\*)

By ② of supremum for  $\bar{S}$ ,

$$\sup \bar{S} - \varepsilon < \bar{s}' \quad \text{for some } \bar{s}' \in \bar{S}$$

By def<sup>n</sup>, we write  $\bar{s}' = -s'$  for some  $s' \in S$

So,  $\sup \bar{S} - \varepsilon < -s' \Rightarrow -\sup \bar{S} + \varepsilon > s'$  for some  $s' \in S$   
which is (\*)

Archimedean Property:  $\mathbb{N}$  is NOT bdd above.

Pf: Suppose NOT, i.e.  $\mathbb{N}$  is bdd above.

By Completeness Property,  $\sup \mathbb{N} =: u \in \mathbb{R}$  exists.

So,  $u - 1 < n'$  for some  $n' \in \mathbb{N}$ .

$$\Rightarrow u < n' + 1 \in \mathbb{N}$$


$\Rightarrow u$  is NOT an upper bd for  $\mathbb{N}$   $\leftarrow$  **Contradiction!**

Corollaries:

(i)  $\inf \left\{ \frac{1}{n} : n \in \mathbb{N} \right\} = 0$

(ii)  $\forall \varepsilon > 0, \exists n \in \mathbb{N}$  s.t.  $0 < \frac{1}{n} < \varepsilon$

(iii)  $\forall \gamma > 0, \exists!$   $n \in \mathbb{N}$  s.t.  $n - 1 < \gamma < n$   
 $\uparrow$   
unique



Ex: Prove these!