

# Math2058 Mathematical Analysis (I)

Chi-Wai Leung

November 30, 2022

1

*Department of Mathematics,  
The Chinese University of Hong Kong, Shatin, Hong Kong  
E-mail: cwleung@math.cuhk.edu.hk*

## 1 Order structure of $\mathbb{R}$

Throughout this section, a number  $L$  means that it is a real number and let  $S$  be a non-empty subset of  $\mathbb{R}$ .

**Definition 1.1** *Using the notation as above:*

(i)  $S$  is said to be bounded above (resp. bounded below) if there is a number  $L$  (resp.  $\ell$ ) such that  $x \leq L$  (resp.  $\ell \leq x$ ) for all  $x \in S$ . In this case, such number  $L$  (resp.  $\ell$ ) is called an upper bound (resp. lower bound) for  $S$ .

Furthermore,  $S$  is said to be bounded if it is both are bounded above and bounded below.

(ii)  $S$  is said to have a maximal element (resp. minimal element) if there is an element  $M \in S$  (resp.  $m \in S$ ) such that  $x \leq M$  (resp.  $m \leq x$ ) for all  $x \in S$ . In this case, write  $\max S$  and  $\min S$  for the maximal element and the minimal element of  $S$  respectively.

**Remark 1.2** (i) It is noted that the maximum of a set may not exist even it is bounded above. For example, if let  $S = \{1 - \frac{1}{n} : n = 1, 2, \dots\}$ , then  $S$  is bounded above but  $\max S$  does not exist.

(ii) It is clear that  $\max S$  exists if and only if  $\min(-S)$  exists, where  $-S = \{-x : x \in S\}$ . In this case, we have  $-\max S = \min(-S)$ .

The following notion plays an important role in mathematics.

**Definition 1.3** *Using the notation as above, a number  $L \in \mathbb{R}$  (rep.  $\ell$ ) is called the **supremum** (resp. the **infimum**) of  $S$  if  $L$  is the least upper bound (resp. the greatest lower bound) for  $S$ . In this case, we write*

$$L := \sup S \quad ; \quad \ell := \inf S.$$

The following result is easy shown by the fact that a number  $L$  is an upper bound for  $S$  if and only if  $-L$  is a lower bound for the set  $-S$ .

---

<sup>1</sup>This is a note of the course Honours Mathematical Analysis I in 2022-23, 1<sup>st</sup> term

**Proposition 1.4** *Using the notation as above, then  $\sup S$  exists if and only if  $\inf(-S)$  exists. In this case, we have*

$$-\sup S = \inf(-S).$$

The following is a very useful result for checking a number whether it is the supremum of a given set. In addition, the technique of the proof is standard.

**Theorem 1.5** *Assume that  $\sup S$  exists. A number  $L = \sup S$  if and only if it satisfies the following two conditions.*

(i)  *$L$  is an upper bound for  $S$ .*

(ii) *For any  $\varepsilon > 0$ , there is an element  $x_0 \in S$  such that  $L - \varepsilon < x_0$ .*

*Similarly, if  $\inf S$  exists, then a number  $\ell = \inf S$  if and only if the following two conditions hold:*

(i')  *$\ell$  is a lower bound for  $S$ ;*

(ii') *For any  $\varepsilon > 0$ , there is an element  $y_0 \in S$  such that  $y_0 < \ell + \varepsilon$ .*

**Proof:** We are going to show the case of supremum first. For showing  $(\Rightarrow)$ , assume that  $L = \sup S$ . It is noted that the condition (i) automatically holds by the definition of supremum. It remains to show that the condition (ii) holds. Let  $\varepsilon > 0$ . Then  $L - \varepsilon < L$ . Since  $L$  is the least upper bound for  $S$ ,  $L - \varepsilon$  is not an upper bound for  $S$ . Therefore, there is an element  $x_0 \in S$  such that  $L - \varepsilon < x_0$  as desired.

Now for showing the converse statement, assume that the conditions (i) and (ii) hold for the number  $L$ . Then by the definition of the supremum, it needs to show that if  $L_1$  is an upper bound for  $S$ , then  $L \leq L_1$ . Suppose not, that is, we assume that there is an upper bound  $L_1$  for  $S$  such that  $L_1 < L$ . Then  $\varepsilon := 1/2(L - L_1) > 0$ . The condition (ii) gives an element  $x_0 \in S$  such that

$$L_1 < \frac{1}{2}(L_1 + L) = L - \varepsilon < x_0 \leq L.$$

The last statement can be obtained by considering  $-S$  in the first assertion above. □

**Axiom of Completeness of  $\mathbb{R}$ :** *Every bounded above non-empty subset of  $\mathbb{R}$  must have the least upper bound, that is, the supremum of a bounded above non-empty subset of  $\mathbb{R}$  must exist.*

**Proposition 1.6** *Let  $A$  and  $B$  be non-empty bounded above subsets of  $\mathbb{R}$ . Put  $A + B := \{x + y : x \in A, y \in B\}$ . Then we have  $\sup(A + B) = \sup A + \sup B$ .*

**Proof:** Note that  $L_1 := \sup A$  and  $L_2 := \sup B$  exist by the Axiom of Completeness. It is clear that  $L_1 + L_2$  is an upper bound for the set  $A + B$ . By using Theorem 1.5, it suffices to show the condition (ii) in Theorem 1.5 holds. Let  $\varepsilon > 0$ . Then by Theorem 1.5, there are elements  $a \in A$  and  $b \in B$  such that  $L_1 - \frac{1}{2}\varepsilon < a$  and  $L_2 - \frac{1}{2}\varepsilon < b$ . Hence, we have  $L_1 + L_2 - \varepsilon < a + b$ . Thus the condition (ii) holds for the set  $A + B$ . The proof is finished. □

**Proposition 1.7** *If  $S$  is a bounded below non-empty subset of  $\mathbb{R}$ , then  $\inf S$  must exist.*

**Proof:** Note that the set  $-S$  is bounded above. Then by the completeness of  $\mathbb{R}$ ,  $\sup(-S)$  exists and hence,  $\inf S = -\sup(-S)$  must exist.  $\square$

**Theorem 1.8 Archimedean Property:** *For each  $x \in \mathbb{R}$ , there is a positive integer  $n$  such that  $x < n$ .*

**Proof:** The proof is shown by the contradiction. Suppose that there is a real number  $M$  such that  $n \leq M$  for all  $n \in \mathbb{Z}_+$ . Thus, the set of all positive integers  $\mathbb{Z}_+$  is bounded above. The Axiom of Completeness tells us that the supremum  $L := \sup \mathbb{Z}_+$  must exist. Then by considering  $\varepsilon = 1$  in Theorem 1.5, there is an element  $m \in \mathbb{Z}_+$  such that  $L - 1 < m$  and hence,  $L < m + 1$ . This implies that  $n < m + 1$  for all  $n \in \mathbb{Z}_+$ . It leads to a contradiction because  $m + 1 \in \mathbb{Z}_+$ .  $\square$

**Corollary 1.9**  $\inf\{1/n : n = 1, 2, \dots\} = 0$ .

**Proof:** Let  $S := \{1/n : n = 1, 2, \dots\}$ . It is noted that 0 is a lower bound for the set  $S$ . By using Theorem 1.5, it needs to show that for any  $\varepsilon > 0$ , there is an element  $a \in S$  such that  $a < 0 + \varepsilon$ . Now let  $\varepsilon > 0$ . Then by Archimedean property, there is a positive integer  $N$  such that  $1/\varepsilon < N$ . Thus, we have  $1/N \in S$  and  $1/N < \varepsilon$  as required. The proof is finished.  $\square$

**Definition 1.10** *We say that a subset  $A$  of  $\mathbb{R}$  is dense in  $\mathbb{R}$  if  $(a, b) \cap A \neq \emptyset$ .*

**Example 1.11** *The set of all integers  $\mathbb{Z}$  is not dense in  $\mathbb{R}$ .*

The following shows that the set of rational numbers a dense subset of  $\mathbb{R}$  which  $\mathbb{Q}$  is an important dense subset.

**Proposition 1.12** *For each pair of real numbers  $a$  and  $b$  with  $a < b$ , then we have  $(a, b) \cap \mathbb{Q} \neq \emptyset$ . In this case, the set of all rational numbers is dense in  $\mathbb{R}$ .*

**Proof:** Note that we may assume  $0 < a < b$ . (Why?) By using Corollary 1.9, there is a positive integer  $N$  such that  $1/N < b - a$  and hence, we have  $1 < Nb - Na$ . On the other hand, let  $p := \max\{k \in \mathbb{N} : k \leq Na\}$ . This implies that  $Na < p + 1$ . In addition, since  $Nb - Na > 1$ , we have  $p + 1 < Nb$ . Therefore, we have  $Na < p + 1 < Nb$ . Thus,  $\frac{p+1}{N} \in \mathbb{Q} \cap (a, b)$ . The proof is finished.  $\square$

Before showing the following proposition, we have a simple but useful observation first.

**Lemma 1.13** *Let  $e, f \in \mathbb{R}$ . Then we have  $e \leq f$  if and only if for all  $\varepsilon > 0$ , we have  $e < f + \varepsilon$ .*

**Proposition 1.14** *There is a unique real number  $x$  such that  $x^2 = 2$ . Consequently, such real number is irrational.*

**Proof:** Let  $S := \{x > 0 : x^2 \leq 2\}$ . Note that  $1 \in S$ , hence,  $S \neq \emptyset$ . On the other hand, if  $x > 2$ , then  $x^2 > 4$ . This implies that the set  $S$  is bounded by 2 and thus, the set  $S$  is bounded above. Then the Axiom of Completeness assures that  $a := \sup S$  exists. We are going to show that  $a^2 = 2$  as required.

We first note that by the characterization of the sup, for each positive integer  $n$ , there is an element  $x_n \in S$  such that  $a - \frac{1}{n} < x_n$ . This implies that

$$a^2 < (x_n + \frac{1}{n})^2 = x_n^2 + \frac{2}{n}x_n + \frac{1}{n^2} < 2 + \frac{4}{n} + \frac{1}{n^2} \quad (1.1)$$

It is noted that we have  $\frac{4}{n} + \frac{1}{n^2} < \frac{5}{n}$  for all positive integer  $n$ . Therefore, we have  $\inf\{\frac{4}{n} + \frac{1}{n^2} : n = 1, 2, \dots\} = 0$  because  $\inf\{1/n : n = 1, 2, \dots\} = 0$ . This implies that for any  $\varepsilon > 0$ , there is a positive integer  $m$  such that  $\frac{4}{m} + \frac{1}{m^2} < \varepsilon$ . Therefore, we have

$$a^2 < 2 + \varepsilon$$

for all  $\varepsilon > 0$ . Lemma 1.13 implies that  $a^2 \leq 2$ .

Finally, it remains to show that  $a^2 < 2$  is impossible. Assume that  $a^2 < 2$ . Then by using the fact  $\inf\{\frac{4}{n} + \frac{1}{n^2} : n = 1, 2, \dots\} = 0$  again, one can choose a positive integer  $N$  such that  $\frac{4}{N} + \frac{1}{N^2} < 2 - a^2$ , and hence, we have  $a^2 + \frac{4}{N} + \frac{1}{N^2} < 2$ . This implies that

$$(a + 1/N)^2 = a^2 + \frac{2}{N}a + \frac{1}{N^2} \leq a^2 + \frac{4}{N} + \frac{1}{N^2} < 2.$$

Thus, we have  $(a + 1/N) \in S$  and  $a < a + 1/N$ . It leads to a contradiction. Therefore,  $a^2 = 2$ .

The uniqueness clearly follows from the fact that if  $a^2 = b^2 = 2$ , then we have  $a^2 - b^2 = (a - b)(a + b) = 0$ .

Now write  $\sqrt{2} := \sup S$ . Then by above we have  $(\sqrt{2})^2 = 2$ . Suppose that  $\sqrt{2} = p/q$  is rational, for some positive integers  $p$  and  $q$ . We have  $p^2 = 2q^2$ . Then by the Unique Prime Factorization theorem, there are natural numbers  $n$  and  $s$  such that  $p = 2^n s$  and  $s$  is not divided by 2. Similarly, there are natural numbers  $m$  and  $t$  such that  $q = 2^m t$  and  $t$  is not divided by 2. Thus, we have

$$2^{2n} s^2 = p^2 = 2q^2 = 2 \cdot 2^{2m} t^2 = 2^{2m+1} t^2.$$

From this we have  $2n = 2m + 1$ . It is impossible. The proof is complete.  $\square$

**Theorem 1.15** For any open interval  $(a, b)$ , we have  $(a, b) \cap \mathbb{Q}^c \neq \emptyset$ , i.e., the set of all irrational numbers is dense in  $\mathbb{R}$ .

**Proof:** We may assume that  $a > 0$  and hence, we have  $\sqrt{2}a < \sqrt{2}b$ . Since  $\mathbb{Q}$  is dense in  $\mathbb{R}$ , there is an element  $r \in \mathbb{Q} \cap (\sqrt{2}a, \sqrt{2}b)$ . Hence, we have

$$a < \frac{r}{\sqrt{2}} < b.$$

Since  $\sqrt{2}$  is irrational and  $r$  is rational, we see that the number  $\frac{r}{\sqrt{2}}$  is irrational as required.  $\square$

## 2 Sequences

A *sequence* of real numbers means that it is a real-valued function  $x$  defined on  $\mathbb{Z}_+$  (or  $\mathbb{N}$ ). Write  $x_n := x(n)$  for  $n = 1, 2, \dots$  and  $x = (x_n)$ .

The following definition plays a very important role in mathematics.

**Definition 2.1** We say that a sequence  $(x_n)$  is convergent if there is a number  $L \in \mathbb{R}$  which satisfies the following condition:

For any  $\varepsilon > 0$ , there is a positive integer  $N = N(\varepsilon)$  (depends on the choice of  $\varepsilon$ ), such that

$$|x_n - L| < \varepsilon \quad \text{whenever } n \geq N.$$

In this case, we say that  $(x_n)$  converges to  $L$  and  $L$  is a **limit** of  $(x_n)$ . If such  $L$  does not exist, we say that  $(x_n)$  is divergent.

**Remark 2.2** Using the notation above, we have:

- (i) A number  $\ell$  is **Not** a limit of  $(x_n)$  if there is  $\varepsilon > 0$  such that for any positive integer  $N$ , we can find a positive integer  $n$  with  $n \geq N$  so that  $|x_n - \ell| \geq \varepsilon$ .

**Warning:** in this case, it does not imply that  $(x_n)$  is divergent !!!!

- (ii) The Definition 2.1 is clearly equivalent to the following statement:  
there is a constant  $C > 0$  such that for any  $\eta > 0$ , there is a positive integer  $N$  satisfying  $|x_n - L| < C\eta$  as  $n \geq N$ .

The following is one of important properties of limits.

**Proposition 2.3** If  $(x_n)$  is a convergent sequence, then its limit is unique.

In this case, we write  $\lim x_n$  for “**the**” limit of  $(x_n)$ .

**Proof:** Let  $L$  and  $L'$  be limits of  $(x_n)$ . Then for any  $\varepsilon > 0$ , there are positive integers  $N$  and  $N'$  such that  $|x_n - L| < \varepsilon$  for any  $n \geq N$  and  $|x_n - L'| < \varepsilon$  for any  $n \geq N'$ . Now if we choose a positive  $m$  so that  $m \geq N$  and  $m \geq N'$ , then we have

$$|L - L'| \leq |L - x_m| + |x_m - L'| < 2\varepsilon.$$

Therefore, we have  $|L - L'| < 2\varepsilon$  for all  $\varepsilon > 0$ . This implies that  $|L - L'| = 0$  and thus,  $L = L'$ . Otherwise, if we choose  $0 < \varepsilon < \frac{1}{4}|L - L'|$ , then it leads to a contradiction.  $\square$

**Example 2.4** Show that if  $x_n := \frac{n+1}{n-1}$  for  $n = 2, 3, \dots$ , then the sequence  $\lim x_n = 1$ .

**Proof:** Note that for each positive integer  $n$  with  $n \geq 2$ , we have  $|x_n - 1| = \frac{2}{n-1}$ . Now let  $\varepsilon > 0$ . Therefore, we have  $|x_n - 1| < \varepsilon$  if and only if  $\frac{2}{\varepsilon} + 1 < n$ . The Archimedean property tells us that there is a positive integer  $N$  such that  $N > \frac{2}{\varepsilon} + 1$ . Hence, we have  $|x_n - 1| < \varepsilon$  as  $n \geq N$ . The proof is complete.  $\square$

**Example 2.5** Let  $x_n = (-1)^n$  for  $n = 1, 2, \dots$ . Show that the sequence  $(x_n)$  is divergent.

**Proof: Warning:** It is clear that neither 1 nor  $-1$  both is the limit of the sequence of  $(x_n)$ . However, we cannot conclude from the Definition 2.1 that the sequence  $(x_n)$  is divergent since the sequence  $(x_n)$  may converge to the number which is other than 1 and  $-1$ .

Now suppose that the sequence  $(x_n)$  is convergent with  $L := \lim x_n$ . Now if for each positive integer  $N$ , put  $A_N := \{x_n : n \geq N\}$ , then  $A_N = \{1, -1\}$ . Therefore, for any positive integer  $N$  the intersection  $(L - 1/4, L + 1/4) \cap A_N$  contains at most one point. This implies that for any positive integer  $N$ , there is  $m \geq N$  such that  $x_m \notin (L - 1/4, L + 1/4)$ , that is,  $|x_m - L| \geq 1/4$ . It leads to a contradiction since  $L$  is the limit of  $(x_n)$  by the assumption.  $\square$

**Example 2.6** Show that if  $x_n = n$  for all  $n = 1, 2, \dots$ , then the sequence  $(x_n)$  is divergent.

**Proof:** suppose not, we assume that the sequence  $(x_n)$  converges to some number  $L$ . Then by Definition 2.1, if we consider  $\varepsilon = 1$ , then there is a positive integer  $N$  such that  $|x_n - L| < 1$  for all  $n \geq N$  and thus,  $n < |L| + 1$  for all  $n \geq N$ . This implies that  $n < |L| + 1$  for all positive integers  $n$ . This contradicts to the Archimedean property.  $\square$

Using the similar idea as the proof of Example 2.6, one can obtain a more general result as follows.

**Proposition 2.7** Every convergent sequence is bounded.

**Proof:** Let  $(x_n)$  be a convergent sequence with the limit  $L$ . If we take  $\varepsilon = 1$  in the Definition 2.1, there is a positive integer  $N$  such that  $|x_n - L| < 1$  for all  $n \geq N$ . Hence, we have  $|x_n| < |L| + 1$  for all  $n \geq N$ . Thus, if we take  $M := \max\{|x_1|, \dots, |x_{N-1}|, |L| + 1\}$ , then we have  $|x_n| \leq M$  for all  $n = 1, 2, \dots$ . Thus,  $(x_n)$  is bounded.  $\square$

**Proposition 2.8** Let  $(x_n)$  and  $(y_n)$  be the convergent sequences. Let  $a := \lim x_n$  and  $b := \lim y_n$ . We have the following assertions.

- (i)  $(x_n + y_n)$  is convergent with  $\lim(x_n + y_n) = a + b$ .
- (ii) The product  $(x_n y_n)$  is convergent with  $\lim x_n y_n = ab$ .
- (iii) If  $y_n \neq 0$  for all  $n$  and  $b \neq 0$ , then the sequence  $(x_n/y_n)$  is convergent and  $\lim x_n/y_n = a/b$ .

**Proof:** For showing (i): let  $\varepsilon > 0$ . Then there is a positive integer  $N$  such that  $|x_n - a| < \varepsilon$  and  $|y_n - b| < \varepsilon$  for all  $n \geq N$ . This implies that

$$|(x_n + y_n) - (a + b)| \leq |x_n - a| + |y_n - b| < 2\varepsilon$$

for all  $n \geq N$ . Thus,  $(x_n + y_n)$  is convergent with  $\lim(x_n + y_n) = a + b$ .

For (ii), let  $\varepsilon > 0$  and let  $N$  be chosen as in Part (i). Since  $(y_n)$  is convergent,  $(y_n)$  is bounded and hence, there is  $M > 0$  such that  $|y_n| \leq M$  for all  $n$ . Hence, the triangle inequality implies that

$$|x_n y_n - ab| \leq |x_n y_n - a y_n| + |a y_n - ab| \leq |x_n - a| |y_n| + |a| |y_n - b| \leq (M + |a|) \varepsilon$$

for all  $n \geq N$ . This implies that  $(x_n y_n)$  is convergent and  $\lim x_n y_n = ab$ .

For showing (iii), it suffices to show that the sequence  $(\frac{1}{y_n})$  converges to  $1/b$  by using Part (ii).

Let  $\varepsilon > 0$  and  $N$  be as in Part (i) again. It is noted that since  $b \neq 0$ , by using the Definition 2.1 there is a positive integer  $N_1 > N$  such that  $|y_n - b| < \frac{|b|}{2}$  for all  $n \geq N_1$ . This gives  $|y_n| > \frac{|b|}{2}$  for all  $n \geq N_1$ . Hence, we have

$$\left| \frac{1}{y_n} - \frac{1}{b} \right| = \frac{|y_n - b|}{|y_n||b|} \leq \frac{2}{|b|^2} \varepsilon$$

for all  $n \geq N_1$ . The proof is complete.  $\square$

**Proposition 2.9** *Let  $(x_n)$  and  $(y_n)$  be the convergent sequences with the limits  $a := \lim x_n$  and  $b := \lim y_n$ . If  $x_n \leq y_n$  for all  $n = 1, 2, \dots$ , then  $a \leq b$ .*

**Proof:** It suffices to show that  $a < b + \varepsilon$  for all  $\varepsilon > 0$ ; otherwise, if  $b < a$ , then by taking  $\varepsilon = a - b > 0$  we have  $a < b + (a - b) = a$  which is impossible. Now let  $\varepsilon > 0$ . Then there is a positive integer  $N$  such that  $|x_N - a| < \varepsilon$  and  $|y_N - b| < \varepsilon$ . This implies that

$$a - \varepsilon < x_N \leq y_N < b + \varepsilon.$$

Thus, we have  $a < b + 2\varepsilon$ . The proof is complete.  $\square$

**Proposition 2.10** *Let  $(x_n), (y_n)$  and  $(z_n)$  be the sequences which satisfy  $x_n \leq y_n \leq z_n$  for all  $n$ . If  $a := \lim x_n = \lim z_n$ , then  $(y_n)$  is convergent and  $\lim y_n = a$ .*

**Proof:** Let  $\varepsilon > 0$ . Then by the Definition 2.1, there is a positive integer  $N$  such that  $|x_n - a| < \varepsilon$  and  $|z_n - a| < \varepsilon$  for all  $n \geq N$ . This implies that

$$a - \varepsilon < x_n \leq y_n \leq z_n < a + \varepsilon$$

for all  $n \geq N$ . Hence, we have  $|y_n - a| < \varepsilon$  for all  $n \geq N$ . The proof is finished.  $\square$

**Proposition 2.11** *let  $S$  be a non-empty bounded above subset of  $\mathbb{R}$ . Then a number  $L = \sup S$  if and only if  $L$  is an upper bound for  $S$  and there is a sequence  $(x_n)$  in  $S$  such that  $\lim x_n = L$ .*

**Proof:** For showing  $(\Rightarrow)$ , assume  $L = \sup S$ . Then  $L$  is an upper bound for  $S$  by the definition. It suffices to show that there is a sequence  $(x_n)$  in  $S$  such that  $\lim x_n = L$ . Recall the characterization of supremum that for any  $\varepsilon > 0$ , there is an element  $x \in S$  such that  $L - \varepsilon < x$ . From this for each positive integer  $n$ , there is an element  $x_n \in S$  such that  $L - \frac{1}{n} < x_n \leq L$ . This implies that  $|x_n - L| < \frac{1}{n}$  for all  $n$  and thus,  $\lim x_n = L$  as required. The converse is clear due to the characterization of supremum again.  $\square$

**Definition 2.12** *A sequence  $(x_n)$  is said to be increasing (resp. decreasing) if  $x_n \leq x_{n+1}$  (resp.  $x_n \geq x_{n+1}$  for all  $n$ ).*

**Theorem 2.13** Let  $(x_n)$  be an increasing (resp. decreasing) sequence. Then  $(x_n)$  is convergent if and only if  $(x_n)$  is bounded. In this case, we have  $\lim x_n = \sup\{x_n : n = 1, 2, \dots\}$  (resp.  $\lim x_n = \inf\{x_n : n = 1, 2, \dots\}$ ).

**Proof:** Assume that  $(x_n)$  is increasing. It is noted that this part ( $\Rightarrow$ ) is always true even  $(x_n)$  is not increasing.

Now for showing the part ( $\Leftarrow$ ), assume that  $(x_n)$  is bounded. Then the set  $S := \{x_n : n = 1, 2, \dots\}$  is bounded. The Axiom of Completeness tells us that  $L := \sup(S)$  exists. We are going to show that  $\lim x_n = L$ . In fact, for any  $\varepsilon > 0$ , there is an element  $x_N \in S$  such that  $L - \varepsilon < x_N$  because  $L = \sup(S)$ . Since  $(x_n)$  is increasing, we have  $L - \varepsilon < x_N \leq x_n \leq L$  for all  $n \geq N$ . Hence,  $|x_n - L| < \varepsilon$  for all  $n \geq N$ . Therefore,  $(x_n)$  converges to  $L$  as desired.

When  $(x_n)$  is decreasing, the assertion can be obtained by considering the sequence  $(-x_n)$ .  $\square$

**Example 2.14** Then the following limit exists

$$e := \lim\left(1 + \frac{1}{n}\right)^n.$$

**Proof:** For each positive integer, let

$$x_n = \left(1 + \frac{1}{n}\right)^n.$$

They by using Proposition 2.13, it suffices to show that  $(x_n)$  is a bounded increasing sequence.

We first claim that  $(x_n)$  is increasing. In fact, by the Binomial Theorem, we see that

$$x_n = 1 + 1 + \sum_{k=2}^n \frac{n(n-1)\cdots(n-k+1)}{k!} \frac{1}{n^k} = \sum_{k=1}^n \frac{1}{k!} \left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)\cdots\left(1 - \frac{k-1}{n}\right). \quad (2.1)$$

It is noted that each term in above is positive and the coefficients of  $\frac{1}{k!}$  for  $2 \leq k \leq n$  in  $x_n$  and  $x_{n+1}$  are

$$\left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)\cdots\left(1 - \frac{k-1}{n}\right) \quad \text{and} \quad \left(1 - \frac{1}{n+1}\right)\left(1 - \frac{2}{n+1}\right)\cdots\left(1 - \frac{k-1}{n+1}\right)$$

respectively. From this we see that  $x_n \leq x_{n+1}$  for all  $n$  and thus, the sequence  $(x_n)$  is increasing.

It remains to show that  $(x_n)$  is bounded. In fact, for each  $2 \leq k \leq n$  we have

$$\frac{1}{k!} \left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)\cdots\left(1 - \frac{k-1}{n}\right) < \frac{1}{2^k}.$$

Then

$$x_n < 1 + 1 + \sum_{k=1}^n \frac{1}{2^k} < 3.$$

The proof is complete.  $\square$

**Remark 2.15** The limit  $e$  in the Example 2.14 above is very important in mathematics which is called the natural base today. It was first induced by Euler. In fact,  $x_n := \left(1 + \frac{1}{n}\right)^n$  is motivated by the Compound interest formula.



**Theorem 2.16 Nested Intervals Theorem** Let  $(I_n := [a_n, b_n])$  be a sequence of closed and bounded intervals. Assume that  $I_1 \supseteq I_2 \supseteq I_3 \supseteq \dots$ . Then we have  $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ . Furthermore, if we further assume that  $\lim_n(b_n - a_n) = 0$ , then there is a unique real number  $c$  such that  $\bigcap_{n=1}^{\infty} I_n = \{c\}$ .

**Proof:** It is noted that since  $(I_n)$  is a decreasing sequence of closed and bounded intervals, we have

$$a_1 \leq a_2 \leq \dots \leq a_n < b_n \leq b_{n-1} \leq \dots \leq b_2 \leq b_1$$

for all positive integers  $n$ . Therefore,  $(a_n)$  and  $(b_n)$  are bounded and they are increasing and decreasing and respectively. This implies that  $(x_n)$  and  $(y_n)$  both are convergent and  $a := \lim a_n = \sup\{a_n : n = 1, 2, \dots\}$  and  $b := \lim b_n = \inf\{b_n : n = 1, 2, \dots\}$ . In addition, we have  $a \leq b$  because  $a_n \leq b_n$  for all  $n$ . Thus, if we fix some  $c$  such that  $a \leq c \leq b$ , then  $c \in \bigcap_{n=1}^{\infty} I_n$  as desired because we have  $a_n \leq a \leq c \leq b \leq b_n$  for all  $n$ .

It remains to show  $\bigcap_{n=1}^{\infty} I_n = \{c\}$  if  $\lim(b_n - a_n) = 0$ . In fact, if  $c' \in \bigcap_{n=1}^{\infty} I_n$ , then we have  $|c - c'| \leq |b_n - a_n|$  for all  $n$ . This implies that  $|c - c'| = 0$  and thus,  $c = c'$ . The proof is finished.  $\square$

**Remark 2.17** The assumption of the boundedness and closeness of the intervals  $I_n$  cannot be removed in the Nest Intervals Theorem.

For example, if  $I_n := (0, \frac{1}{n})$  and  $J_n := [n, \infty)$ , for all  $n = 1, 2, \dots$ , then  $\bigcap I_n = \bigcap J_n = \emptyset$ .

### 3 Subsequences

**Definition 3.1** A subsequence  $(x_{n_k})_{k=1}^{\infty}$  of a sequence  $(x_n)$  means that  $(n_k)_{k=1}^{\infty}$  is a sequence of positive integers satisfying  $n_1 < n_2 < \dots < n_k < n_{k+1} < \dots$ , that is, such sequence  $(n_k)$  can be viewed as a strictly increasing function  $\mathbf{n} : k \in \{1, 2, \dots\} \mapsto n_k \in \{1, 2, \dots\}$ .

**Remark 3.2** In this case, note that for each positive integer  $N$ , there is  $K \in \mathbb{N}$  such that  $n_K \geq N$  and thus we have  $n_k \geq N$  for all  $k \geq K$ .

**Proposition 3.3** If  $(x_n)$  is a convergent sequence, then any subsequence  $(x_{n_k})$  of  $(x_n)$  converges to the same limit. In this case, we have  $\lim_k x_{n_k} = \lim x_n$ .

**Proof:** We assume that  $\lim x_n = a \in \mathbb{R}$  exists. Let  $(x_{n_k})$  be a subsequence of  $(x_n)$ . We claim that  $\lim x_{n_k} = a$ . Let  $\varepsilon > 0$ . In fact, since  $\lim x_n = a$ , there is a positive integer  $N$  such that  $|a - x_n| < \varepsilon$  for all  $n \geq N$ . Note that by the definition of a subsequence, there is a positive integer  $K$  such that  $n_k \geq N$  for all  $k \geq K$ . Hence, we see that  $|a - x_{n_k}| < \varepsilon$  for all  $k \geq K$ . Thus we have  $\lim_{k \rightarrow \infty} x_{n_k} = a$ . The proof is complete.  $\square$

**Theorem 3.4 Bolzano-Weierstrass Theorem** (write B-W Theorem for short):  
Every bounded sequence has a convergent subsequence.

**Proof:** We give two different proofs in here, however, each proof basically is due to the Axiom of Completeness.

Let  $(x_n)$  be a bounded sequence and put  $X := \{x_n : n = 1, 2, \dots\}$ . The Theorem clearly holds if  $X$  is a finite set. In fact in this case, there must have an element  $x_m$  appears infinite many times. Hence, we can choose a subsequence  $(x_{n_k})$  so that  $x_{n_k} \equiv x_m$  for all  $k = 1, 2, \dots$

Thus we may assume that the set  $X$  is infinite.

**Method 1:**

Since  $(x_n)$  is bounded, there is a closed and bounded interval  $I_1 = [a_1, b_1]$  such that  $x_n \in I_1$  for all  $n$ . Put  $x_{n_1} := x_1$ .

It is noted that one of the following sets must be infinite:

$$A_2 := \{n \in \mathbb{Z}_+ : x_n \in [a_1, \frac{a_1 + b_1}{2}]\}; \quad B_2 := \{n \in \mathbb{Z}_+ : x_n \in [\frac{a_1 + b_1}{2}, b_1]\}.$$

We may assume that the set  $A_2$  is infinite. Hence there is an element  $n_2 \in A_2$  such that  $n_1 < n_2$ . Put  $I_2 := [a_2, b_2] = [a_1, \frac{a_1 + b_1}{2}]$ . Thus  $x_{n_2} \in I_2$ . Similarly, one of the following sets is infinite:

$$A_3 := \{n \in \mathbb{Z}_+ : x_n \in [a_2, \frac{a_2 + b_2}{2}]\}; \quad B_3 := \{n \in \mathbb{Z}_+ : x_n \in [\frac{a_2 + b_2}{2}, b_2]\}.$$

In addition, we may assume that the set  $A_3$  is infinite. Hence, there is an element  $n_3 \in A_3$  such that  $n_1 < n_2 < n_3$ . Put  $I_3 := [a_3, b_3] = [a_2, \frac{a_2 + b_2}{2}]$ . Thus,  $x_{n_3} \in I_3$ . By repeating the same step, we can get a decreasing sequence of a closed and bounded intervals  $I_k = [a_k, b_k]$  and a subsequence  $(x_{n_k})$  of  $(x_n)$  such that the following conditions hold:

1.  $\lim(b_k - a_k) = \lim \frac{1}{2^k}(b_1 - a_1) = 0$ .
2.  $x_{n_k} \in I_k$  for all  $k = 1, 2, \dots$

The Nest Intervals Theorem tells us that there is a number  $c$  such that  $c \in I_k$  for all  $k$  and hence, we have  $|x_{n_k} - c| \leq (b_k - a_k) = \frac{1}{2^k}(b_1 - a_1) \rightarrow 0$ . Therefore the subsequence  $(x_{n_k})$  is convergent as required. The proof is finished.

**Method 2**

This method is the Weierstrass' original proof.

Recall our assumption that the set  $X = \{x_n : n = 1, 2, \dots\}$  is infinite. Let

$$S := \{x \in \mathbb{R} : (x, \infty) \cap X \text{ is infinite}\}.$$

We first note that since  $(x_n)$  is bounded, there are real numbers  $m$  and  $M$  so that  $m \leq x_n \leq M$  for all  $n$ . Since the set  $X = \{x_n : n = 1, 2, \dots\}$  is infinite, the set  $S$  is a bounded above non-empty set because  $m \in S$  and  $x \leq M$  for all  $x \in S$ . The Axiom of Completeness implies that  $L := \sup(S)$  must exist. We want to show that there is a subsequence  $(x_{n_k})$  of  $(x_n)$  which converges to  $L$ .

**Claim:** For any  $\varepsilon > 0$ , there is an element  $u \in S$  such that  $|u - L| < \varepsilon$  and  $(u, L + \varepsilon] \cap X$  is infinite.

In fact, if let  $\varepsilon > 0$ , then by the characterization of the supremum there is an element  $u \in S$  such that  $L - \varepsilon < u$ . Since  $u \in S$ , we have  $(u, \infty) \cap X$  is infinite. It implies that the set  $(u, L + \varepsilon] \cap X$  must be infinite, otherwise,  $(L + \varepsilon, \infty) \cap X$  is infinite and thus,  $L + \varepsilon \in S$  by the construction of  $S$ . It leads to a contradiction because  $L$  is an upper bound for  $S$ . Thus, the **Claim** follows.

Now for  $\varepsilon = 1$ , then there is  $u_1 \in S$  such that  $L - 1 < u_1 < L + 1$ . Then by the Claim above, choose  $x_{n_1} \in (u_1, L + 1]$  and hence,  $L - 1 < x_{n_1} \leq L + 1$ . Next, we considering  $\varepsilon = 1/2$ , then there is an element  $u_2 \in S$  such that the set  $(u_2, L + 1/2]$  is infinite by the Claim above again. Therefore we can find  $x_{n_2}$  such that  $n_1 < n_2$  and  $L - 1/2 < u_2 < x_{n_2} \leq L + 1/2$ . By repeating the same step and considering  $\varepsilon = \frac{1}{k}$  for  $k = 1, 2, \dots$  in the **Claim** above, we can get a subsequence  $(x_{n_k})$  of  $(x_n)$  such that  $L - \frac{1}{k} < x_{n_k} \leq L + \frac{1}{k}$  for all  $k = 1, 2, \dots$ . Therefore,  $(x_{n_k})$  is a convergent subsequence of  $(x_n)$  with the limit  $L$ . The proof is complete.  $\square$

**Remark 3.5** The assumption of the boundedness of  $(x_n)$  cannot be removed. For example, let  $x_n = n$  for all  $n = 1, 2, \dots$ . Then  $(x_n)$  does not have a convergent subsequence because  $|x_n - x_m| \geq 1$  for  $n \neq m$ .

**Proposition 3.6** Let  $(x_n)$  be a bounded sequence. For each positive integer  $n$ , put

$$a_n := \inf\{x_k : k \geq n\} \quad \text{and} \quad b_n := \sup\{x_k : k \geq n\}.$$

Then we have the following assertions.

- (i) The limits  $\lim a_n$  and  $\lim b_n$  always exist with  $\lim a_n \leq \lim b_n$ . In this case, we write  $\underline{\lim} x_n := \lim a_n$  (called the *lim inf* of  $(x_n)$ ) and  $\overline{\lim} x_n = \lim b_n$  (called the *lim sup* of  $(x_n)$ ).
- (ii)  $(x_n)$  is convergent if and only if  $\underline{\lim} x_n = \overline{\lim} x_n$ . In this case, we have  $\lim x_n = \underline{\lim} x_n = \overline{\lim} x_n$ .
- (iii) There exists a subsequence  $(x_{n_k})$  of  $(x_n)$  such that  $\lim x_{n_k} = \overline{\lim} x_n$ . Consequently, the Bolzano-Weierstrass Theorem holds.

**Proof:** For showing part (i), we note that if  $a_n \leq x_k$  for all  $k \geq n$ , then  $a_n \leq x_k$  for  $k \geq n+1$ . Thus, we have  $a_n \leq a_{n+1}$  for all  $n$ . Similarly, we have  $b_{n+1} \geq b_n$ . Thus, we have  $a_1 \leq \dots \leq a_n \leq a_{n+1} \leq b_{n+1} \leq b_n \leq \dots \leq b_1$  for all  $n$ . This implies that  $(a_n)$  and  $(b_n)$  both are bounded monotone sequences. Therefore,  $\lim a_n$  and  $\lim b_n$  both exist. In fact, we have

$$\underline{\lim} x_n = \sup_n \inf_{k \geq n} x_k \leq \overline{\lim} x_n = \inf_n \sup_{k \geq n} x_k.$$

For part (ii), we first assume that  $l := \lim x_n$  exists. Thus, for any  $\varepsilon > 0$ , there is a positive integer  $N$  such that  $l - \varepsilon < x_n < l + \varepsilon$  for all  $n \geq N$ . Then by the definition of  $a_n$  and  $b_n$ , we have

$$l - \varepsilon \leq a_n \leq b_n \leq l + \varepsilon$$

for all  $n \geq N$ . Thus, we have  $|b_n - a_n| \leq 2\varepsilon$  for all  $n \geq N$ . By taking  $n \rightarrow \infty$ , this gives  $|\overline{\lim} x_n - \underline{\lim} x_n| \leq 2\varepsilon$  for all  $\varepsilon > 0$ , and hence, we have  $\overline{\lim} x_n = \underline{\lim} x_n$ .

Now for showing the converse ( $\Leftarrow$ ), we assume that we have  $l := \overline{\lim} x_n = \underline{\lim} x_n$ . Then for any  $\varepsilon$ , there is a positive integer  $N$  so that  $l - \varepsilon < a_n \leq b_n < l + \varepsilon$  for all  $n \geq N$ . Since we always have  $a_n \leq x_k \leq b_n$  for all  $k \geq n$ . Therefore, we have  $l - \varepsilon < x_k < l + \varepsilon$  for all  $k \geq N$  and hence,  $\lim x_k = l$ .

For proving part (iii), we are going to construct a subsequence  $(x_{n_k})$  of  $(x_n)$  so that  $\lim x_{n_k} = \overline{\lim} x_n$ . Let  $L := \overline{\lim} x_n$ . It is noted that for any  $\varepsilon > 0$ , there is a positive integer  $N$  so that  $L - \varepsilon < b_n := \sup_{k \geq n} x_k < L + \varepsilon$  for all  $n \geq N$ . This implies that  $x_k < L + \varepsilon$  for all  $k \geq N$ .

If we fix  $n \geq N$ , since  $L - \varepsilon < b_n$  for all  $n \geq N$ , we can choose  $\eta > 0$  such that  $L - \varepsilon < b_n - \eta$ . Using the characterization of surpeming, we have  $L - \varepsilon < b_n - \eta < x_m$  for some  $m \geq n$ . Therefore, we have shown that

$$\forall \varepsilon > 0, \exists N, \forall n \geq N, \exists m \geq n \quad \text{so that} \quad L - \varepsilon < x_m < L + \varepsilon. \quad (3.1)$$

Now for considering  $\varepsilon = 1$  in 3.1, there is  $N_1$  so that  $L - 1 < x_{n_1} < L + 1$  for some  $n_1 \geq N_1$ . Next, for considering  $\varepsilon = 1/2$  in 3.1, there is  $N_2$  so that for any  $n \geq N_2$ , we have  $L - 1/2 < x_m < L + 1/2$  for some  $m \geq n$ . Thus, if we choose  $n > N_2$  and  $n > n_1$ , then there is  $n_2 \geq n$  so that  $n_2 > n_1$  and  $L - 1/2 < x_{n_2} < L + 1/2$ .

Similarly, if we take  $\varepsilon = 1/3$ , there is a positive integer  $N_3$  so that for any  $n \geq N_3$  we have  $L - 1/3 < x_m < L + 1/3$  for some  $m \geq n$ . Therefore, if we take  $n > N_3$  and  $n > n_2$ , then there is  $n_3 \geq n$  such that  $L - 1/3 < x_{n_3} < L + 1/3$  and  $n_3 > n_2$ .

To repeat the same steps, we get a strictly increasing sequence of positive integers  $(n_k)$  so that  $L - 1/k < x_{n_k} < L + 1/k$  for all  $k$ . Thus,  $(x_{n_k})$  is a convergent subsequence with the limit  $L$ . The proof is complete.  $\square$

**Proposition 3.7** *Let  $(x_n)$  and  $(y_n)$  be bounded sequences. Then we have*

$$(i) \overline{\lim}(-a_n) = -\underline{\lim}a_n.$$

$$(ii) \overline{\lim}(ax_n) = a\overline{\lim}x_n \text{ for } a \geq 0.$$

$$(iii) \underline{\lim}x_n + \underline{\lim}y_n \leq \underline{\lim}(x_n + y_n) \leq \overline{\lim}(x_n + y_n) \leq \overline{\lim}x_n + \overline{\lim}y_n$$

**Proof:** Parts (i) and (ii) are clear. We want to show part (iii) and claim that

$$\overline{\lim}(x_n + y_n) \leq \overline{\lim}x_n + \overline{\lim}y_n.$$

Let  $b := \overline{\lim}x_n$  and  $c := \overline{\lim}y_n$ . Let  $\varepsilon > 0$ . Then there is a positive integer  $N$  such that  $b_n < b + \varepsilon$  and  $c_n < c + \varepsilon$  for all  $n \geq N$ . This implies that  $x_k + y_k \leq b_n + c_n < b + c + 2\varepsilon$  for all  $k \geq n \geq N$ . Therefore, we have  $\sup_{k \geq n}(x_k + y_k) < b + c + 2\varepsilon$  for all  $n \geq N$  and thus,  $\overline{\lim}(x_n + y_n) = \lim_n \sup_{k \geq n}(x_k + y_k) < b + c + 2\varepsilon$  for all  $\varepsilon > 0$ . This gives  $\overline{\lim}(x_n + y_n) = \lim_n \sup_{k \geq n}(x_k + y_k) < b + c$  as desired.

By considering the sequences  $(-x_n)$  and  $(-y_n)$  in above, we see that  $\underline{\lim}x_n + \underline{\lim}y_n \leq \underline{\lim}(x_n + y_n)$ . the proof is complete.  $\square$

**Remark 3.8** It is noted that in general we don't have the equality  $\overline{\lim}(x_n + y_n) = \overline{\lim}x_n + \overline{\lim}y_n$ . For example, if we let  $x_n = (-1)^{n+1}$  and  $y_n = (-1)^n$ , then  $\overline{\lim}(x_n + y_n) < \overline{\lim}x_n + \overline{\lim}y_n$ .

## 4 Compact Sets

Motivated by the Bolzano-Weierstrass Theorem, the following notation plays a very important role in Mathematics.

**Definition 4.1** *A subset  $A$  of  $\mathbb{R}$  is said to be compact if for any sequence  $(x_n)$  in  $A$ , there is a convergent subsequence  $(x_{n_k})$  of  $(x_n)$  such that  $\lim x_{n_k} \in A$ .*

**Example 4.2** *Clearly,  $\mathbb{R}$  and  $(0, 1)$  are not compact.*

**Proposition 4.3** *Every closed and bounded interval is compact.*

**Proof:** Recall a closed and bounded interval that it is a set  $[a, b] := \{x : a \leq x \leq b\}$  for some  $-\infty < a < b < \infty$ .

Let  $(x_n)$  be a sequence in  $[a, b]$ . Then  $(x_n)$  is a bounded sequence. The Bolzano-Weierstrass Theorem gives a convergent subsequence  $(x_{n_k})$ . It is noted since  $a \leq x_{n_k} \leq b$  for all  $k = 1, 2, \dots$ , we have  $a \leq \lim x_{n_k} \leq b$ . Thus,  $\lim x_{n_k} \in [a, b]$  as desired.  $\square$

**Remark 4.4** However, a compact set need not be a closed and bounded interval. For example,  $[0, 1] \cup \{2\}$  is a compact set but it is not an interval.

In the remainder of this section, we give a characterization of a compact set.

**Definition 4.5** Let  $A$  be a subset of  $\mathbb{R}$ . A point  $x$  is called a *limit point* (or *cluster point*) of  $A$  if for any  $\varepsilon > 0$ , there is an element  $a \in A$  such that  $0 < |x - a| < \varepsilon$ , i.e., there is an element  $a \in A$  with  $x \neq a$  such that  $|x - a| < \varepsilon$ . We write  $D(A)$  for the set of all limit points of  $A$ .

Furthermore,  $A$  is said to be *closed* if  $D(A) \subseteq A$ .

**Example 4.6**

- (i) If  $A = (0, 1] \cup \{2\}$ , then  $D(A) = [0, 1]$ . Hence,  $A$  is not closed since  $0 \in D(A) \setminus A$ .
- (ii) If  $A = \mathbb{Z}$ , then  $D(A) = \emptyset$  and thus,  $\mathbb{Z}$  is a closed set.

**Proposition 4.7** Let  $A$  be a subset of  $\mathbb{R}$ . Then the following statements are equivalent.

- (i)  $A$  is closed.
- (ii) If  $(x_n)$  is a sequence in  $A$  and is convergent, then  $\lim x_n \in A$ .

**Proof:** For  $(i) \Rightarrow (ii)$ , assume that  $A$  is closed but the condition  $(ii)$  does not hold. Then there is a convergent sequence  $(x_n)$  in  $A$  but the limit  $l := \lim x_n \notin A$ . Since  $A$  is closed,  $D(A) \subseteq A$ . Thus,  $l$  is not a limit point of  $A$ . This implies that there is  $\delta > 0$  so that  $((l - \delta, l + \delta) \setminus \{l\}) \cap A = \emptyset$ . Since  $\lim x_n = l$ , there is a positive integer  $N$  such that  $|x_N - l| < \delta$ . Note that we have  $l \neq x_N$  because  $l \notin A$ . Hence,  $x_N \in ((l - \delta, l + \delta) \setminus \{l\}) \cap A$  which leads to a contradiction. Therefore,  $(ii)$  holds.

For  $(ii) \Rightarrow (i)$ , let  $z \in D(A)$ . Then for any  $\varepsilon > 0$ , there is an element  $x \in A$  such that  $0 < |x - z| < \varepsilon$ . Therefore, for each positive integer  $n$ , there is an element  $x_n \in A$  such that  $0 < |x_n - z| < 1/n$  and thus,  $z := \lim x_n$ . The assumption  $(i)$  implies that  $z \in A$ . Therefore,  $D(A) \subseteq A$ . The proof is complete.  $\square$

**Definition 4.8** For a subset  $A$  of  $\mathbb{R}$ , put

$$\bar{A} = A \cup D(A).$$

The set  $\bar{A}$  is called the *closure* of  $A$ .

**Example 4.9** We have the following examples.

1.  $\overline{(0, 1]} = [0, 1]$ .
2.  $\overline{\mathbb{Q}} = \mathbb{R}$ .
3.  $\overline{\mathbb{Z}} = \mathbb{Z}$ .

**Proposition 4.10** Let  $A$  be a subset of  $\mathbb{R}$ . Then we have the following assertions.

1.  $\bar{A}$  is closed.
2.  $A$  is closed if and only if  $\bar{A} = A$ .
3.  $z \in \bar{A}$  if and only if for any  $\delta > 0$ , there is an element  $a \in A$  so that  $|z - a| < \delta$  if and only if there is a convergent sequence  $(x_n)$  in  $A$  so that  $z = \lim x_n$ .
4.  $\bar{A}$  is the smallest closed set containing  $A$ , i.e., if  $B$  is a closed set containing  $A$ , then  $\bar{A} \subseteq B$ .

**Proof:** For showing part (1), we need to show that  $D(\bar{A}) \subseteq \bar{A}$ . Suppose not, assume that there is an element  $z \in D(\bar{A})$  but  $z \notin \bar{A}$ . Since  $z \notin \bar{A}$ , there is  $\delta > 0$  such that  $(z - \delta, z + \delta) \cap A = \emptyset$ . On the other hand, there is an element  $b \in (z - \delta, z + \delta) \cap \bar{A}$  because  $z \in D(\bar{A})$ . Now choose  $r > 0$  such that  $(b - r, b + r) \subseteq (z - \delta, z + \delta)$ . Using the definition of limit points again, we can find some element  $a \in A$  such that  $a \in (b - r, b + r)$  and thus,  $a \in (z - \delta, z + \delta) \cap A$ . It leads to a contradiction because  $(z - \delta, z + \delta) \cap A = \emptyset$  by the choice of  $\delta$ .

Parts (2)-(4) can be shown by the definition of limit points directly. Try to do it by yourself.  $\square$

Recall that a subset  $A$  of  $\mathbb{R}$  is said to be dense in  $\mathbb{R}$  if for any open interval  $I$ , we have  $I \cap A \neq \emptyset$ .

**Proposition 4.11** *Let  $A$  be a subset of  $\mathbb{R}$ . Then  $A$  is dense in  $\mathbb{R}$  if and only if  $\bar{A} = \mathbb{R}$ .*

**Proof:** For showing  $(\Rightarrow)$ : assume that  $A$  is a dense set. Let  $z \in \mathbb{R}$ . Then for any  $\delta > 0$ , we have  $(z - \delta, z + \delta) \cap A \neq \emptyset$  by the definition of a dense set. Hence, there is  $a \in A$  such that  $|z - a| < \delta$ . Thus,  $z \in \bar{A}$  by Proposition 4.10(3) above.

Conversely, assume that  $\bar{A} = \mathbb{R}$ . Let  $I$  be an open interval. We want to show  $I \cap A$  is non-empty. Fix an element  $z \in I$ . Since  $I$  is an open interval, we can choose  $\delta > 0$  such that  $(z - \delta, z + \delta) \subseteq I$ . Since  $\bar{A} = \mathbb{R}$ , by using Proposition 4.10(3) again, there is an element  $a \in A$  such that  $|z - a| < \delta$ . Therefore,  $a \in (z - \delta, z + \delta) \cap A$  and hence,  $I \cap A \neq \emptyset$ . The proof is finished.  $\square$

**Theorem 4.12** *Let  $A$  be a subset of  $\mathbb{R}$ . Then  $A$  is compact if and only if  $A$  is a closed and bounded subset.*

**Proof:** For showing the necessary part, we assume that  $A$  is compact.

We first claim that  $A$  is bounded. Suppose that  $A$  is unbounded. If we fix an element  $x_1 \in A$ , then there is  $x_2 \in A$  such that  $|x_1 - x_2| > 1$ . Using the unboundedness of  $A$ , we can find an element  $x_3 \in A$  such that  $|x_3 - x_k| > 1$  for  $k = 1, 2$ . To repeat the same step, we can find a sequence  $(x_n)$  in  $A$  such that  $|x_n - x_m| > 1$  for  $n \neq m$ . Thus  $A$  has no convergent subsequence. Thus  $A$  must be bounded.

Finally, we show that  $A$  is closed. Let  $(x_n)$  be a sequence in  $A$  and it is convergent. It needs to show that  $\lim_n x_n \in A$ . Note that since  $A$  is compact,  $(x_n)$  has a convergent subsequence  $(x_{n_k})$  such that  $\lim_k x_{n_k} \in A$ . Then by Proposition 3.3, we see that  $\lim_n x_n = \lim_k x_{n_k} \in A$ . The proof is finished.

Conversely, we suppose that  $A$  is closed and bounded. Let  $(x_n)$  be a sequence in  $A$  and thus  $(x_n)$  is a bounded sequence in  $\mathbb{R}$ . Then by the Bolzano-Weierstrass Theorem,  $(x_n)$  has a convergent subsequence  $(x_{n_k})$ . Since  $A$  is closed,  $\lim_k x_{n_k} \in A$ . Therefore,  $A$  is compact.

□

**Example 4.13** Let  $A = \{1/n : n = 1, 2, \dots\} \cup \{0\}$ . Then  $A$  is a compact set.  $A$  is clearly bounded. Then by Theorem 4.12, it suffices to show that the set  $A$  is closed. Clearly,  $0 \in D(A)$ . We are going to show  $D(A) = \{0\}$ . In fact, if  $z \neq 0$ , clearly we can find some  $r > 0$  such that the intersection  $(z - r, z + r) \cap A$  contains at most one point. Therefore, if  $z \neq 0$ , then  $z \notin D(A)$ . Thus,  $D(A) = \{0\}$ . Hence, the set  $A$  is closed as desired.

In the rest of this section, we are going to use another description of a compact set.

For convenience, we call a collection of open intervals  $\{J_\alpha : \alpha \in \Lambda\}$  an *open intervals cover* of a given subset  $A$  of  $\mathbb{R}$ , where  $\Lambda$  is an arbitrary non-empty index set, if each  $J_\alpha$  is an open interval (not necessary bounded) and

$$A \subseteq \bigcup_{\alpha \in \Lambda} J_\alpha.$$

**Theorem 4.14 Heine-Borel Theorem:** Any closed and bounded interval  $[a, b]$  satisfies the following condition which is called *the Heine-Borel Property*.

(HB) Given any open intervals cover  $\{J_\alpha\}_{\alpha \in \Lambda}$  of  $[a, b]$ , there are finitely many  $J_{\alpha_1}, \dots, J_{\alpha_N}$  such that  $[a, b] \subseteq J_{\alpha_1} \cup \dots \cup J_{\alpha_N}$

**Proof:** Suppose that  $[a, b]$  does not satisfy Heine-Borel Property. Then there is an open intervals cover  $\{J_\alpha\}_{\alpha \in \Lambda}$  of  $[a, b]$  but it has no finite sub-cover. Let  $I_1 := [a_1, b_1] = [a, b]$  and  $m_1$  the mid-point of  $[a_1, b_1]$ . Then by the assumption,  $[a_1, m_1]$  or  $[m_1, b_1]$  cannot be covered by finitely many  $J_\alpha$ 's. We may assume that  $[a_1, m_1]$  cannot be covered by finitely many  $J_\alpha$ 's. Put  $I_2 := [a_2, b_2] = [a_1, m_1]$ . To repeat the same steps, we can obtain a sequence of closed and bounded intervals  $I_n = [a_n, b_n]$  with the following properties:

- (a)  $I_1 \supseteq I_2 \supseteq I_3 \supseteq \dots$ ;
- (b)  $\lim_n (b_n - a_n) = 0$ ;
- (c) each  $I_n$  cannot be covered by finitely many  $J_\alpha$ 's.

Then by the Nested Intervals Theorem, there is an element  $\xi \in \bigcap_n I_n$  such that  $\lim_n a_n = \lim_n b_n = \xi$ . In particular, we have  $a = a_1 \leq \xi \leq b_1 = b$ . Hence, there is  $\alpha_0 \in \Lambda$  such that  $\xi \in J_{\alpha_0}$ . Since  $J_{\alpha_0}$  is open, there is  $\varepsilon > 0$  such that  $(\xi - \varepsilon, \xi + \varepsilon) \subseteq J_{\alpha_0}$ . On the other hand, there is  $N \in \mathbb{N}$  such that  $a_N$  and  $b_N$  in  $(\xi - \varepsilon, \xi + \varepsilon)$  because  $\lim_n a_n = \lim_n b_n = \xi$ . Thus we have  $I_N = [a_N, b_N] \subseteq (\xi - \varepsilon, \xi + \varepsilon) \subseteq J_{\alpha_0}$ . It contradicts to the Property (c) above. The proof is complete.

□

**Remark 4.15** The assumption of the closeness and boundedness of an interval in Heine-Borel Theorem is essential.

For example, notice that  $\{J_n := (1/n, 1) : n = 1, 2, \dots\}$  is an open interval covers of  $(0, 1)$  but you cannot find finitely many  $J_n$ 's to cover the open interval  $(0, 1)$ .

**Lemma 4.16** A subset  $A$  is a closed subset of  $\mathbb{R}$  if and only if for each element  $x \notin A$ , there is  $r > 0$  such that  $(x - r, x + r) \cap A = \emptyset$ .

The following is a very important feature of a compact set.

**Theorem 4.17** Let  $A$  be a subset of  $\mathbb{R}$ . Then the following statements are equivalent.

(i) For any open intervals cover  $\{J_\alpha\}_{\alpha \in \Lambda}$  of  $A$ , we can find finitely many  $J_{\alpha_1}, \dots, J_{\alpha_N}$  such that  $A \subseteq J_{\alpha_1} \cup \dots \cup J_{\alpha_N}$ .

(ii)  $A$  is compact.

(iii)  $A$  is closed and bounded.

**Proof:** The result will be shown by the following path

$$(i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i).$$

For  $(i) \Rightarrow (ii)$ , assume that the condition (i) holds but  $A$  is not compact. Then there is a sequence  $(x_n)$  in  $A$  such that  $(x_n)$  has no subsequence which has the limit in  $A$ . Put  $X = \{x_n : n = 1, 2, \dots\}$ . Then  $X$  is infinite. Note that for each element  $a \in A$ , there is  $\delta_a > 0$  such that  $J_a := (a - \delta_a, a + \delta_a) \cap X$  is finite. Indeed, if there is an element  $a \in A$  such that  $(a - \delta, a + \delta) \cap X$  is infinite for all  $\delta > 0$ , then  $(x_n)$  has a convergent subsequence with the limit  $a$ . On the other hand, we have  $A \subseteq \bigcup_{a \in A} J_a$ . Then by the compactness of  $A$ , we can find finitely many  $a_1, \dots, a_N$  such that  $A \subseteq J_{a_1} \cup \dots \cup J_{a_N}$ . Hence, we have  $X \subseteq J_{a_1} \cup \dots \cup J_{a_N}$ . Then by the choice of  $J_a$ 's,  $X$  must be finite. This leads to a contradiction. Therefore,  $A$  must be compact.

The implication  $(ii) \Rightarrow (iii)$  follows immediately from Theorem 4.12.

Finally we want to show  $(iii) \Rightarrow (i)$ . Suppose that  $A$  is closed and bounded. Then we can find a closed and bounded interval  $[a, b]$  such that  $A \subseteq [a, b]$ . Now let  $\{J_\alpha\}_{\alpha \in \Lambda}$  be an open intervals cover of  $A$ . Note that for each element  $x \in [a, b] \setminus A$ , there is  $\delta_x > 0$  such that  $(x - \delta_x, x + \delta_x) \cap A = \emptyset$  since  $A$  is closed by using Lemma 4.16. If we put  $I_x = (x - \delta_x, x + \delta_x)$  for  $x \in [a, b] \setminus A$ , then we have

$$[a, b] \subseteq \bigcup_{\alpha \in \Lambda} J_\alpha \cup \bigcup_{x \in [a, b] \setminus A} I_x.$$

Using the Heine-Borel Theorem 4.14, we can find finitely many  $J_\alpha$ 's and  $I_x$ 's, say  $J_{\alpha_1}, \dots, J_{\alpha_N}$  and  $I_{x_1}, \dots, I_{x_K}$ , such that  $A \subseteq [a, b] \subseteq J_{\alpha_1} \cup \dots \cup J_{\alpha_N} \cup I_{x_1} \cup \dots \cup I_{x_K}$ . Note that  $I_x \cap A = \emptyset$  for each  $x \in [a, b] \setminus A$  by the choice of  $I_x$ . Therefore, we have  $A \subseteq J_{\alpha_1} \cup \dots \cup J_{\alpha_N}$  and hence  $A$  is compact.

The proof is complete. □

**Remark 4.18** In fact, the condition in Theorem 4.17(i) is the usual definition of a *compact set* for a general topological space. More precisely, if a set  $A$  satisfies the Definition 4.1, then  $A$  is said to be *sequentially compact*. Theorem 4.17 tells us that the notation of the compactness and the sequentially compactness are the same as in the case of a subset of  $\mathbb{R}$ . However, these two notations are different for a general topological space.



## 5 Cauchy sequences

The following notation is the landmark in the development of the 20th century mathematics.

**Definition 5.1** A sequence  $(x_n)$  is called a Cauchy sequence if it satisfies the following condition:

for any  $\varepsilon > 0$ , there is a positive integer  $N$  so that  $|x_m - x_n| < \varepsilon$  whenever  $m, n \geq N$ .

**Remark 5.2** According to the definition of a Cauchy sequence, a sequence  $(x_n)$  is not a Cauchy sequence if there is  $\varepsilon > 0$  so that for any positive integer  $N$ , we can find some  $m, n \geq N$  such that  $|x_m - x_n| \geq \varepsilon$ .

**Theorem 5.3 Cauchy Criterion:** A sequence  $(x_n)$  is convergent if and only if it is a Cauchy sequence.

**Proof:** The necessary part is clear. In fact, if  $(x_n)$  is a convergent sequence with the limit  $L$ , then for any  $\varepsilon > 0$ , there is a positive integer  $N$  such that  $|x_n - L| < \varepsilon$  for all  $n \geq N$ . Therefore, we have

$$|x_m - x_n| \leq |x_m - L| + |L - x_n| < 2\varepsilon \quad \text{as } m, n \geq N.$$

Conversely, we assume that  $(x_n)$  is a Cauchy sequence.

We first Claim that  $(x_n)$  is a bounded sequence. In fact, since  $(x_n)$  is a Cauchy, we can find a positive integer  $N_1$  such that  $|x_m - x_{N_1}| < 1$  for all  $m \geq N_1$  and thus,  $|x_m| < 1 + |x_{N_1}|$  for all  $m \geq N_1$ . Therefore, we have  $|x_m| \leq \max(|x_1|, \dots, |x_{N_1-1}|, |x_{N_1}| + 1)$  for all positive integers  $m$ .

The Bolzano-Weierstrass Theorem tells us that  $(x_n)$  has a convergent subsequence  $(x_{n_k})$ . Let  $L := \lim_k x_{n_k}$ . If we show that  $L$  is the limit of  $(x_n)$ , then the proof is finished.

Let  $\varepsilon > 0$ . Then there is a positive integer  $N$  such that  $|x_m - x_n| < \varepsilon$  as  $m, n \geq N$ . On the other hand, since  $L = \lim_k x_{n_k}$ , we can choose  $K$  large enough such that  $|L - x_{n_K}| < \varepsilon$  and  $n_K > N$ . This implies that for any  $n \geq N$ , we have

$$|x_n - L| < |x_n - x_{n_K}| + |x_{n_K} - L| < 2\varepsilon.$$

The proof is complete. □

**Example 5.4** Let  $s_n = \sum_{k=1}^n 1/k$ . Then  $(s_n)$  is not a Cauchy sequence and thus,  $(s_n)$  is divergent.

In fact, it is noted that for  $n \leq m$ , we have

$$|s_m - s_n| = \frac{1}{n+1} + \dots + \frac{1}{m} \geq \frac{m-n}{m}.$$

Hence, we always have  $|s_{2n} - s_n| \geq \frac{1}{2}$  for all  $n$ . Thus, if we take  $\varepsilon = 1/2$ , then for any positive integer  $N$  by taking  $n = N$  and  $m = 2N$ , we have  $|s_{2N} - s_N| > 1/2 = \varepsilon$ . Hence,  $(s_n)$  is not a Cauchy sequence.

**Remark 5.5** A sequence  $(x_n)$  properly converges to  $+\infty$  (resp.  $-\infty$ ) if for any  $M > 0$ , there is a positive integer  $N$  so that  $x_n > M$  (resp.  $x_n < -M$ ) for all  $n \geq M$ . In this case, we write  $\lim x_n = \infty$  (resp.  $\lim x_n = -\infty$ ). **Warning!!!** In this case, the sequence  $(x_n)$  is still divergent since  $\infty$  is **NOT** a real number, hence,  $\infty$  is not the limit of  $(x_n)$ .

Note that the sequence  $(s_n)$  in Example 5.4 properly converges to  $+\infty$ . From this we see that the sequence  $(\sum_{k=1}^n \frac{1}{n^\alpha})_{n=1}^\infty$  also diverges properly to  $+\infty$  if  $\alpha \leq 1$ .

However, a divergent sequence may not converge properly to  $\infty$ , for example, if we take  $x_n = 0$  as  $n$  is odd; otherwise,  $x_n = n$ .

**Example 5.6** Let  $t_n = \sum_{k=1}^n \frac{1}{k^2}$ . Then the sequence  $(t_n)$  is convergent. Using the Cauchy Theorem, we need to show that  $(t_n)$  is a Cauchy sequence.

It is noted that for  $n \leq m$ , we have

$$|t_m - t_n| = \sum_{k=n+1}^m \frac{1}{k^2} \leq \sum_{k=n+1}^m \frac{1}{(k-1)k} = \sum_{k=n+1}^m (\frac{1}{k-1} - \frac{1}{k}) = \frac{1}{n} - \frac{1}{m} < \frac{1}{n}.$$

Thus, if we are given  $\varepsilon > 0$ , then we choose a positive integer  $N$  so that  $\frac{1}{n} < \varepsilon$  for all  $n \geq N$ . Therefore,  $|t_m - t_n| < \varepsilon$  whenever  $m \geq n \geq N$ . The proof is complete.

**Remark 5.7** We have the following implications in  $\mathbb{R}$ .

Axiom of Completeness  $\Rightarrow$  Bounded Monotone Convergent Theorem (Theorem 2.13)  $\Rightarrow$   
Nested Intervals Theorem  $\Rightarrow$  Bolzano-Weierstrass Theorem  $\Rightarrow$  Cauchy Theorem.

**Everything is due to the Axiom of Completeness.**

## 6 Appendix: Bolzano-Weierstrass Theorem and Cauchy Criterion in $\mathbb{R}^m$

Throughout this section, for each element  $x \in \mathbb{R}^m$ , we write  $x := (x(1), \dots, x(m))$  and put

$$\|x\| := \sqrt{\sum_{k=1}^m |x(k)|^2}.$$

We call  $\|x\|$  the norm of  $x$ . Clearly, we have

$$\max_{1 \leq k \leq m} |x(k)| \leq \|x\| \leq \sqrt{m} \max_{1 \leq k \leq m} |x(k)| \tag{6.1}$$

for all  $x \in \mathbb{R}^m$ .

For each element  $x$  and  $y$  in  $\mathbb{R}^m$ , the distance between  $x$  and  $y$  is defined by  $\|x - y\|$ . In this case, one can define naturally the notation of a convergent sequence in  $\mathbb{R}^m$  as in the  $\mathbb{R}$  case. More precisely, we have the following definition.

**Definition 6.1** A sequence  $(x_n)$  in  $\mathbb{R}^m$  is said to be convergent if there is an element  $u \in \mathbb{R}^m$  such that  $\lim_n \|x_n - u\| = 0$ . Clearly such element is unique if it exists. In this case we call  $u$  the limit of  $(x_n)$  and write  $u := \lim x_n$ .

By using Eq 6.1, we have the following immediately.

**Lemma 6.2** *Using the notation as above, a sequence  $(x_n)$  converges to  $u$  in  $\mathbb{R}^m$  if and only if the sequence  $(x_n(k))$  converges to  $u(k)$  in  $\mathbb{R}$  for all  $k = 1, \dots, m$ .*

Naturally, one can also have the following notation as in the  $\mathbb{R}$  case.

**Definition 6.3** (i) *A point  $z$  in  $\mathbb{R}^m$  is said to be a limit point of a subset  $A$  of  $\mathbb{R}^m$  if for every  $r > 0$ , there is a point  $a \in A$  such that  $0 < \|z - a\| < r$ . Also,  $A$  is said to be a closed set if  $A$  contains all its limit points.*

(ii) *A Cauchy sequence  $(x_n)$  in  $\mathbb{R}^m$  means that if for every  $\varepsilon > 0$ , there is a positive integer  $N$  such that  $\|x_m - x_n\| < \varepsilon$  whenever  $m, n \geq N$ .*

**Theorem 6.4 Cauchy Criterion:** *A sequence in  $\mathbb{R}^m$  is convergent if and only if it is a Cauchy sequence.*

**Proof:** The Eq 6.1 tells us that if a sequence  $(x_n)$  in  $\mathbb{R}^m$  is a Cauchy sequence, then so is for each coordinate sequence  $(x_n(k))$  in  $\mathbb{R}$  for  $k = 1, \dots, m$ . Hence the result is obtained immediately by using the Cauchy criterion for the real case.  $\square$

In the rest of this section, we are going to show the Bolzano-Weierstrass Theorem in the higher dimensional case. Recall that a subsequence of  $(x_n)$  in  $\mathbb{R}^m$  is a sequence in  $\mathbb{R}^m$  given by a strictly increasing function  $\phi : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ , i.e., it is  $(x_{\phi(j)})_{j=1}^{\infty}$ .

**Theorem 6.5 Bolzano-Weierstrass Theorem:** *Every bounded sequence in  $\mathbb{R}^m$  has a convergent subsequence.*

**Proof:** Let  $(x_n)$  be a bounded sequence in  $\mathbb{R}^m$ . Thanks to the Eq6.1 again, each coordinate sequence  $(x_n(k))$  is also a bounded sequence of real numbers for  $k = 1, \dots, m$ . As  $k = 1$ , by the Bolzano-Weierstrass for the real sequence case, there is a convergent subsequence  $(x_{\phi_1(j)}(1))_{j=1}^{\infty}$  of  $(x_n(1))$ , where  $\phi_1 : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$  is a strictly increasing function. As  $k = 2$ , we consider the subsequence  $(x_{\phi_1(j)})$  of  $(x_n)$ . Using the Bolzano-Weierstrass for the real sequence case again, there is a convergent subsequence  $(x_{\phi_2(j)}(2))_{j=1}^{\infty}$  of  $(x_{\phi_1(j)}(2))$ , where  $\phi_2 : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$  is a strictly increasing function. Next we consider the subsequence  $(x_{\phi_2 \circ \phi_1(j)})$  of  $(x_{\phi_1(j)})$ , and so is the subsequence of  $(x_n)$ , for the case of  $k = 3$ . To repeat the same step, we get a subsequence  $(x_{\phi_m \circ \dots \circ \phi_1(j)})$  of  $(x_n)$  so that each coordinate sequence  $(x_{\phi_m \circ \dots \circ \phi_1(j)}(k))$  is convergent for all  $k = 1, \dots, m$ . Then by the Eq 6.1 again, the subsequence  $(x_{\phi_m \circ \dots \circ \phi_1(j)})$  is convergent in  $\mathbb{R}^m$  as desired.  $\square$

## 7 Limits of functions

Throughout this section let  $f$  be a real-valued function defined on a subset  $A$  of  $\mathbb{R}$ .

A point  $x_0$  is called a *limit point* of  $A$  if for any  $r > 0$ , there is some element  $a \in A$  such that  $0 < |x_0 - a| < r$ . We write  $D(A)$  for the set of all limit points of  $A$ . Note that a limit point of  $A$  may not sit in  $A$ .

**Definition 7.1** Let  $c \in D(A)$ . A number  $L$  is said to be a limit of  $f$  at  $c$  (note that  $f(c)$  may not be defined!!) if for any  $\varepsilon$ , there is  $\delta = \delta(\varepsilon) > 0$  (depends the choice of  $\varepsilon$ ) such that

$$|f(x) - L| < \varepsilon \quad \text{whenever } x \in A \text{ and } 0 < |x - c| < \delta .$$

(Note: we only consider those points in  $A$  which are very close to  $c$  but do not equal to  $c$ !!!)

**Remark 7.2** A number  $L$  is not a limit of  $f$  at  $c$  means if there is  $\varepsilon > 0$  so that for any  $\delta$ , we can find some  $x' \in A$  with  $|x' - c| < \delta$  but  $|f(x') - L| \geq \varepsilon$ .

**Proposition 7.3** Using the notation as above if  $f$  has a limit at  $c$ , then its limit is unique. Consequently, if we write  $\lim_{x \rightarrow c} f(x)$  for the limit of  $f$  at  $c$ , then this notation is well defined.

**Proof:** Let  $L'$  be a another limit of  $f$  at  $c$ . Let  $\varepsilon > 0$ . Then by the definition above, there are some positive numbers  $\delta$  and  $\delta'$  so that  $|f(x) - L| < \varepsilon$  for any  $x \in A$  with  $0 < |x - c| < \delta$ . Similarly, we have  $|f(x) - L'| < \varepsilon$  for any  $x \in A$  with  $0 < |x - c| < \delta'$ . Since  $c \in D(A)$ , we can find some  $a \in A$  such that  $0 < |c - a| < \delta''$ , where  $\delta'' = \min(\delta, \delta')$ . This gives

$$|L - L'| \leq |L - f(a)| + |f(a) - L'| < 2\varepsilon.$$

Since  $\varepsilon > 0$  is arbitrary, we have  $L = L'$  as desired. □

**Example 7.4** Let  $A = (0, \infty)$ . Define  $f(x) := x^2 \sin \frac{1}{x}$ .

(i) Show that  $\lim_{x \rightarrow 0} f(x) = 0$ .

In fact, it is noted that  $|x^2| \leq |x|$  for all  $x \in (0, 1)$ . Let  $\varepsilon > 0$ . Thus, if we take  $0 < \delta = \min(\varepsilon, 1)$ , then we have

$$|f(x) - 0| \leq |x^2| \leq |x| < \varepsilon$$

whenever  $x > 0$  with  $|x - 0| < \delta$ .

(ii) Using the  $\varepsilon$ - $\delta$  notation, show that  $\lim_{x \rightarrow 0} f(x) \neq 1$ .

Note that if we take  $\varepsilon = 1/2$ , then for any  $\delta > 0$ , we choose a positive integer  $N$  such that  $0 < |\frac{1}{N\pi} - 0| < \delta$ , and we have

$$|f(\frac{1}{N\pi}) - 1| = 1 > \varepsilon.$$

Therefore, 1 is not the limit of  $f$  at 0.

**Proposition 7.5** Using the notation as above, let  $c$  be a limit point of  $A$ . Then the following are equivalent.

(i)  $\lim_{x \rightarrow c} f(x)$  exists.

(ii) If  $(x_n)$  is a convergent sequence in  $A \setminus \{c\}$  with  $\lim x_n = c$ , then the sequence  $(f(x_n))$  is convergent.

In this case,  $\lim_{x \rightarrow c} f(x) = \lim f(x_n)$  whenever a convergent sequence  $(x_n)$  in  $A \setminus \{c\}$  with  $\lim x_n = c$ .

**Proof:** For showing  $(i) \Rightarrow (ii)$ , we assume that  $L = \lim_{x \rightarrow c} f(x)$  exists. Let  $(x_n)$  be a convergent sequence in  $A \setminus \{c\}$  with the limit  $c$ . Let  $\varepsilon > 0$ . Then by the definition of the limit of a function, we can find  $\delta > 0$  such that  $|f(x) - L| < \varepsilon$  whenever  $x \in A$  with  $0 < |x - c| < \delta$ . On the other hand, since  $\lim x_n = c$  and  $x_n \neq c$  for all  $n$ , there is a positive integer  $N$  such that  $0 < |x_n - c| < \delta$  for all  $n \geq N$  and thus,  $|f(x_n) - L| < \varepsilon$  for all  $n \geq N$ . Thus, the condition  $(ii)$  holds.

Suppose that the condition  $(ii)$  holds. We first claim that the sequence  $(f(x_n))$  converges to the same limit whenever  $(x_n)$  is a convergent sequence in  $A \setminus \{c\}$  such that  $\lim x_n = c$ . In fact, suppose that there are two sequences  $(x_n)$  and  $(y_n)$  in  $A \setminus \{c\}$  with  $\lim x_n = \lim y_n = c$  such that  $u := \lim f(x_n)$  and  $v := \lim f(y_n)$  are not the same. Put  $w_n := x_k$  if  $n = 2k$  and  $w_n := y_k$  if  $n = 2k + 1$  is odd. Then  $\lim w_n = c$ . The assumption  $(ii)$  implies that  $\lim f(w_n)$  exists. However,  $(f(x_n))$  and  $(f(y_n))$  both are the subsequences of  $(f(w_n))$  and they converge to the different limits. It leads to a contradiction.

Now we fix a sequence  $(x_n)$  in  $A \setminus \{c\}$  such that  $\lim x_n = c$ . Then by the assumption  $L := \lim f(x_n)$  exists. Suppose that  $L$  is not the limit of  $f(x)$  at  $c$ . Thus, there is  $\varepsilon > 0$  such that for any  $\delta > 0$ , we can find some  $x' \in A$  with  $0 < |x' - c| < \delta$  but  $|f(x') - L| \geq \varepsilon$ . From this, we see that for each positive integer  $n$ , there is  $x'_n \in A$  with  $0 < |x'_n - c| < 1/n$  but  $|f(x'_n) - L| \geq \varepsilon$ . Thus, the sequence  $(x'_n)$  sits in  $A \setminus \{c\}$  and converges to  $c$  but  $L$  is not the limit of the sequence  $(f(x'_n))$ . This contradicts to the above observation. Therefore, the part  $(i)$  holds.  $\square$

Proposition 7.5, together with Proposition 2.8, we have the following assertion immediately.

**Proposition 7.6** *Let  $f$  and  $g$  be the functions defined on  $A$ . Let  $c$  be a limit point of  $A$ . Assume that  $L := \lim_{x \rightarrow c} f(x)$  and  $R := \lim_{x \rightarrow c} g(x)$  both exist. Then we have the following statements.*

1.  $\lim_{x \rightarrow c} (f + g)(x)$  exists and  $\lim_{x \rightarrow c} (f + g)(x) = L + R$
2.  $\lim_{x \rightarrow c} (f \cdot g)(x)$  exists and  $\lim_{x \rightarrow c} (f \cdot g)(x) = L \cdot R$ .
3. if we further assume that  $g(x) \neq 0$  for all  $x \in A$  and  $R \neq 0$ , then  $\lim_{x \rightarrow c} (f/g)(x)$  exists and  $\lim_{x \rightarrow c} (f/g)(x) = L/R$ .

The following result is regarded as the Cauchy criterion in the case of functions.

**Proposition 7.7** *Using the notation as before,  $\lim_{x \rightarrow c} f(x)$  exists if and only if for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $|f(x') - f(x'')| < \varepsilon$  whenever  $x', x'' \in A$  with  $0 < |x' - c| < \delta$  and  $0 < |x'' - c| < \delta$ .*

**Proof:** For showing  $(\Rightarrow)$  we assume that  $L := \lim_{x \rightarrow c} f(x)$  exists. Let  $\varepsilon > 0$ . Then there is  $\delta > 0$  such that  $|f(x) - L| < \varepsilon$  as  $x \in A$  with  $0 < |x - c| < \delta$ . Thus, if  $x', x'' \in A$  with  $0 < |x' - c| < \delta$  and  $0 < |x'' - c| < \delta$ , we see that

$$|f(x') - f(x'')| \leq |f(x') - L| + |L - f(x'')| < 2\varepsilon.$$

Hence, the necessary condition holds.

Note that since  $c$  is a limit point of  $A$ , we can find a sequence Let  $(x_n)$  in  $A \setminus \{c\}$  such that  $\lim x_n = c$ . Then the necessary condition implies that  $(f(x_n))$  is a Cauchy sequence. In fact,

for any  $\varepsilon > 0$ , the necessary condition above gives  $\delta > 0$  so that  $|f(x') - f(x'')| < \varepsilon$  whenever  $x', x'' \in A$  with  $0 < |x' - c| < \delta$  and  $0 < |x'' - c| < \delta$ . Since  $\lim x_n = c$ , there is a positive integer  $N$  such that  $|x_n - c| < \delta$  for all  $n \geq N$  and hence, we have  $|f(x_n) - f(x_m)| < \varepsilon$  for all  $m, n \geq N$ . Thus,  $(f(x_n))$  is a Cauchy sequence. Hence,  $\lim f(x_n)$  exists. The proof is complete by using Proposition 7.5.

□

**Definition 7.8** Using the notation as before, let  $f$  be a function defined on  $A$  and let  $c$  be a limit point of  $A$ .

1. We say that  $f$  diverges to  $+\infty$  (resp.  $-\infty$ ) as  $x$  tends to  $c$  if for any  $M > 0$ , there is  $\delta > 0$  such that  $f(x) > M$  (resp.  $f(x) < -M$ ) as  $x \in A$  with  $0 < |x - c| < \delta$ . In this case, write  $\lim_{x \rightarrow c} f(x) = +\infty$  (resp.  $\lim_{x \rightarrow c} f(x) = -\infty$ ).
2. We further suppose that  $A$  is not bounded above. We say that  $f$  has a limit  $L$  as  $x$  tends to  $+\infty$  if for any  $\varepsilon > 0$ , there is a positive number  $R > 0$  such that  $|f(x) - L| < \varepsilon$  as  $x \in A$  with  $x > R$ . In this case, a limit must be unique if it exists. Write  $\lim_{x \rightarrow \infty} f(x) = L$ .  
Similarly, one can define the notion  $\lim_{x \rightarrow -\infty} f(x) = L$  when  $A$  is not bounded below. For simply, when we are talking about notion  $\lim_{x \rightarrow \infty} f(x)$ ,  $A$  has been assumed to be unbounded above in advance.
3. Similarly, one can give a suitable definition for the notion:  $\lim_{x \rightarrow \infty} f(x) = +\infty$ .

**Proposition 7.9** Using the notation as before, let  $f, g$  be the functions defined on  $A$ .

- (i) If  $\lim_{x \rightarrow c} f(x) = +\infty$  and  $\lim_{x \rightarrow c} g(x)$  exists, then  $\lim_{x \rightarrow c} (f + g)(x) = +\infty$ .
- (ii) If  $\lim_{x \rightarrow c} f(x) = +\infty$  and  $\lim_{x \rightarrow c} g(x) > 0$  exists, then  $\lim_{x \rightarrow c} (f \cdot g)(x) = +\infty$ .
- (iii) If  $\lim_{x \rightarrow \infty} f(x) = +\infty$  and  $\lim_{x \rightarrow \infty} g(x) > 0$  exists, then  $\lim_{x \rightarrow \infty} (f \cdot g)(x) = +\infty$ .

**Proof:** For showing part (ii), let  $M > 0$ . Since  $l := \lim_{x \rightarrow c} g(x) > 0$ , there is  $\delta_1 > 0$  so that  $g(x) > l - \frac{l}{2} = \frac{l}{2} > 0$  for all  $x \in A$  with  $0 < |x - c| < \delta_1$ . Moreover,  $\lim_{x \rightarrow c} f(x) = +\infty$ , and so we can find  $0 < \delta < \delta_1$  such that  $f(x) > \frac{2M}{l}$  as  $x \in A$  with  $0 < |x - c| < \delta$  and hence in this case, we have

$$f(x)g(x) > \frac{2M}{l} \cdot \frac{l}{2} = M.$$

Part (iii) follows.

Using the similar argument, try to finish the proof by yourself. □

**Remark 7.10** The assumption of the non-zero limits in Proposition 7.9(ii) and (iii) cannot be removed. For example, by considering  $f(x) := 1/x; g(x) := x$  for  $x > 0$ , note that  $\lim_{x \rightarrow 0} f(x) = \infty$  and  $\lim_{x \rightarrow 0} g(x) = 0$  but  $f(x)g(x) = 1$  for all  $x > 0$ .

**Example 7.11** Let  $p(x) := a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  be a polynomial of degree  $n > 0$ , where  $x \in \mathbb{R}$ . If the leading coefficient  $a_n$  of  $p$  is positive, then  $\lim_{x \rightarrow +\infty} f(x) = +\infty$ . In fact, since  $a_n \neq 0$ , we see that

$$p(x) = a_n x^n \left( 1 + \frac{a_{n-1}}{a_n} x^{-1} + \frac{a_{n-2}}{a_n} x^{-2} + \dots + \frac{a_0}{a_n} x^{-n} \right)$$

for all  $x > 0$ . In addition, since  $a_n > 0$  and  $n > 0$ , clearly we have  $\lim_{x \rightarrow +\infty} a_n x^n = +\infty$ . The result follows immediately from Proposition 7.9.

**Definition 7.12** A point  $c$  is called a right (resp. left) limit point of  $A$  if for any  $r > 0$ , there is some  $x \in A$  such that  $0 < x - c < r$  (resp.  $0 < c - x < r$ ), i.e.,  $(c, c + r) \cap A \neq \emptyset$  (resp.  $(c - r, c) \cap A \neq \emptyset$ ). Write  $D_r(A)$  (resp.  $D_l(A)$ ) for the set of right (resp. left) limit points of  $A$ .

Clearly, we have  $D_r(A) \cup D_l(A) = D(A)$ .

**Example 7.13** We have the following examples.

1. If  $A = (0, 1) \cup \{2\}$ , then  $D_r(A) = [0, 1)$  and  $D_l(A) = (0, 1]$ .
2. If  $A = \{1, 1/2, 1/3, \dots\}$ , then  $D_r(A) = \{0\}$  and  $D_l(A) = \emptyset$ .

**Definition 7.14** Using the notation as above, let  $c \in D_r(A)$ . We say that  $f$  has a right (resp. left) limit  $L$  of  $f$  at  $c$  if for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $|f(x) - L| < \varepsilon$  for all  $x \in A$  with  $0 < x - c < \delta$  (resp.  $0 < c - x < \delta$ ).

It is noted that if a right (resp. left) limit exists, then it is unique.

We write  $\lim_{x \rightarrow c+} f(x)$  and  $\lim_{x \rightarrow c-} f(x)$  for the right and left limit respectively.

**Example 7.15** Let  $A = \mathbb{R} \setminus \{0\}$ . Define  $f(x) = 1$  if  $x > 0$ ; otherwise,  $f(x) = -1$ . Then  $\lim_{x \rightarrow 0+} f(x) = 1$  and  $\lim_{x \rightarrow 0-} f(x) = -1$ . This function is called the sign function.

We always denote it by  $\text{sgn}(x)$ .

**Proposition 7.16** Let  $c \in D_r(A) \cap D_l(A)$ . Then  $\lim_{x \rightarrow c} f(x)$  exists if and only if  $\lim_{x \rightarrow c+} f(x)$  and  $\lim_{x \rightarrow c-} f(x)$  both exist and  $\lim_{x \rightarrow c+} f(x) = \lim_{x \rightarrow c-} f(x)$ .

In this case, we have  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c+} f(x) = \lim_{x \rightarrow c-} f(x)$ .

**Proposition 7.17** Let  $f(x)$  be a function defined on  $(0, \infty)$  and  $g(x) = f(1/x)$ . Then  $\lim_{x \rightarrow +\infty} f(x)$  exists if and only if  $\lim_{x \rightarrow 0+} g(x)$  exists.

In this case, we have  $\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow 0+} g(x)$ .

## 8 Continuous functions

Throughout this section, let  $A$  be a non-empty subset of  $\mathbb{R}$  and let  $f$  be a function defined on  $A$ .

**Definition 8.1** Let  $c \in A$ . We say that a function  $f$  is continuous at  $c$  if for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $|f(x) - f(c)| < \varepsilon$  whenever  $x \in A$  with  $|x - c| < \delta$ .

Furthermore,  $f$  is said to be continuous on  $A$  if it is continuous at every point in  $A$ .

**Remark 8.2** Using the notation as above, note that

1. A function  $f$  is discontinuous at  $c$  if there is  $\varepsilon > 0$  so that for any  $\delta > 0$ , we can find some  $x \in A$  satisfying  $|x - c| < \delta$  but  $|f(x) - f(c)| \geq \varepsilon$ .

2. If a point  $c \in A$  is not a limit point of  $A$ , then a function  $f$  is continuous automatically at  $c$ . In fact, if  $c \in A$  is not a limit point of  $A$ , then there is  $r > 0$  such that  $(c-r, c+r) \cap A = \{c\}$ . Therefore, for any  $\varepsilon > 0$ , we can choose  $\delta = r$  in the Definition 8.1 above.

**Definition 8.3** Let  $A$  be subset of  $\mathbb{R}$ . A subset  $E$  of  $A$  is called an open subset of  $A$  if for each  $c \in E$ , there is  $r > 0$  such that  $(c-r, c+r) \cap A \subseteq E$ .

**Remark 8.4 Warning !!!!:** Notice that if  $E$  an open subset of  $A$ , it does not imply that  $E$  is an open subset of  $\mathbb{R}$ .

For example, if we consider  $A = [0, 1]$  and  $E = (0, 1]$ , then  $E$  is an open subset of  $A$  but it is not an open subset of  $\mathbb{R}$ .

The following result is directly obtained from the definition of a continuous function.

**Proposition 8.5** Using the notation as above, then a function  $f$  is continuous on  $A$  if and only the pre-image  $f^{-1}(V) := \{x \in A : f(x) \in V\}$  of any open subset  $V$  of  $\mathbb{R}$  is an open subset of  $A$ .

**Proposition 8.6** Let  $c \in A$ . Then we have the following assertions.

- (i) If  $c \in A$  is a limit point of  $A$ , then  $f$  is continuous at  $c$  if and only if  $\lim_{x \rightarrow c} f(x) = f(c)$ .
- (ii)  $f$  is continuous at  $c$  if and only if whenever a sequence  $(x_n)$  in  $A$  with  $\lim x_n = c$ , we have  $\lim f(x_n) = f(c)$ .

**Proof:** Part (i) follows directly from the Definition 8.1.

Part (ii) can be obtained by using a similar argument as in Proposition 7.6. Try to do it by yourself. □

**Proposition 8.7** Let  $c \in A$  and let  $f, g$  be functions defined on  $A$ . If  $f, g$  are continuous at  $c$ , then we have the following assertions.

- (i) The function  $f + g$  is continuous at  $c$ .
- (ii) The product  $f \cdot g$  is continuous at  $c$ .
- (iii) Moreover, if  $g(x) \neq 0$  for all  $x \in A$ , then  $f/g$  is continuous at  $c$ .
- (iv) Moreover, if the image of  $f$  is contained in a subset  $B$  of  $\mathbb{R}$  and  $h : B \rightarrow \mathbb{R}$  is continuous at  $f(c)$ , then the composition  $h \circ f$  is continuous at  $c$ .

**Proof:** The above assertions follows immediately from Propositions 2.8 and 8.6. Alternatively, they can be shown directly by the definition.

For showing part (ii), since  $g$  is continuous at  $c$ , there is  $\delta_1 > 0$  such that  $|f(x) - f(c)| < 1$  and hence,  $|f(x)| < 1 + |f(c)|$  for all  $x \in A$  with  $|x - c| < \delta_1$ . Using the continuity of  $f$  and  $g$  at  $c$ , there exists  $0 < \delta < \delta_1$  so that  $|f(x) - f(c)| < \varepsilon$  and  $|g(x) - g(c)| < \varepsilon$  as  $x \in A$  and  $|x - c| < \delta$ . Therefore, we have

$$|f(x)g(x) - f(c)g(c)| \leq |f(x)g(x) - f(c)g(x)| + |f(c)g(x) - f(c)g(c)| \leq \varepsilon(1 + |g(c)| + |f(c)|)$$



as  $x \in A$  and  $|x - c| < \delta$ . Part (ii) follows.

By using part (ii), we need to show that the function  $1/g(x)$  is continuous at  $c$ . Note that we may assume  $g(c) > 0$  (otherwise by considering  $-g(x)$ ).  $g(x)$  is continuous at  $c$ , and so there is  $\delta_1 > 0$  so that  $|g(x) - g(c)| < g(c)/2$  and hence,  $g(x) > g(c)/2$  for all  $x \in A$  and  $|x - c| < \delta_1$ . Now let  $\varepsilon > 0$ , there is  $0 < \delta < \delta_1$  so that  $|g(x) - g(c)| < \delta$  as  $x \in A$  and  $|x - c| < \delta$ . Therefore, we have

$$\left| \frac{1}{g(x)} - \frac{1}{g(c)} \right| = \frac{|g(x) - g(c)|}{g(x)g(c)} \leq \frac{2\varepsilon}{g(c)^2}$$

for all  $x \in A$  with  $|x - c| < \delta$ . The proof of (iii) is complete.

The last assertion follows clearly from the definition. □

Before showing the following important result, we first recall that a subset  $A$  is said to be compact if for any sequence  $(x_n)$  in  $A$  has a convergent subsequence  $(x_{n_k})$  such that  $\lim_k x_{n_k} \in A$ . Moreover,  $A$  is compact if and only if it is a closed and bounded set.

**Theorem 8.8** *If  $f$  is a continuous function defined on a compact set  $A$ , then  $f$  is a bounded function. Moreover, there are  $x_1$  and  $x_2$  in  $A$  such that  $f(x_1) = \min\{f(x) : x \in A\}$  and  $f(x_2) = \max\{f(x) : x \in A\}$ .*

**Proof:** First, we show that the set  $f(A)$  is bounded above. We give two different methods about the claim.

**Method I:**

Suppose that  $f$  is not bounded above. Then for each positive integer  $n$ , there is  $x_n \in A$  such that  $f(x_n) \geq n$ .  $A$  is compact, and so there is a convergent subsequence  $(x_{n_k})$  with  $c := \lim x_{n_k} \in A$ . Note that since  $f$  is continuous at  $c$ , we see that the sequence  $(f(x_{n_k}))$  converges to  $f(c)$  and thus,  $(f(x_{n_k}))$  is a bounded sequence but  $f(x_{n_k}) \geq n_k$  for all  $k$ . It leads to a contradiction because  $n_k \rightarrow \infty$  as  $k \rightarrow \infty$ .

**Method II:**

Notice that since  $f$  is continuous on  $A$ , then for each element  $x \in A$ , there is  $\delta(x) > 0$  such that  $f(u) < f(x) + 1$  as  $u \in A$  with  $|u - x| < \delta(x)$ . Note that if we put  $J_x := (x - \delta(x), x + \delta(x))$  for  $x \in A$ , then we have  $A \subseteq \bigcup_{x \in A} J_x$ . Using the compactness of  $A$ , there are finitely many  $x_1, \dots, x_N \in A$  such that  $A \subseteq J_{x_1} \cup \dots \cup J_{x_N}$ . Then implies that  $f(x) \leq \max(1 + f(x_1), \dots, 1 + f(x_N))$  for all  $x \in A$ , so the set  $f(A)$  is bounded above.

Next, we want to show that  $f(a) = \max\{f(x) : x \in A\}$  for some  $a \in A$ .

In fact, note that since the set  $\{f(x) : x \in A\}$  is bounded above,  $L := \sup\{f(x) : x \in A\}$  exists. Thus, there exists a sequence  $(x_n)$  in  $A$  such that  $\lim f(x_n) = L$ . Using the compactness of  $A$ , there is a convergent subsequence  $(x_{n_k})$  of  $(x_n)$  with  $a := \lim x_{n_k} \in A$ . Thus, we have  $f(a) = \lim f(x_{n_k})$  and thus,  $f(a) = L$  as desired.

By considering  $-f$ , we get  $f(x_1) = \min\{f(x) : x \in A\}$  for some  $x_1 \in A$ . The proof is complete. □

**Remark 8.9** The assumption of compactness in Theorem 8.8 cannot be removed.

For example if  $A = [1, \infty)$  and  $f(x) = 1/x$  for  $x \in A$ , then there is no points attains its minimum on  $A$  although  $f$  is a bounded function.

**Theorem 8.10** *If  $f$  is a continuous function defined on a compact set, then the image  $f(A) := \{f(x) : x \in A\}$  is compact.*

**Proof: Method I:**

It suffices to show that  $f(A)$  is a closed and bounded set. We have shown that  $f(A)$  is bounded by Theorem 8.8. We need to show that  $f(A)$  is closed. By applying Proposition 4.7, we need to claim that if  $(x_n)$  is a sequence in  $A$  so that  $(f(x_n))$  is convergent, then the limit  $L := \lim f(x_n) \in f(A)$ . Indeed, by the compactness of  $A$ ,  $(x_n)$  has a convergent subsequence  $(x_{n_k})$  such that  $c := \lim x_{n_k} \in A$ .  $f$  is continuous at  $c$ , and so  $\lim f(x_{n_k}) = f(c)$  and thus,  $L = f(c) \in f(A)$  as required.

**Method II:**

Let  $\{J_i\}_{i \in I}$  be an open cover of  $f(A)$ . Since  $f$  is continuous on  $A$ , Proposition 8.5 implies that for each element  $a \in A$ , there are  $\delta_a > 0$  and  $i_a \in I$  such that  $f((a - \delta_a, a + \delta_a) \cap A) \subseteq J_{i_a}$ . Note that we have

$$A \subseteq \bigcup_{a \in A} (a - \delta_a, a + \delta_a).$$

Then by the compactness of  $A$ , there are finitely many  $a_1, \dots, a_N$  in  $A$  such that

$$A \subseteq \bigcup_{k=1}^N (a_k - \delta_{a_k}, a_k + \delta_{a_k}).$$

Therefore, we have

$$f(A) \subseteq \bigcup_{k=1}^N f((a_k - \delta_{a_k}, a_k + \delta_{a_k}) \cap A) \subseteq \bigcup_{k=1}^N J_{i_{a_k}}.$$

□

**Remark 8.11** In general, the image of a closed set under a continuous map is not necessarily closed. For example,  $A = [1, \infty)$  and  $f(x) = 1/x, x \in A$ . Note that  $A$  is a closed set but  $f(A) = (0, 1]$  is not closed.

**Definition 8.12** *Two subsets  $A$  and  $B$  are said to be homeomorphic if there is a bijection  $f$  from  $A$  onto  $B$  such that  $f$  and the inverse  $f^{-1}$  both are continuous. In this case,  $f$  is called a homeomorphism.*

**Proposition 8.13** *Suppose that  $A$  and  $B$  are homeomorphic. If  $A$  is compact, then so is  $B$ .*

**Proof:** It can be shown directly by Theorem 8.10. □

**Example 8.14** *By applying Theorem 8.10, it is impossible to find a continuous surjection from  $[0, 1]$  onto  $[0, 1)$  because  $[0, 1]$  is compact but  $[0, 1)$  is not. Therefore,  $[0, 1]$  is not homeomorphic to  $[0, 1)$ .*

**Proposition 8.15** *Let  $A$  and  $B$  be non-empty subsets of  $\mathbb{R}$ . Let  $f : A \rightarrow B$  be a continuous bijection. If  $A$  is compact, then  $f$  is a homeomorphism, i.e., the inverse  $f^{-1}$  is continuous.*

**Proof:** Put  $y = f(x)$  and  $g(y) = f^{-1}(x)$ ,  $x \in A$ . Suppose that the function  $g$  is discontinuous at some  $b \in B$ . Then, there is  $\varepsilon > 0$  so that for any  $\delta > 0$ , there is  $y \in B$  so that  $|y - b| < \delta$  but  $|g(y) - g(b)| \geq \varepsilon$ . By considering  $\delta = 1/n$  for  $n = 1, 2, \dots$ . Therefore, there is a sequence  $(y_n)$  in  $B$  so that  $\lim y_n = b$  and  $|g(y_n) - g(b)| \geq \varepsilon$  for all  $n$ . Let  $x_n = g(y_n) \in A$ . Then by the compactness of  $A$ ,  $(x_n)$  has a convergent subsequence  $(x_{n_k})$  such that  $a := \lim x_{n_k} \in A$ . Note that  $b = \lim y_{n_k} = \lim f(x_{n_k}) = f(a)$  because  $f$  is continuous and  $\lim y_n = b$ . Thus,  $a = g(b)$ . Therefore, we have  $\lim g(y_{n_k}) = \lim x_{n_k} = a = g(b)$  which leads to a contradiction because  $|g(y_n) - g(b)| \geq \varepsilon$  for all  $n$ .  $\square$

**Remark 8.16** The assumption of compactness of the domain on Proposition 8.15 cannot be removed. For example, by considering  $A = [0, 1) \cup [1, 2]$  and  $B = [0, 2]$ , a function  $f : A \rightarrow B$  is defined by  $f(x) = x$  for  $x \in [0, 1)$  and  $f(x) = x - 1$  for  $x \in [1, 2]$ . Then  $f$  is a continuous bijection but its inverse is discontinuous at  $y = 1$ . Note that  $A$  is non-compact in this case.

**Theorem 8.17 Intermediate Value Theorem** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a continuous function. Assume that  $f(a) < L < f(b)$ . Then there is  $c \in (a, b)$  such that  $f(c) = L$ .

**Proof:** If we consider the function  $x \in [a, b] \mapsto f(x) - L$ , then we may assume that  $L = 0$ , i.e.,  $f(a) < 0 < f(b)$ . We want to show that there is  $c \in (a, b)$  so that  $f(c) = 0$ .

**Method 1:**

Let  $S := \{x \in [a, b] : f(x) > 0\}$ . Note that  $S$  is non-empty a bounded below set since  $b \in S$  and  $x > b$  for all  $x \in S$ . Thus,  $c := \inf S$  exists. We will show that  $f(c) = 0$ . Note that for each positive integer  $n$ , there is  $x_n \in S$  satisfying  $c \leq x_n < c + 1/n$ , and so  $\lim x_n = c$ . Since  $a \leq x_n \leq b$  for all  $n$ , we see that  $c \in [a, b]$ . By the continuity of  $f$  and  $f(x_n) > 0$  for all  $n$ , we have  $\lim f(x_n) = f(c)$  and  $f(c) \geq 0$ . We want to show that it is impossible if  $f(c) > 0$ . Note that  $c > a$  since  $f(a) < 0$ . Therefore, there is  $\delta > 0$  such that  $a < c - \delta$  and  $|f(x) - f(c)| < f(c)/2$  as  $x \in [a, b]$  with  $|x - c| < \delta$ . Thus, if we fix a point  $x_1$  such that  $a < c - \delta < x_1 < c \leq b$ , then we have  $f(x_1) > f(c)/2 > 0$ . This implies that  $x_1 \in S$  and  $x_1 < c$ . It is a contradiction because  $c$  is a lower bound for the set  $S$ . Therefore,  $f(c) = 0$ .

**Method 2:**

Let  $[a_1, b_1] = [a, b]$ . We want to construct inductively a sequence of closed and bounded intervals  $\{[a_k, b_k]\}_{k=1}^n$ , where  $1 \leq n \leq +\infty$ , satisfying the following conditions.

1.  $[a_1, b_1] \supseteq [a_2, b_2] \supseteq \dots$ .
2.  $b_k - a_k = \frac{1}{2}(b_{k-1} - a_{k-1})$ , for all  $2 \leq k \leq n$ .
3.  $f(a_k) < 0 < f(b_k)$ , for all  $1 \leq k \leq n$ .

Suppose that the sequence of closed and bounded intervals  $([a_k, b_k])$  has been constructed for  $1 \leq k \leq n$ . We want to construct  $[a_{n+1}, b_{n+1}]$  so that it satisfies the conditions (1) – (3) above. Put  $m_n := \frac{a_n + b_n}{2}$ . If  $f(m_n) = 0$ , then the result follows. Otherwise, if  $f(m_n) > 0$ , then we put  $[a_{n+1}, b_{n+1}] = [a_n, m_n]$ . If  $f(m_n) < 0$ , then we put  $[a_{n+1}, b_{n+1}] = [m_n, b_n]$ . Therefore, if  $f(m_n) \neq 0$  for all  $n = 1, 2, \dots$ , then we have an infinite sequence of  $([a_k, b_k])$  satisfying the conditions (1) – (3) above. By applying the Nested Intervals Theorem in this case, we have  $\bigcap_{k=1}^{\infty} [a_k, b_k] = \{c\}$  for some  $c \in [a, b]$ . Note that we have  $\lim a_k = \lim b_k = c$ .  $f$  is continuous at  $c$ , so we have  $f(c) = \lim f(a_k) = \lim f(b_k)$ . From this, together with the condition (3) above, we have  $f(c) = 0$ . The proof is complete.  $\square$

Recall that an interval is a non-empty subset of  $\mathbb{R}$  which is one of the following forms.

1. (Bounded case):  $[a, b]$ ;  $[a, b)$ ;  $(a, b]$  and  $(a, b)$  for  $a < b$ .
2. (Unbounded case):  $[a, +\infty)$ ;  $(a, +\infty)$ ;  $(-\infty, b]$ ;  $(-\infty, b)$  and  $\mathbb{R}$ .

**Proposition 8.18** *Let  $A$  be subset of  $\mathbb{R}$ . Assume that  $A$  has at least two points. Then the followings are equivalent.*

1.  $A$  is an interval.
2. For each pair of elements  $a, b \in A$  with  $a < b$ , we have  $[a, b] \subseteq A$ .

**Proof:** (1)  $\Rightarrow$  (2) is clear. We want to show (2)  $\Rightarrow$  (1). Assume that the condition (2) holds.

First, we assume that  $A$  is bounded. Then  $L := \sup A$  and  $l := \inf A$  both exist. Then  $x \in [l, L]$  for any  $x \in A$ , so  $A \subseteq [l, L]$ . Now, if  $L$  and  $l$  are in  $A$ , then the condition (2) implies that  $[l, L] \subseteq A$  and thus,  $A = [l, L]$ . By using the similar argument for the other cases, i.e.,  $l \in A$  and  $L \notin A$ ;  $l \notin A$  and  $L \in A$ ;  $l \notin A$  and  $L \notin A$ , we see that  $A$  is equal to  $[l, L)$ ;  $(l, L]$  and  $(l, L)$  respectively.

Similarly, the result can be obtained in the unbounded case. □

**Theorem 8.19** *Let  $f$  be a continuous function defined on  $A$ . If  $A$  is an interval, then so is its image  $f(A)$ .*

**Proof:** By using Proposition 8.18, we need to show that  $[c, d] \subseteq f(A)$  whenever  $c, d \in f(A)$  with  $c < d$ . In fact, let  $f(a) = c$  and  $f(b) = d$  for some  $a, b \in A$ . We may assume that  $a < b$ . Note that since  $A$  is an interval, we have  $[a, b] \subseteq A$ . By applying the Intermediate Value Theorem, for any element  $L \in [c, d]$ , there is an element  $x_1$  between  $a$  and  $b$  such that  $L = f(x_1) \in f(A)$ , and hence  $[c, d] \subseteq f(A)$ . The proof is complete. □

**Example 8.20** *By Theorem 8.19, there is no continuous surjections from  $[0, 1]$  onto  $[0, 1] \cup [2, 3]$ . Hence, the set  $[0, 1]$  is not homeomorphic to  $[0, 1] \cup [2, 3]$ .*

## 9 Uniform continuous functions

Throughout this section, let  $f$  be a function defined on a non-empty subset of  $\mathbb{R}$ .

**Definition 9.1** *A function  $f$  is said to be uniformly continuous on  $A$  if for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $|f(x) - f(y)| < \varepsilon$  whenever  $x, y \in A$  with  $|x - y| < \delta$ .*

**Remark 9.2** *A function  $f$  is not uniformly continuous on  $A$  if there is  $\varepsilon > 0$  such that for any  $\delta > 0$ , there are  $x, y \in A$  with  $|x - y| < \delta$  but  $|f(x) - f(y)| \geq \varepsilon$ .*

**Example 9.3** (i) Let  $f(x) = x^2$  for  $x \in [0, \infty)$ . Then  $f$  is not uniformly continuous on  $[0, \infty)$ . In fact for any positive integer  $n$ , we have

$$\left|f\left(n + \frac{1}{n}\right) - f(n)\right| = (2n + 1/n)(1/n) = 2 + \frac{1}{n^2} \geq 2.$$

Therefore, if we let  $\varepsilon = 2$ , then for any  $\delta > 0$ , we choose a positive integer  $n$  such that  $1/n < \delta$ , so  $n + 1/n$  and  $n$  in  $[0, \infty)$  with  $|n + 1/n - n| < \delta$  but  $|f(n + \frac{1}{n}) - f(n)| \geq 2$ . Thus,  $f$  is not uniformly continuous on  $[0, \infty)$ .

Note that from this example we see that a continuous function need not be uniformly continuous on its domain.

(ii) Let  $f(x) = x^2$  for  $x \in [0, 1]$ . Then  $f$  is uniformly continuous on  $[0, 1]$ . In fact for  $x, y \in [0, 1]$  we have

$$|f(x) - f(y)| = |x - y||x + y| \leq 2|x - y|.$$

Let  $\varepsilon > 0$ . Then we can choose  $0 < \delta < \varepsilon/2$ , so we have  $|f(x) - f(y)| \leq 2|x - y| < \varepsilon$  whenever  $x, y \in [0, 1]$  with  $|x - y| < \delta$ . Thus,  $f$  is uniformly continuous on  $[0, 1]$ .

**Theorem 9.4** Let  $f$  be a continuous function on  $A$ . If  $A$  is compact, then  $f$  is uniformly continuous on  $A$ .

**Proof: Method I:**

Suppose that  $A$  is compact but  $f$  is not uniformly continuous on  $A$ . Then there is  $\varepsilon > 0$  such that for any  $\delta > 0$ , there are  $x, y \in A$  with  $|x - y| < \delta$  but  $|f(x) - f(y)| \geq \varepsilon$ . Consider  $\delta = 1/n$  for  $n = 1, 2, \dots$ . Then for any positive integer  $n$ , there are  $x_n$  and  $y_n$  such that  $|x_n - y_n| < 1/n$  but  $|f(x_n) - f(y_n)| \geq \varepsilon$ .

Then by the compactness of  $A$ , the sequence  $(x_n)$  has a convergent subsequence  $(x_{n_k})$  such that  $a := \lim_k x_{n_k}$ . By applying the compactness of  $A$ , the sequence  $(y_{n_k})$  has a convergent subsequence  $(y_{n_{k_i}})$  with  $b := \lim_i y_{n_{k_i}} \in A$ . Note that we still have  $a := \lim_i x_{n_{k_i}}$ . Since  $|x_{n_{k_i}} - y_{n_{k_i}}| < 1/n_{k_i}$  for all  $i = 1, 2, \dots$ , we have  $a = b$ . Hence, we have

$$\lim_i f(x_{n_{k_i}}) = f(a) = f(b) = \lim_i f(y_{n_{k_i}}),$$

and so we have

$$0 < \varepsilon \leq |f(x_{n_{k_i}}) - f(y_{n_{k_i}})| \rightarrow 0 \quad \text{as } i \rightarrow \infty.$$

It leads to a contradiction.

**Method II:**

Let  $\varepsilon > 0$ . Let  $a \in A$ . Since  $f$  is continuous at  $a$ , there is  $\delta_a > 0$  so that  $|f(x) - f(a)| < \varepsilon$  as  $x \in A$  and  $|x - a| < \delta_a$ . Put  $J_a := (a - \frac{\delta_a}{2}, a + \frac{\delta_a}{2})$ . Then we have  $A \subseteq \bigcup_{a \in A} J_a$ . By using

the compactness of  $A$ , there are finitely many  $a_1, \dots, a_N \in A$  such that  $A \subseteq \bigcup_{k=1}^N J_{a_k}$ . Take

$0 < \delta < \frac{1}{2}\delta_{a_k}$  for all  $k = 1, \dots, N$ . Let  $x, x' \in A$  with  $|x - x'| < \delta$ . Note that  $x \in J_{a_i}$  for some  $1 \leq i \leq N$ . Then we have  $|x - a_i| < \frac{1}{2}\delta_{a_i} < \delta_{a_i}$  and  $|x' - a_i| \leq |x' - x| + |x - a_i| < \delta_{a_i}$ . Thus, we have

$$|f(x) - f(x')| \leq |f(x) - f(a_i)| + |f(a_i) - f(x')| < 2\varepsilon.$$

The proof is complete. □

**Theorem 9.5 Weierstrass approximation theorem** *Let  $f(x)$  be a real-valued continuous function defined on  $[0, 1]$ . Then for any  $\varepsilon > 0$ , there is a polynomial function  $p(x)$  such that  $|f(x) - p(x)| < \varepsilon$  for all  $x \in [0, 1]$ .*

**Proof:** Before showing the result, let us recall a class of polynomial functions in the following, called *Bernstein polynomials*. For each positive integer  $n$ , set

$$B_n(x) := \sum_{k=0}^n a_k \binom{n}{k} x^k (1-x)^{n-k}, \quad \text{where } a_k \in \mathbb{R}$$

for  $x \in [0, 1]$ . Using the binomial theorem, clearly we have

$$\sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} = 1 \quad (9.1)$$

for all  $x \in [0, 1]$ . Taking  $x(1-x) \frac{d}{dx}$  in Eq 9.1, we have

$$\sum_{k=0}^n \binom{n}{k} [x^k (1-x)^{n-k}] (k - nx) = 0.$$

Again taking  $\frac{1}{n^2} x(1-x) \frac{d}{dx}$  on both side and applying the product rule of derivatives, we have

$$\sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} \left(x - \frac{k}{n}\right)^2 = \frac{x(1-x)}{n}.$$

This gives

$$\sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} \left(x - \frac{k}{n}\right)^2 \leq \frac{1}{4n} \quad \text{for all } x \in [0, 1] \quad (9.2)$$

since  $\max\{x(1-x) : x \in [0, 1]\} = 1/4$ .

Now let

$$p_n(x) := \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} f\left(\frac{k}{n}\right), \quad \text{for } x \in [0, 1].$$

The theorem is obtained if we show that for any  $\varepsilon > 0$ , there is a positive integer  $N$  such that  $|f(x) - p_n(x)| < \varepsilon$  for all  $n \geq N$  and for all  $x \in [0, 1]$ . Note that using the Eq9.1, we see that

$$|f(x) - p_n(x)| \leq \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} |f(x) - f\left(\frac{k}{n}\right)| \quad (9.3)$$

for all  $x \in [0, 1]$  and for all  $n = 1, 2, \dots$ . On the other hand, since  $f$  is continuous on  $[0, 1]$ ,  $f$  is uniformly continuous on  $[0, 1]$ . Thus, for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $|f(u) - f(v)| < \varepsilon$  whenever  $u, v \in [0, 1]$  with  $|u - v| < \delta$ . Now for any  $x \in [0, 1]$  and  $n = 1, 2, \dots$ , we put

$$A := \sum_{k: |x - \frac{k}{n}| < \delta} \binom{n}{k} x^k (1-x)^{n-k} |f(x) - f\left(\frac{k}{n}\right)|$$

and

$$B := \sum_{k: |x - \frac{k}{n}| \geq \delta} \binom{n}{k} x^k (1-x)^{n-k} |f(x) - f\left(\frac{k}{n}\right)|.$$

Then the Eq 9.3 can be written as the following

$$|f(x) - p_n(x)| \leq A + B.$$

Notice that by the choice of  $\delta$ , we have

$$A \leq \varepsilon \sum_{k: |x - \frac{k}{n}| < \delta} \binom{n}{k} x^k (1-x)^{n-k} = \varepsilon.$$

We are now going to estimate the value  $B$ . By using Eq 9.2, we have

$$\delta^2 \sum_{k: |x - \frac{k}{n}| \geq \delta} \binom{n}{k} x^k (1-x)^{n-k} \leq \sum_{k: |x - \frac{k}{n}| \geq \delta} \binom{n}{k} x^k (1-x)^{n-k} \left(x - \frac{k}{n}\right)^2 \leq \frac{1}{4n}.$$

This gives

$$\sum_{k: |x - \frac{k}{n}| \geq \delta} \binom{n}{k} x^k (1-x)^{n-k} \leq \frac{1}{4n\delta^2}.$$

Since  $f$  is continuous,  $f$  is bounded on  $[0, 1]$ . If we put  $M := \sup |f(x)|$ , then we can now conclude that

$$B \leq 2M \sum_{k: |x - \frac{k}{n}| \geq \delta} \binom{n}{k} x^k (1-x)^{n-k} \leq \frac{M}{2n\delta^2}$$

for all  $x \in [0, 1]$  and for all  $n = 1, 2, \dots$ . Therefore, if we choose a positive integer  $N$  so that  $\frac{M}{2n\delta^2} < \varepsilon$  for all  $n \geq N$ , then  $|f(x) - p_n(x)| \leq A + B < 2\varepsilon$  for all  $n \geq N$  and for all  $x \in [0, 1]$  as desired. The proof is complete.  $\square$

**Definition 9.6** Let  $A$  be a non-empty subset of  $\mathbb{R}$ . A function  $f : A \rightarrow \mathbb{R}$  is called a Lipschitz function if there is a constant  $C > 0$  such that  $|f(x) - f(y)| \leq C|x - y|$  for all  $x, y \in A$ . In this case.

Furthermore, if we can find such  $0 < C < 1$ , then we call  $f$  a contraction.

Clearly we have the following property.

**Proposition 9.7** Every Lipschitz function is uniformly continuous on its domain.

**Example 9.8** (i) : The sine function  $f(x) = \sin x$  is a Lipschitz function on  $\mathbb{R}$  since we always have  $|\sin x - \sin y| \leq |x - y|$  for all  $x, y \in \mathbb{R}$ .

(ii) : Define a function  $f$  on  $[0, 1]$  by  $f(x) = x \sin(1/x)$  for  $x \in (0, 1]$  and  $f(0) = 0$ . Then  $f$  is continuous on  $[0, 1]$  and thus  $f$  is uniformly continuous on  $[0, 1]$ , but note that  $f$  is not a Lipschitz function. In fact, for any  $C > 0$ , if we consider  $x_n = \frac{1}{2n\pi + (\pi/2)}$  and  $y_n = \frac{1}{2n\pi}$ , then  $|f(x_n) - f(y_n)| > C|x_n - y_n|$  if and only if

$$\frac{2}{\pi} \cdot \frac{(2n\pi + \frac{\pi}{2})(2n\pi)}{2n\pi + \frac{\pi}{2}} = 4n > C.$$

Therefore, for any  $C > 0$ , there are  $x, y \in [0, 1]$  such that  $|f(x) - f(y)| > C|x - y|$  and hence  $f$  is not a Lipschitz function on  $[0, 1]$ .

**Proposition 9.9** Let  $A$  be a non-empty closed subset of  $\mathbb{R}$ . If  $f : A \rightarrow A$  is a contraction, then there is a unique fixed point of  $f$ , i.e., there is a point  $a \in A$  such that  $f(a) = a$ .

**Proof:** First we show the existence.  $f$  is a contraction on  $A$ , so there is  $0 < C < 1$  such that  $|f(x) - f(y)| \leq C|x - y|$  for all  $x, y \in A$ . Fix  $x_1 \in A$ . Since  $f(A) \subseteq A$ , we can inductively define a sequence  $(x_n)$  in  $A$  by  $x_{n+1} = f(x_n)$  for  $n = 1, 2, \dots$ . Note that we have

$$|x_{n+1} - x_n| = |f(x_n) - f(x_{n-1})| \leq C|x_n - x_{n-1}|$$

for all  $n = 2, 3, \dots$ . This gives

$$|x_{n+1} - x_n| \leq C^{n-1}|x_2 - x_1|$$

for  $n = 2, 3, \dots$ . Thus, for any  $n, p = 1, 2, \dots$ , we see that

$$|x_{n+p} - x_n| \leq \sum_{i=n}^{n+p-1} |x_{i+1} - x_i| \leq |x_2 - x_1| \sum_{i=n}^{n+p-1} C^{i-1}.$$

Since  $0 < C < 1$ , for any  $\varepsilon > 0$ , there is  $N$  such that  $\sum_{i=n}^{n+p-1} C^{i-1} < \varepsilon$  for all  $n \geq N$  and  $p = 1, 2, \dots$ . Therefore,  $(x_n)$  is a Cauchy sequence and thus the limit  $a := \lim_n x_n$  exists.  $A$  is closed, so we have  $a \in A$  and hence  $f$  is continuous at  $a$ . On the other hand, since  $x_{n+1} = f(x_n)$ , we have  $a = f(a)$  by taking  $n \rightarrow \infty$ .

Finally, we show the uniqueness of the fixed point. In fact, if  $a$  and  $b$  are the fixed points of  $f$  and  $a \neq b$ , then we have  $|a - b| = |f(a) - f(b)| \leq C|a - b| < |a - b|$  because  $0 < C < 1$ . It leads to a contradiction. The proof is complete.  $\square$

**Remark 9.10** Proposition 9.9 does not hold if  $f$  is not a contraction. For example, if we consider  $f(x) = x - 1$  for  $x \in \mathbb{R}$ , clearly we have  $|f(x) - f(y)| = |x - y|$  and  $f$  has no fixed point in  $\mathbb{R}$ .

**Proposition 9.11** Let  $f$  be a continuous function defined on  $(a, b)$ . The the followings are equivalent.

- (i) There exists a continuous function  $F : [a, b] \rightarrow \mathbb{R}$  such that  $F(x) = f(x)$  for all  $x \in (a, b)$ .
- (ii)  $f$  is uniformly continuous on  $(a, b)$ .
- (iii) The limits  $\lim_{x \rightarrow a^+} f(x)$  and  $\lim_{x \rightarrow b^-} f(x)$  both exist.

In this case, this continuous extension  $F$  is uniquely determined by  $f$ . In fact,  $F(a) = \lim_{x \rightarrow a^+} f(x)$  and  $F(b) = \lim_{x \rightarrow b^-} f(x)$ .

**Proof:** For (i)  $\Rightarrow$  (ii), we assume that (i) holds. Then by Theorem 9.4,  $F$  is uniformly continuous on  $[a, b]$ , so  $f = F|_{(a, b)}$  is uniformly continuous on  $(a, b)$ .

For (ii)  $\Rightarrow$  (iii), we are going to show that  $\lim_{x \rightarrow b^-} f(x)$  exists.

It suffices to show that the sequence  $(f(x_n))$  converges to the same limit whenever any sequence  $(x_n)$  in  $(a, b)$  that converges to  $b$ .

First, we claim that  $(f(x_n))$  is a Cauchy sequence for any such sequence  $(x_n)$  in  $(a, b)$ . Let  $\varepsilon > 0$ . Then by the assumption (ii), there is  $\delta > 0$  such that  $|f(x) - f(y)| < \varepsilon$  as



$x, y \in (a, b)$  with  $|x - y| < \delta$ . Now since  $\lim x_n = b$  and thus  $(x_n)$  is a Cauchy sequence. Therefore, we can find a positive  $N$  such that  $|x_m - x_n| < \delta$  when  $m, n \geq N$ . This gives  $|f(x_m) - f(x_n)| < \varepsilon$  as  $m, n \geq N$ . The claim follows and thus, the limit  $\lim_{n \rightarrow \infty} f(x_n)$  exists. Next we want to show that if  $(x_n)$  and  $(y_n)$  both are the sequences in  $(a, b)$  that converge to  $b$ , then  $\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} f(y_n)$ . Let  $L = \lim_{n \rightarrow \infty} f(x_n)$  and  $L' = \lim_{n \rightarrow \infty} f(y_n)$ . Let  $\varepsilon > 0$  and let  $\delta$  be given by the uniform continuity of  $f$ . Since  $\lim x_n = \lim y_n$ , we can choose a positive integer  $N$  large enough so that  $|x_N - y_N| < \delta$ . In addition, such  $N$  satisfies  $|f(x_N) - L| < \varepsilon$  and  $|f(y_N) - L'| < \varepsilon$  because  $L = \lim_{n \rightarrow \infty} f(x_n)$  and  $L' = \lim_{n \rightarrow \infty} f(y_n)$ . This implies that

$$|L - L'| \leq |L - f(x_N)| + |f(x_N) - f(y_N)| + |f(y_N) - L'| < 3\varepsilon$$

for all  $\varepsilon > 0$ . Thus,  $L = L'$  and hence, the limit  $\lim_{x \rightarrow b^-} f(x)$  exists.

The proof of the case  $\lim_{x \rightarrow a^+} f(x)$  is similar.

Finally, we show (iii)  $\Rightarrow$  (i). Define  $F(a) := \lim_{x \rightarrow a^+} f(x)$ ;  $F(b) := \lim_{x \rightarrow b^-} f(x)$  and  $F(x) := f(x)$  for  $x \in (a, b)$ . Note that  $F$  is continuous on  $[a, b]$ . In fact, we have  $F(a) = \lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^+} F(x)$  and  $F(b) = \lim_{x \rightarrow b^-} f(x) = \lim_{x \rightarrow b^-} F(x)$ . Thus,  $F$  is continuous at  $x = a$  and  $b$ .

The last assertion follows immediately from the continuity of  $F$ . The proof is complete.  $\square$

**Remark 9.12** Indeed, in the proof of Proposition 9.11 (i)  $\Rightarrow$  (ii) above, we have shown the following fact. Suppose that  $f$  is uniformly continuous function defined on  $A$ . If  $(x_n)$  is a Cauchy sequence in  $A$ , then so is the sequence  $(f(x_n))$ . We can use this simple observation to see a function "NOT" being uniformly continuous on its domain.

Note the assumption of the uniform continuity of  $f$  is essential in here by considering the simple example that  $f(x) = \frac{1}{x}$ ,  $x \in A := (0, 1]$  and  $x_n = \frac{1}{n}$ ,  $n = 1, 2, \dots$

**Definition 9.13** A function  $s : [a, b] \rightarrow \mathbb{R}$  is called a step function (resp. piecewise linear) if there exist finitely many points  $a = x_0 < x_1 < \dots < x_n = b$  such that  $s$  is a constant on each  $(x_{k-1}, x_k)$  (resp. linear on  $[x_{k-1}, x_k]$ , i.e.  $s(x) = m_k x + b_k$ ) for all  $k = 1, \dots, n$ .

**Proposition 9.14** If  $f$  is a continuous function defined on a closed and bounded interval  $[a, b]$ , then it can be uniformly approximated by step functions (resp. piecewise linear functions), that is, for each  $\varepsilon > 0$ , there exists a step function  $s$  (resp. piecewise linear function) defined on  $[a, b]$  such that  $|f(x) - s(x)| < \varepsilon$  for all  $x \in [a, b]$ .

**Proof:** By using Theorem 9.4, we first note that  $f$  is uniformly continuous on  $[a, b]$ . Let  $\varepsilon > 0$ . Then there is  $\delta > 0$  so that  $|f(x) - f(y)| < \varepsilon$  whenever  $x, y \in (a, b)$  with  $|x - y| < \delta$ . If we choose a partition  $a = x_0 < \dots < x_n = b$  on  $[a, b]$  such that  $|x_k - x_{k-1}| < \delta$  for  $k = 1, \dots, n$ . Now if we let  $s(x) := f(x_{k-1})$  when  $x \in [x_{k-1}, x_k)$ , then  $s$  is the step function as desired. Using the similar argument, the result is obtained for the case of piecewise linear functions.  $\square$

## 10 Monotone Functions

Using the notation given as before,  $f$  is a function defined on a subset  $A$  of  $\mathbb{R}$ .  $f$  is called a *monotone function* if it is either increasing or decreasing. The following results also hold for decreasing functions by considering  $-f$  instead. Recall that  $c$  is a *right (resp. left) limit point* of  $A$  if for any  $r > 0$  we have  $(c, c+r) \cap A \neq \emptyset$  (resp.  $(c-r, c) \cap A \neq \emptyset$ ).

**Proposition 10.1** *Let  $f$  be an increasing function on  $A$ . Let  $c \in A$ . Put*

$$L(c) := \inf\{f(x) : x \in A, x > c\} \quad \text{if } \{x \in A, x > c\} \neq \emptyset.$$

*Similarly, we put*

$$l(c) := \sup\{f(x) : x \in A, x < c\} \quad \text{if } \{x \in A : x < c\} \neq \emptyset.$$

*If  $c$  is a right (resp. left) limit point of  $A$ , then  $L(c) = f(c+) := \lim_{x \rightarrow c+} f(x)$  (resp.  $l(c) = f(c-) := \lim_{x \rightarrow c-} f(x)$ ).*

**Proof:** First, we want to prove that if  $c$  is a right limit point of  $A$ , then the right limit  $f(c+)$  exists. Since  $c$  is a right limit point of  $A$ ,  $\{f(x) : x \in A, x > c\} \neq \emptyset$ .  $f$  is increasing, so  $f(c)$  is a lower bound of the set  $\{f(x) : x \in A, x > c\}$ . The Axiom of Completeness implies that  $L(c) := \inf\{f(x) : x \in A, x > c\}$  exists and  $f(c) \leq L(c)$ . Thus, for any  $\varepsilon > 0$ , there is  $x_1 \in A$  with  $x_1 > c$  such that  $f(x_1) < L(c) + \varepsilon$ . Hence, if we take  $0 < \delta < x_1 - c$ , then  $L(c) - \varepsilon < L(c) \leq f(x) \leq f(x_1) < L(c) + \varepsilon$  whenever  $x \in (c, c + \delta)$ . Thus,  $L(c) = f(c+)$  as desired.

The proof for the case of a left limit point is similar. □

**Proposition 10.2** *Using the notation given as in Proposition 10.1, let  $f$  be a strictly increasing (not necessary continuous) function defined on an interval  $I$ , i.e.  $f(x_1) < f(x_2)$  if and only if  $x_1 < x_2$  as  $x_1, x_2 \in I$ . Let  $g : f(I) \rightarrow I$  be the inverse of  $f$ . If  $d \in f(I)$ , then  $g(d) = L(d) := \inf\{g(y) : y \in f(I), y > d\}$  (resp.  $g(d) = l(d) := \sup\{g(y) : y \in f(I), y > d\}$ ) provided  $L(d)$  (resp.  $l(d)$ ) exists.*

*In addition, if  $d$  is a right (resp. left) limit point of  $f(I)$ , then  $g(d) = g(d+)$  (resp.  $g(d) = g(d-)$ ).*

*Consequently, the inverse function  $g : f(I) \rightarrow I$  is continuous.*

**Proof:**

Note that  $g$  is also strictly increasing on  $f(I)$ . Let  $c := g(d)$ , hence  $c \in I$  and  $f(c) = d$ .  $g$  is increasing, so  $g(d) \leq L(d)$  whenever  $L(d)$  exists. We now suppose that  $g(d) < L(d)$ , thus we can choose a point  $z$  such that  $c = g(d) < z < L(d)$ . Then by the definition of  $L(d)$ , there is  $y_1 \in f(I)$  with  $y_1 > d$ . Thus, we have  $z < L(d) \leq g(y_1)$ . If we let  $x_1 = g(y_1)$ , then  $x_1 \in I$  and  $c < z < L(d) \leq x_1$ .  $I$  is an interval, so  $z \in (c, x_1) \subseteq I$ . Thus,  $f(z) > f(c) = d$ , so  $f(z) \in \{y \in f(I) : y > d\}$ . This implies that  $z = g(f(z)) \geq L(d)$ . It leads to a contradiction because  $c < z < L(d)$  by the choice of  $z$ . Therefore,  $g(d) = L(d)$ .

Similarly, we also have a contradiction if  $l(d) < g(d)$ . Hence  $l(d) = g(d)$ .

Finally, we want to show that  $g$  is continuous at  $d$  in the following cases.

If  $d$  is an isolated point of  $f(I)$ , then  $g$  is automatically continuous at  $d$ .

If  $d$  is a right limit point of  $f(I)$  but is not a left limit point of  $f(I)$ , then by Proposition 10.1, we have  $g(d) = L(d) = g(d+)$ . Therefore,  $g$  is continuous at  $d$ . Similarly, if  $d$  is a left

limit point of  $f(I)$  but is not a right limit point of  $f(I)$ , then we have  $g(d) = l(d) = g(d-)$ , hence  $g$  is continuous at  $d$ .

Finally, if  $d$  is a right and left limit point of  $f(I)$ . Then, we have  $g(d) = g(d+) = g(d-)$  and so  $g$  is continuous at  $d$ . The proof is complete.  $\square$

**Proposition 10.3** *Let  $f$  be an increasing function defined on  $A$  and let  $D$  be the set of discontinuous points of  $f$ . Then  $D$  is a countable set.*

**Proof:** For each integer  $n$ , we put  $D_n := \{x \in D : n-1 \leq f(x) \leq n\}$ . Then  $D = \bigcup_{n \in \mathbb{Z}} D_n$ . Therefore, it suffices to show that each  $D_n$  is countable.

We now fix  $D_m$ . By using Proposition 10.1, we first note that  $c \in D_m$  if and only if  $f(c) - f(c-) > 0$  or  $f(c+) - f(c) > 0$ . Put  $J(c-) := [f(c-), f(c)]$  and  $J(c+) := [f(c), f(c+)]$ . Then  $J(c+)$  or  $J(c-)$  is an interval. Therefore, if we put  $\alpha(c)$  is the length of  $(J(c-) \cup J(c+))$  for  $c \in D_m$ , then  $\alpha(c) > 0$ . On the other hand, if  $c_1, c_2 \in D_m$  with  $c_1 < c_2$ , then  $J(c_1+) \cap J(c_2-)$  has at most one point if they exist. Thus, we have

$$0 < \sum_{c \in D_m} \alpha(c) \leq m - (m-1) = 1.$$

Since  $\alpha(c) > 0$  for all  $c \in D_m$ , the set  $D_m$  need to be countable. In fact, note that we have

$$D_m = \bigcup_{k \in \mathbb{Z}_+} \{c \in D_m : \alpha(c) \geq 1/k\}.$$

Thus, if  $D_m$  is uncountable, then there exists a positive integer  $k$  so that  $R := \{c \in D_m : \alpha(c) \geq 1/k\}$  is infinite. Therefore,  $\sum_{c \in R} \alpha(c)$  is infinite. It leads to a contradiction.  $\square$

Let  $I$  be an interval. We call a function  $f : I \rightarrow \mathbb{R}$  *locally bounded* at  $c \in I$  if there is  $r > 0$  such that the function  $f$  is bounded on  $(c-r, c+r) \cap I$ . Notice that if  $\lim_{x \rightarrow c} f(x)$  exists, then  $f$  is locally bounded at  $c$ .

**Proposition 10.4** *Now if  $f$  is a locally bounded at  $c \in I$ , then the following always exist.*

$$\overline{\lim}_{x \rightarrow c} f(x) := \lim_{\delta \rightarrow 0^+} \left( \sup_{0 < |x-c| < \delta} f(x) \right) \quad \text{and} \quad \underline{\lim}_{x \rightarrow c} f(x) := \lim_{\delta \rightarrow 0^+} \left( \inf_{0 < |x-c| < \delta} f(x) \right).$$

Moreover, we have  $\underline{\lim}_{x \rightarrow c} f(x) \leq \overline{\lim}_{x \rightarrow c} f(x)$ .

**Proof:** For simply, we assume that  $I$  is an open interval.

For  $\delta > 0$ , put  $L(\delta) := \sup_{0 < |x-c| < \delta} f(x)$  and  $\ell(\delta) := \inf_{0 < |x-c| < \delta} f(x)$ . Since  $f$  is locally bounded at  $c$ ,  $L(\delta)$  and  $\ell(\delta)$  are defined for arbitrary small  $\delta > 0$ . Moreover, the functions  $L(\delta)$  is increasing and  $\ell(\delta)$  is decreasing. Then by Proposition 10.1, we see that  $\overline{\lim}_{x \rightarrow c} f(x) = \inf_{\delta > 0} ( \sup_{0 < |x-c| < \delta} f(x) )$  and  $\underline{\lim}_{x \rightarrow c} f(x) = \sup_{\delta > 0} ( \inf_{0 < |x-c| < \delta} f(x) )$ .

On the other hand, since  $\ell(\delta) \leq L(\delta)$  for all  $\delta > 0$ , we have  $\underline{\lim}_{x \rightarrow c} f(x) \leq \overline{\lim}_{x \rightarrow c} f(x)$ .  $\square$

**Example 10.5** *Define  $f(x) = \sin \frac{1}{x}$  for  $x \neq 0$ . Then  $\overline{\lim}_{x \rightarrow 0} f(x) = 1$  and  $\underline{\lim}_{x \rightarrow 0} f(x) = -1$ . Hence, the case  $\underline{\lim}_{x \rightarrow 0} f(x) < \overline{\lim}_{x \rightarrow 0} f(x)$  may occur.*

**Proposition 10.6** *Using the notation as in Proposition 10.4, the following are equivalent.*

(i)  $\lim_{x \rightarrow c} f(x)$  exists.

(ii)  $\underline{\lim}_{x \rightarrow c} f(x) = \overline{\lim}_{x \rightarrow c} f(x)$ .

In this case, we have  $\lim_{x \rightarrow c} f(x) = \underline{\lim}_{x \rightarrow c} f(x) = \overline{\lim}_{x \rightarrow c} f(x)$ .

**Proof:** Let  $L := \lim_{x \rightarrow c} f(x)$  if it exists.

For showing (i)  $\Rightarrow$  (ii), let  $\varepsilon > 0$ , then there is  $\delta > 0$  such that  $L - \varepsilon < f(x) < L + \varepsilon$  whenever  $x \in I$  with  $0 < |x - c| < \delta$ . Using the notation as in Proposition 10.4, then we have  $L - \varepsilon \leq \ell(\delta_1) \leq L(\delta_1) \leq L + \varepsilon$  for all  $0 < \delta_1 < \delta$ . Taking  $\delta_1 \rightarrow 0+$ , we have

$$L - \varepsilon \leq \underline{\lim}_{x \rightarrow c} f(x) \leq \overline{\lim}_{x \rightarrow c} f(x) \leq L + \varepsilon$$

for all  $\varepsilon > 0$ . This gives  $\underline{\lim}_{x \rightarrow c} f(x) = \overline{\lim}_{x \rightarrow c} f(x) = L$  as desired.

(ii)  $\Rightarrow$  (i) follows immediately from the simple observation that  $\ell(\delta) \leq f(x) \leq L(\delta)$  for all  $\delta > 0$  and for all  $x \in I$  with  $0 < |x - c| < \delta$ .  $\square$

**Definition 10.7** *A function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be of bounded variation if it satisfies the following condition.*

$$\|f\|_{BV} := \sup \left\{ \sum_{k=1}^n |f(x_k) - f(x_{k-1})| : a = x_0 < \dots < x_n = b \right\} < \infty.$$

We call  $\|f\|_{BV}$  the total variation of  $f$  and write  $BV[a, b]$  for the set of all functions of bounded variation.

**Proposition 10.8** *We keep all notation as given above. Then we have*

(i) *The space  $BV[a, b]$  is a vector space.*

(ii) *Every function of bounded variation is bounded.*

(iii) *Every monotone function is of bounded variation.*

(iv) *Every Lipschitz function is of bounded variation.*

**Proof:** Parts (i), (iii) and (iv) are clear.

For showing Part (ii), let  $f \in BV[a, b]$  and  $x \in (a, b)$ . Then by considering the partition  $a < x < b$ , we have  $|f(x) - f(a)| + |f(b) - f(x)| \leq \|f\|_{BV}$ . This gives  $2|f(x)| \leq |f(a)| + |f(b)| + \|f\|_{BV}$  for all  $x \in (a, b)$  and hence,  $f$  is bounded on  $[a, b]$ .  $\square$

**Remark 10.9** The part (iii) in Proposition 10.8 tells us that a function of bounded variation may not be continuous. On the other hand, a bounded function is not necessary to be of bounded variation.

**Example 10.10** *Let*

$$f(x) = \begin{cases} \sin(\frac{1}{x}) & \text{if } x \in (0, \frac{2}{\pi}]; \\ 0 & \text{if } x = 0. \end{cases}$$

*Notice that  $f$  is bounded on  $[0, \frac{2}{\pi}]$  but  $f \notin BV[0, \frac{2}{\pi}]$ . In fact, for each positive integer  $n$ , if we let  $x_k := \frac{1}{\frac{\pi}{2} + (n-k)\pi}$  for  $1 \leq k \leq n$  and consider the partition on  $[0, \frac{2}{\pi}] : 0 = x_0 < x_1 < \dots < x_{n-1} < x_n = \frac{2}{\pi}$ , then we have*

$$\sum_{k=1}^n |f(x_k) - f(x_{k-1})| \geq \sum_{k=2}^n |f(x_k) - f(x_{k-1})| \geq 2(n-1)$$

*for all  $n \geq 2$ . This implies that  $f$  is not of bounded variation.*

**Proposition 10.11** *Suppose that  $f \in BV[a, b]$ . Put*

$$\phi(x) = \begin{cases} \sup\{\sum_{k=1}^n |f(x_k) - f(x_{k-1})| : a = x_0 < \dots < x_n = x\} & \text{if } x \in (a, b]; \\ 0 & \text{if } x = a. \end{cases}$$

*Then the functions  $\phi$  and  $\phi - f$  are increasing on  $[a, b]$ . Consequently, a function  $f : [a, b] \rightarrow \mathbb{R}$  is of bounded variation if and only if  $f = u - v$  for some increasing functions  $u$  and  $v$  on  $[a, b]$ .*

**Proof:** We first notice that since  $f \in BV[a, b]$ , the function  $\phi$  is well defined. In fact, we have  $0 \leq \phi(x) \leq \|f\|_{BV}$  for all  $x \in [a, b]$ . Let  $a < x < y \leq b$ . For any partition on  $[a, x] : a = x_0 < \dots < x_n = x$ , we see that

$$\sum_{k=1}^n |f(x_k) - f(x_{k-1})| \leq \sum_{k=1}^n |f(x_k) - f(x_{k-1})| + |f(y) - f(x)| \leq \phi(y).$$

This implies that  $\phi(x) \leq \phi(y)$  and

$$\phi(x) \leq \phi(y) - |f(y) - f(x)| \leq \phi(y) - (f(y) - f(x)).$$

Thus, we have

$$\phi(x) - f(x) \leq \phi(y) - f(y).$$

Therefore, the function  $\phi - f$  is increasing on  $[a, b]$ .

For the last assertion, if  $f \in BV[a, b]$ , then we have  $f = \phi - (\phi - f)$  and so  $f$  can be written as the difference of increasing functions as desired. Conversely, since  $BV[a, b]$  is a vector space and every increasing function is of bounded variation, the difference of increasing functions is a function of bounded variation. The proof is complete.  $\square$

## References

- [1] R.G. Bartle and R. Sherbert, Introduction to real analysis, 4th edition. John Wiley & Sons, Inc. (2011).