

Lecture 1:

Image transformation

Let \mathcal{I} = Collection of images of size N and range of intensity $[0, M]$.

$$= \{ f \in M_{N \times N}(\mathbb{R}) : 0 \leq f(i,j) \leq M ; 1 \leq i, j \leq N \}$$

(for simplicity, assume f is a square image; can be general $N_1 \times N_2$ image)

Image transformation = $\mathcal{O} : \mathcal{I} \rightarrow \mathcal{I}$ (transform one image to another)

Imaging problems

① Find a suitable transformation $T \Rightarrow g := T(f)$ becomes good!

② Given a ^{noisy image} g and transformation T , find original clean image f .

$$g = T(f) + n \quad \begin{matrix} \uparrow & \uparrow & \leftarrow \\ \text{Known} & \text{Known} & \text{Unknown} \end{matrix} \quad \begin{matrix} \uparrow & \uparrow & \leftarrow \\ \text{Unknown} & \text{Unknown} & \text{Unknown} \end{matrix} \quad (\text{Inverse problem})$$

Definition: (Linear image transformation)

$\mathcal{O}: \mathcal{I} \rightarrow \mathcal{I}$ is linear $\Leftrightarrow \mathcal{O}(af + g) = a\mathcal{O}(f) + \mathcal{O}(g)$ for $\forall f, g \in \mathcal{I}; \forall a \in \mathbb{R}$

Take $f \in \mathcal{I}$. Let

$$f = \begin{pmatrix} f(1,1) & \dots & f(1,N) \\ f(2,1) & \dots & f(2,N) \\ \vdots & \ddots & \vdots \\ f(i,j) & \dots & \vdots \\ f(N,1) & \dots & f(N,N) \end{pmatrix} = \sum_{i=1}^N \sum_{j=1}^N \begin{pmatrix} 0 & \dots & 0 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & f(i,j) & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 0 & \dots & 0 \end{pmatrix} = \sum_{i=1}^N \sum_{j=1}^N f(x,y) \begin{pmatrix} 0 & \dots & 0 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 1 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 0 & \dots & 0 \end{pmatrix}$$

$\downarrow y$ $\leftarrow x$

Let $g = \mathcal{O}(f)$. Assume \mathcal{O} is linear, then:

$$g(\alpha, \beta) = \sum_{x=1}^N \sum_{y=1}^N f(x,y) \left[\mathcal{O} \left(\begin{pmatrix} 0 & \dots & 0 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 1 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 0 & \dots & 0 \end{pmatrix} \right) \right]_{\alpha, \beta}$$

$$= \sum_{x=1}^N \sum_{y=1}^N f(x,y) h(x, \alpha, y, \beta) \quad \text{where}$$

$$h(x, \alpha, y, \beta) = [\mathcal{O}(P_{xy})]_{\alpha, \beta}; P_{xy} = \begin{pmatrix} 0 & \dots & 0 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 1 & \dots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \dots & 0 & \dots & 0 \end{pmatrix} \leftarrow x^{th} \quad \downarrow y^{th}$$

Remark: $h(x, \alpha, y, \beta)$ = how much input value at (x, y) influence the output value at (α, β) .



Pixel (x, y) affecting pixel (α, β) by a weight $h(x, \alpha, y, \beta)$.

Definition: (Point spread function)

$h(\cdot, \alpha, \cdot, \beta)$ is called the PSF at (α, β) .

Fix α, β . Let x, y as variables!

Definition: (Shift-invariant)

A PSF is shift invariant if:

$$h(x, \alpha, y, \beta) = h(\alpha - x, \beta - y) \text{ for } \forall 1 \leq x, y, \alpha, \beta \leq N$$

Definition: (Convolution) Let $f, g \in \mathcal{I}$.

Convolution of f and g is defined as $f * g(\alpha, \beta) = \sum_{x=1}^N \sum_{y=1}^N f(x, y) g(\alpha-x, \beta-y)$

(Assume f and g are periodically extended: $\begin{cases} f(x+iN, y+jN) = f(x, y) \\ g(x+iN, y+jN) = g(x, y) \end{cases} \forall i, j \in \mathbb{Z}$)

Theorem: If a PSF is shift-invariant, then the operator \mathcal{O} is a convolution with the input image.

Proof: Let $g := \mathcal{O}(f)$.
$$\begin{aligned} g(\alpha, \beta) &= \sum_{x=1}^N \sum_{y=1}^N f(x, y) \underbrace{h(x, \alpha, y, \beta)}_{h(\alpha-x, \beta-y)} \\ &= f * h(\alpha, \beta) \end{aligned}$$

Remark:

- $f * h = h * f$ (exercise)
- Convolution is important for understanding image blur.

Definition: (Separable)

A PSF is Separable if: $h(x, \alpha, y, \beta) = h_c(x, \alpha) h_r(y, \beta)$ for $\forall 1 \leq x, y, \alpha, \beta \leq N$.

Theorem: Suppose PSF is separable. Then:

The operator \mathcal{O} consists of two matrix multiplication.

Proof:

$$g(\alpha, \beta) = \sum_{x=1}^N \sum_{y=1}^N f(x, y) h(x, \alpha, y, \beta)$$

Recall: $(AB)_{ij} = \sum_{k=1}^N A_{ik} B_{kj}$

$$= \sum_{x=1}^N \sum_{y=1}^N f(x, y) h_c(x, \alpha) h_r(y, \beta)$$

$$= \sum_{x=1}^N h_c(x, \alpha) \left(\sum_{y=1}^N f(x, y) h_r(y, \beta) \right)$$

matrix multiplication matrix multiplication