

# MATH2068 MATHEMATICAL ANALYSIS II (2021-22)

CHI-WAI LEUNG

## 1. DIFFERENTIATION

Throughout this section, let  $I$  be an open interval (not necessarily bounded) and let  $f$  be a real-valued function defined on  $I$ .

**Definition 1.1.** Let  $c \in I$ . We say that  $f$  is differentiable at  $c$  if the following limit exists:

$$\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}.$$

In this case, we write  $f'(c)$  for the above limit and we call it the derivative of  $f$  at  $c$ . We say that if  $f$  is differentiable on  $I$  if  $f'(x)$  exists for every point  $x$  in  $I$ .

**Proposition 1.2.** Let  $c \in I$ . Then  $f'(c)$  exists if and only if there is a function  $\varphi$  defined on  $I$  such that the function  $\varphi$  is continuous at  $c$  and

$$f(x) - f(c) = \varphi(x)(x - c)$$

for all  $x \in I$ .

In this case,  $\varphi(c) = f'(c)$ .

*Proof.* Assume that  $f'(c)$  exists. Define a function  $\varphi : I \rightarrow \mathbb{R}$  by

$$\varphi(x) = \begin{cases} \frac{f(x) - f(c)}{x - c} & \text{if } x \neq c; \\ f'(c) & \text{if } x = c. \end{cases}$$

Clearly, we have  $f(x) - f(c) = \varphi(x)(x - c)$  for all  $x \in I$ . We want to show that the function  $\varphi$  is continuous at  $c$ . In fact, let  $\varepsilon > 0$ , by the definition of the limit of a function, there is  $\delta > 0$  such that

$$\left| f'(c) - \frac{f(x) - f(c)}{x - c} \right| < \varepsilon$$

whenever  $x \in I$  with  $0 < |x - c| < \delta$ . Therefore, we have  $|f'(c) - \varphi(x)| < \varepsilon$  as  $x \in I$  with  $0 < |x - c| < \delta$ . Since  $\varphi(c) = f'(c)$ , we have  $|f'(c) - \varphi(x)| < \varepsilon$  as  $x \in I$  with  $|x - c| < \delta$ , hence the function  $\varphi$  is continuous at  $c$  as desired.

The converse is clear since  $\varphi(x) = \frac{f(x) - f(c)}{x - c}$  if  $x \neq c$ . The proof is complete.  $\square$

**Proposition 1.3.** Using the notation as above, if  $f$  is differentiable at  $c$ , then  $f$  is continuous at  $c$ .

*Proof.* By using Proposition 1.2, if  $f'(c)$  exists, then there is a function  $\varphi$  defined on  $I$  such that the function  $\varphi$  is continuous at  $c$  and we have  $f(x) - f(c) = \varphi(x)(x - c)$  for all  $x \in I$ . This implies that  $\lim_{x \rightarrow c} f(x) = f(c)$ , so  $f$  is continuous at  $c$  as desired.  $\square$

**Remark 1.4.** In general, the converse of Proposition 1.3 does not hold, for example, the function  $f(x) := |x|$  is a continuous function on  $\mathbb{R}$  but  $f'(0)$  does not exist.

**Proposition 1.5.** *Let  $f$  and  $g$  be the functions defined on  $I$ . Assume that  $f$  and  $g$  both are differentiable at  $c \in I$ . We have the following assertions.*

- (i)  $(f + g)'(c)$  exists and  $(f + g)'(c) = f'(c) + g'(c)$ .
- (ii) The product  $(f \cdot g)'(c)$  exists and  $(f \cdot g)'(c) = f'(c)g(c) + f(c)g'(c)$ .
- (iii) If  $g(c) \neq 0$ , then we have  $(\frac{f}{g})'(c)$  exists and  $(\frac{f}{g})'(c) = \frac{f'(c)g(c) - f(c)g'(c)}{g(c)^2}$ .

*Proof.* Part (i) clearly follows from the definition of the limit of a function.

For showing Part (ii), note that we have

$$\frac{f(x)g(x) - f(c)g(c)}{x - c} = \frac{f(x) - f(c)}{x - c}g(x) + f(c)\frac{g(x) - g(c)}{x - c}$$

for all  $x \in I$  with  $x \neq c$ . From this, together with Proposition 1.3, Part (ii) follows.

For Part (iii), by using Part (ii), it suffices to show that  $(\frac{1}{g})'(c) = -\frac{g'(c)}{g(c)^2}$ . In fact,  $g'(c)$  exists, so  $g$  is continuous at  $c$ . Since  $g(c) \neq 0$ , there is  $\delta_1 > 0$  so that  $g(x) \neq 0$  for all  $x \in I$  with  $|x - c| < \delta_1$ . Then we have

$$\frac{1}{x - c} \left( \frac{1}{g(x)} - \frac{1}{g(c)} \right) = \frac{1}{x - c} \left( \frac{g(c) - g(x)}{g(x)g(c)} \right)$$

for all  $x \in I$  with  $0 < |x - c| < \delta_1$ . By taking  $x \rightarrow c$ , we see that  $(\frac{1}{g})'(c)$  exists and  $(\frac{1}{g})'(c) = \frac{-g'(c)}{g(c)^2}$ . The proof is complete.  $\square$

**Proposition 1.6. (Chain Rule):** *Let  $f, g$  be functions defined on  $\mathbb{R}$ . Let  $d = f(c)$  for some  $c \in \mathbb{R}$ . Suppose that  $f'(c)$  and  $g'(d)$  exist. Then the derivative of composition  $(g \circ f)'(c)$  exists and  $(g \circ f)'(c) = g'(d)f'(c)$ .*

*Proof.* By using Proposition 1.2, we want to find a function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$g \circ f(x) - g \circ f(c) = \varphi(x)(x - c)$$

for all  $x \in \mathbb{R}$  and the function  $\varphi(x)$  is continuous at  $c$ , and so  $(g \circ f)'(c) = \varphi(c)$ .

Let  $y = f(x)$ . By using Proposition 1.2 again, there is a function and  $\beta(y)$  so that  $g(y) - g(d) = \beta(y)(y - d)$  for all  $y \in \mathbb{R}$  and  $\beta(y)$  is continuous at  $d$ . Similarly, there is a function  $\alpha(x)$  we have  $f(x) - f(c) = \alpha(x)(x - c)$  for all  $x \in \mathbb{R}$  and  $\alpha(x)$  is continuous at  $c$ . These two equations imply that

$$g \circ f(x) - g \circ f(c) = \beta(f(x))(f(x) - f(c)) = \beta(f(x))\alpha(x)(x - c)$$

for all  $x \in \mathbb{R}$ . Let  $\varphi(x) := \beta(f(x)) \cdot \alpha(x)$  for  $x \in \mathbb{R}$ . Since  $\beta(d) = g'(d)$  and  $\alpha(c) = f'(c)$ , we see that  $\varphi(c) = \beta(f(c))\alpha(c) = g'(d)f'(c)$ . It remains to show that the function  $\varphi$  is continuous at  $c$ . In fact,  $f'(c)$  exists, so  $f$  is continuous at  $c$ , and hence the composition  $\beta \circ f(x)$  is continuous at  $c$ . In addition, the function  $\alpha$  is continuous at  $c$ . Therefore, the function  $\varphi := (\beta \circ f) \cdot \alpha$  is continuous at  $c$ , and so  $(g \circ f)'(c)$  exists with  $(g \circ f)'(c) = \varphi(c) = g'(d)f'(c)$ . The proof is complete.  $\square$

**Proposition 1.7.** *Let  $I$  and  $J$  be open intervals. Let  $f$  be a strictly increasing function from  $I$  onto  $J$ . Let  $d = f(c)$  for  $c \in I$ . Assume that  $f'(c)$  exists and the inverse of  $f$ , write  $g := f^{-1}$ , is continuous at  $d$ . If  $f'(c) \neq 0$ , then  $g'(d)$  exists and  $g'(d) = \frac{1}{f'(c)}$ .*

*Proof.* Let  $y = f(x)$ . Note that by using Proposition 1.2, there is a function  $F$  on  $I$  such that  $f(x) - f(c) = F(x)(x - c)$  for all  $x \in I$  and  $F$  is continuous at  $c$  with  $F(c) = f'(c) \neq 0$ .  $F$  is continuous at  $c$ , so there are open intervals  $I_1$  and  $J_1$  such that  $c \in I_1 \subseteq I$  and  $d \in f(I_1) = J_1$ , moreover,  $F(x) \neq 0$  for all  $x \in I_1$ . Note that since  $f(x) - f(c) = F(x)(x - c)$ , we have  $y - d = f(g(y)) - f(g(c)) = F(g(y))(g(y) - g(d))$  for all  $y \in J_1$ . Since  $F(x) \neq 0$  for all  $x \in I_1$ , we have  $g(y) - g(d) = F(g(y))^{-1}(y - d)$  for all  $y \in J_1$ . Note that the function  $F(g(y))^{-1}$  is continuous at  $d$ . Thus,  $g'(d)$  exists and  $g'(d) = F(g(d))^{-1} = \frac{1}{f'(c)}$  as desired.  $\square$

**Definition 1.8.** Let  $D$  be a non-empty subset of  $\mathbb{R}$  and let  $g$  be a real-valued function defined on  $D$ .

(i) We say that  $g$  has an absolute maximum (resp. absolute minimum) at a point  $c \in D$  if  $g(c) \geq g(x)$  (resp.  $g(c) \leq g(x)$ ) for all  $x \in D$ .

In this case,  $c$  is called an absolute extreme point of  $g$ .

(ii) We say that  $g$  has a local maximum (resp. local minimum) at a point  $c \in D$  if there is  $r > 0$  such that  $(c - r, c + r) \subseteq D$  and  $g(c) \geq g(x)$  (resp.  $g(c) \leq g(x)$ ) for all  $x \in (c - r, c + r)$ .

In this case,  $c$  is called a local extreme point of  $g$ .

**Remark 1.9.** Note that an absolute extreme point of a function  $g$  need not be a local extreme point, for example if  $g(x) := x$  for  $x \in [0, 1]$ , then  $g$  has an absolute maximum point at  $x = 1$  of  $g$  but 1 is not a local maximum point of  $g$ .

**Proposition 1.10.** Let  $I$  be an open interval and let  $f$  be a function on  $I$ . Assume that  $f$  has a local extreme point at  $c \in I$  and  $f'(c)$  exists. Then  $f'(c) = 0$ .

*Proof.* Without lost the generality, we may assume that  $f$  has local minimum at  $c$ . Then there is  $r > 0$  such that  $f(x) \geq f(c)$  for  $x \in (c - r, c + r) \subseteq I$ . Since  $f'(c)$  exists, by using Proposition 1.2, there is a function  $\varphi$  defined on  $I$  such that  $f(x) - f(c) = \varphi(x)(x - c)$  for all  $x \in I$  and  $\varphi$  is continuous at  $c$  with  $\varphi(c) = f'(c)$ . Thus, we have  $\varphi(x)(x - c) \geq 0$  for all  $x \in (c - r, c + r)$ . From this we see that  $\varphi(x) \geq 0$  as  $x \in (c, c + r)$ , similarly,  $\varphi(x) \leq 0$  as  $x \in (c - r, c)$ . The function  $\varphi$  is continuous at  $c$ , so  $\varphi(c) = 0$  and hence  $f'(c) = \varphi(c) = 0$  as desired.  $\square$

**Proposition 1.11. Rolle's Theorem:** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a continuous function. Assume that  $f'(x)$  exists for all  $x \in (a, b)$  and  $f(a) = f(b)$ . Then there is a point  $c \in (a, b)$  such that  $f'(c) = 0$ .

*Proof.* Recall a fact that every continuous function defined a compact attains absolute points, that is, there are  $c_1$  and  $c_2$  such that  $f(c_1) = \min_{x \in [a, b]} f(x)$  and  $f(c_2) = \max_{x \in [a, b]} f(x)$ , hence,  $f(c_1) \leq f(x) \leq f(c_2)$  for all  $x \in [a, b]$ . If  $f(c_1) = f(c_2)$ , then  $f(x) \equiv f(c_1) = f(c_2)$  for all  $x \in [a, b]$ , so  $f'(x) \equiv 0$  for all  $x \in (a, b)$ .

Otherwise, suppose that  $f(c_1) < f(c_2)$ . Since  $f(a) = f(b)$ , we have  $c_1 \in (a, b)$  or  $c_2 \in (a, b)$ . We may assume that  $c_1 \in (a, b)$ . Then  $x = c_1$  is a local minimum point of  $f$ . Therefore,  $f'(c_1) = 0$  by using Proposition 1.10.  $\square$

**Theorem 1.12. Main Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is a continuous function and is differentiable on  $(a, b)$ , then there is a point  $c \in (a, b)$  such that  $f(b) - f(a) = f'(c)(b - a)$ .

*Proof.* Define a function  $\varphi : [a, b] \rightarrow \mathbb{R}$  by

$$\varphi(x) = f(x) - f(a) - \frac{f(b) - f(a)}{b - a}(x - a)$$

for  $x \in [a, b]$ . Note that the function  $\varphi$  is continuous on  $[a, b]$  with  $\varphi(a) = \varphi(b) = 0$ , in addition,  $\varphi'(x)$  exists for all  $x \in (a, b)$ . The Rolle's Theorem implies that there is a point  $c \in (a, b)$  such that

$$0 = \varphi'(c) = f'(c) - \frac{f(b) - f(a)}{b - a}.$$

The proof is complete.  $\square$

**Corollary 1.13.** Assume that  $f : [a, b] \rightarrow \mathbb{R}$  is a continuous function and is differentiable on  $(a, b)$ . If  $f' \equiv 0$  on  $(a, b)$ , then  $f$  is a constant function.

*Proof.* Fix any point  $z \in (a, b)$ . Let  $x \in (z, b]$ . By using the Mean Value Theorem, there is a point  $c \in (z, x)$  such that  $f(x) - f(z) = f'(c)(x - z)$ . If  $f' \equiv 0$  on  $(a, b)$ , so  $f(x) = f(z)$  for all  $x \in [z, b]$ . Similarly, we have  $f(x) = f(z)$  for all  $x \in [a, z]$ . The proof is complete.  $\square$

**Definition 1.14.** We call a function  $f$  is a  $C^1$ -function on  $I$  if  $f'(x)$  exists and continuous on  $I$ . In addition, we define the  $n$ -derivatives of  $f$  by  $f^{(n)}(x) := f^{(n-1)}(x)$  for  $n \geq 2$ , provided it exists. In this case, we say that  $f$  is a  $C^n$ -function on  $I$ . In particular, we call  $f$  a  $C^\infty$ -function (or smooth function) if  $f$  is a  $C^n$ -function for all  $n = 1, 2, \dots$

For example, the exponential function  $\exp x$  is a very important example of smooth function on  $\mathbb{R}$ .

**Corollary 1.15. Inverse Mapping Theorem:** Let  $f$  be a  $C^1$ -function on an open interval  $I$  and let  $c \in I$ . Assume that  $f'(c) \neq 0$ . Then there is  $r > 0$  such that the function  $f$  is a strictly monotone function on  $(c - r, c + r) \subseteq I$ . If we let  $J := f(c - r, c + r)$ , then the inverse function  $g := f^{-1} : J \rightarrow (c - r, c + r)$  is also a  $C^1$ -function.

*Proof.* We may assume that  $f'(c) > 0$ .  $f'(x)$  is continuous on  $I$ , so there is  $r > 0$  such that  $f'(x) > 0$  for all  $x \in (c - r, c + r) \subseteq I$ . For any  $x_1$  and  $x_2$  in  $(c - r, c + r)$  with  $x_1 < x_2$ , by using the Mean Value Theorem, we have  $f(x_2) - f(x_1) = f'(v)(x_2 - x_1)$  for some  $v \in (x_1, x_2)$ , and hence  $f(x_2) > f(x_1)$ . Therefore the restriction of  $f$  on  $(c - r, c + r)$  is a strictly increasing function, thus, it is an injection. Let  $J := f((c - r, c + r))$ . Then  $J$  is an interval by the Intermediate Value Theorem. Moreover,  $J$  is an open interval because  $f$  is strictly increasing. Also, if we let  $g = f^{-1}$  on  $J$ , then  $g$  is continuous on  $J$  due to the fact that every continuous bijection on a compact set is a homeomorphism. Therefore, by Proposition 1.7, we see that  $g'(y)$  exists on  $J$  and  $g'(y) = \frac{1}{f'(x)}$  for  $y = f(x)$  and  $x \in (c - r, c + r)$ . Therefore,  $g$  is a  $C^1$  function on  $J$ . The proof is complete.  $\square$

**Proposition 1.16. Cauchy Mean Value Theorem:** Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be continuous functions with  $g(a) \neq g(b)$ . Assume that  $f, g$  are differentiable functions on  $(a, b)$  and  $g'(x) \neq 0$  for all  $x \in (a, b)$ . Then there is a point  $c \in (a, b)$  such that  $\frac{f(b)-f(a)}{g(b)-g(a)} = \frac{f'(c)}{g'(c)}$ .

*Proof.* Define a function  $\psi$  on  $[a, b]$  by  $\psi(x) = f(x) - f(a) - \frac{f(b)-f(a)}{g(b)-g(a)}(g(x) - g(a))$  for  $x \in [a, b]$ . Then by using the similar argument as in the Mean Value Theorem, the result follows.  $\square$

**Theorem 1.17. Lagrange Remainder Theorem:** Let  $f$  be a  $C^{(n+1)}$  function defined on  $(a, b)$ . Let  $x_0 \in (a, b)$ . Then for each  $x \in (a, b)$ , there is a point  $c$  between  $x_0$  and  $x$  such that

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{f^{(n+1)}(c)}{(n+1)!} (x - x_0)^{n+1}.$$

*Proof.* We may assume that  $x_0 < x < b$ . **Case:** We first assume that  $f^{(k)}(x_0) = 0$  for all  $k = 0, 1, \dots, n$ . Put  $g(t) = (t - x_0)^{n+1}$  for  $t \in [x_0, x]$ . Then  $g'(t) = (n+1)(t - x_0)^n$  and  $g(x_0) = 0$ . Then by the Cauchy Mean Value Theorem, there is  $x_1 \in (x_0, x)$  such that  $\frac{f(x)}{g(x)} = \frac{f(x)-f(x_0)}{g(x)-g(x_0)} = \frac{f'(x_1)}{g'(x_1)}$ . Using the same step for  $f'$  and  $g'$  on  $[x_0, x_1]$ , there is  $x_2 \in (x_0, x_1)$  such that  $\frac{f'(x_1)}{g'(x_1)} = \frac{f'(x_1)-f'(x_0)}{g'(x_1)-g'(x_0)} = \frac{f^{(2)}(x_2)}{g^{(2)}(x_2)}$ . To repeat the same step, there are  $x_1, x_2, \dots, x_{n+1}$  in  $(a, b)$  such that  $x_k \in (x_0, x_{k-1})$  for  $k = 1, 2, \dots, n+1$  and

$$\frac{f(x)}{g(x)} = \frac{f'(x_1)}{g'(x_1)} = \dots = \frac{f^{(n+1)}(x_{n+1})}{g^{(n+1)}(x_{n+1})}.$$

In addition, note that  $g^{n+1}(x_{n+1}) = (n+1)!$ . Therefore, we have  $\frac{f(x)}{g(x)} = \frac{f^{(n+1)}(x_{n+1})}{(n+1)!}$ , and hence  $f(x) = \frac{f^{(n+1)}(x_{n+1})}{(n+1)!} (x - x_0)^{n+1}$ . Note  $x_{n+1} \in (x_0, x)$  and thus, the result holds for this case.

For the general case, put  $G(x) = f(x) - \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$  for  $x \in (a, b)$ . Note that we have  $G(x_0) = G'(x_0) = \dots = G^{(n)}(x_0) = 0$ . Then by the Claim above, there is a point  $c \in (x_0, x)$  such that  $G(x) = \frac{G^{(n+1)}(c)}{(n+1)!}$ . Since  $G^{(n+1)}(c) = f^{(n+1)}(c)$ ,  $f(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k + \frac{f^{(n+1)}(c)}{(n+1)!}$ . The proof is complete.  $\square$

**Example 1.18.** Recall that the exponential function  $e^x$  is defined by

$$e^x := \sum_{k=0}^{\infty} \frac{x^k}{k!} := \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{x^k}{k!}$$

for  $x \in \mathbb{R}$ . Note that the above limit always exists for all  $x \in \mathbb{R}$  (shown in the last chapter).

Show that the natural base  $e$  is an irrational number.

Put  $f(x) := e^x$  for  $x \in \mathbb{R}$ . It is a known fact  $f$  is a  $C^\infty$  function and  $f^{(n)}(x) = e^x$  for all  $x \in \mathbb{R}$ . Fix any  $x > 0$ . Then by the Lagrange Theorem, for each positive integer  $n$ , there is  $c_n \in (0, x)$  such that

$$f(x) = \sum_{k=0}^n \frac{x^k}{k!} + \frac{e^{c_n}}{(n+1)!} x^{n+1}.$$

In particular, taking  $x = 1$ , we have

$$0 < \frac{e^{c_n}}{(n+1)!} = e - \sum_{k=0}^n \frac{1}{k!} < \frac{3}{(n+1)!}$$

for all positive integer  $n$ . Now if  $e = p/q$  for some positive integers  $p$  and  $q$ , and thus, we have

$$0 < \frac{p}{q} - \sum_{k=0}^n \frac{1}{k!} < \frac{3}{(n+1)!}$$

for all  $n = 1, 2, \dots$ . Now we can choose  $n$  large enough such that  $(n!)^2 \in \mathbb{N}$ . It leads to a contradiction because we have

$$0 < (n!)^2 \left( \frac{p}{q} - \sum_{k=0}^n \frac{1}{k!} \right) < \frac{3(n!)}{(n+1)!} = \frac{3}{n+1} < 1.$$

Therefore,  $e$  is irrational.

**Proposition 1.19.** Let  $f$  be a  $C^2$  function on an open interval  $I$  and  $x_0 \in I$ . Assume that  $f'(x_0) = 0$ . Then  $f$  has local maximum (resp. local minimum) at  $x_0$  if  $f^{(2)}(x_0) < 0$  (resp.  $f^{(2)}(x_0) > 0$ ).

*Proof.* We assume that  $f^{(2)}(x_0) > 0$ . We want to show that  $x_0$  is a local minimum point of  $f$ . The proof of another case is similar. Note that for any  $x \in I \setminus \{x_0\}$ . Then by the Lagrange Theorem, there is a point  $c$  between  $x_0$  and  $x$  such that

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{1}{2} f^{(2)}(x_0)(x - x_0)^2 = f(x_0) + \frac{1}{2} f^{(2)}(x_0)(x - x_0)^2.$$

$f^{(2)}$  is continuous at  $x_0$  and  $f^{(2)}(x_0) > 0$ , and so there is  $r > 0$  such that  $f^{(2)}(x) > 0$  for all  $x \in (x_0 - r, x_0 + r) \subseteq I$ . Therefore, we have

$$f(x) = f(x_0) + \frac{1}{2} f^{(2)}(x)(x - x_0)^2 \geq f(x_0)$$

for all  $x \in (x_0 - r, x_0 + r)$  and thus,  $x_0$  is a local minimum point of  $f$  as desired.  $\square$

**Proposition 1.20. L'Hospital's Rule:** Let  $f$  and  $g$  be the differentiable functions on  $(a, b)$  and let  $c \in (a, b)$ . Assume that  $f(c) = g(c) = 0$ , in addition,  $g'(x) \neq 0$  and  $g(x) \neq 0$  for all  $x \in (a, b) \setminus \{c\}$ . If the limit  $L := \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$  exists, then so does  $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ , moreover, we have  $L = \lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ .

*Proof.* Fix  $c < x < b$ . Then by the Cauchy Mean Value Theorem, there is a point  $x_1 \in (c, x)$  such that

$$\frac{f(x)}{g(x)} = \frac{f(x) - f(c)}{g(x) - g(c)} = \frac{f'(x_1)}{g'(x_1)}$$

$x_1 \in (c, x)$ , so if  $L := \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$  exists, then  $\lim_{x \rightarrow c^+} \frac{f(x)}{g(x)}$  exists and is equal to  $L$ .

Similarly, we also have  $\lim_{x \rightarrow c^-} \frac{f(x)}{g(x)} = L$ . The proof is finished.  $\square$

**Proposition 1.21.** Let  $f$  be a function on  $(a, b)$  and let  $c \in (a, b)$ .

(i) If  $f'(c)$  exists, then the following limit exists (also called the symmetric derivatives of  $f$  at  $c$ ):

$$f'(c) = \lim_{t \rightarrow 0} \frac{f(c+t) - f(c-t)}{2t}.$$

(ii) If  $f^{(2)}(c)$  exists, then

$$f^{(2)}(c) = \lim_{t \rightarrow 0} \frac{f(c+t) - 2f(c) + f(c-t)}{t^2}.$$

*Proof.* For showing (i), note that we have

$$f'(c) = \lim_{t \rightarrow 0^+} \frac{f(c+t) - f(c)}{t} = \lim_{t \rightarrow 0^-} \frac{f(c+t) - f(c)}{t}.$$

Putting  $t = -s$  into the second equality above, we see that

$$f'(c) = \lim_{s \rightarrow 0^+} \frac{f(c-s) - f(c)}{-s}.$$

To sum up the two equations above, we have

$$f'(c) = \lim_{t \rightarrow 0^+} \frac{f(c+t) - f(c-t)}{2t}.$$

Similarly, we have  $f'(c) = \lim_{t \rightarrow 0^-} \frac{f(c+t) - f(c-t)}{2t}$ . Part (i) follows.

For showing Part (ii), let  $h(t) := f(c+t) - 2f(c) + f(c-t)$  for  $t \in \mathbb{R}$ . Then  $h(0) = 0$  and  $h'(t) = f'(c+t) - f'(c-t)$ . By using the L'Hospital's Rule and Part (i), we have

$$\lim_{t \rightarrow 0} \frac{f(c+t) - 2f(c) + f(c-t)}{t^2} = \lim_{t \rightarrow 0} \frac{h'(t)}{(t^2)'} = \lim_{t \rightarrow 0} \frac{f'(c+t) - f'(c-t)}{2t} = f^{(2)}(c).$$

The proof is complete.  $\square$

**Definition 1.22.** A function  $f$  defined on  $(a, b)$  is said to be convex if for any pair  $a < x_1 < x_2 < b$ , we have

$$f((1-t)x_1 + tx_2) \leq (1-t)f(x_1) + tf(x_2)$$

for all  $t \in [0, 1]$ .

**Proposition 1.23.** Let  $f$  be a  $C^2$  function on  $(a, b)$ . Then  $f$  is a convex function if and only if  $f^{(2)}(x) \geq 0$  for all  $x \in (a, b)$ .

*Proof.* For showing ( $\Rightarrow$ ): assume that  $f$  is a convex function. Fix a point  $c \in (a, b)$ .  $f$  is convex, so we have  $f(c) = f(\frac{1}{2}(c+t) + \frac{1}{2}(c-t)) \leq \frac{1}{2}f(c+t) + \frac{1}{2}f(c-t)$  for all  $t \in \mathbb{R}$  with  $c \pm t \in (a, b)$ . By Proposition 1.21, we have

$$f^{(2)}(c) = \lim_{t \rightarrow 0} \frac{f(c+t) - 2f(c) + f(c-t)}{t^2}.$$

Therefore, we have  $f^{(2)}(c) \geq 0$ .

For ( $\Leftarrow$ ), assume that  $f^{(2)}(x) \geq 0$  for all  $x \in (a, b)$ . Fix  $a < x_1 < x_2 < b$  and  $t \in [0, 1]$ . Let  $c := (1-t)x_1 + tx_2$ . Then by the Lagrange Remainder Theorem, there are points  $z_1 \in (x_1, c)$  and  $z_2 \in (c, x_2)$  such that

$$f(x_2) = f(c) + f'(c)(x_2 - c) + \frac{1}{2}f^{(2)}(z_2)(x_2 - c)^2$$

and

$$f(x_1) = f(c) + f'(c)(x_1 - c) + \frac{1}{2}f^{(2)}(z_1)(x_1 - c)^2.$$

These two equations implies that

$$(1-t)f(x_1) + tf(x_2) = f(c) + (1-t)\frac{1}{2}f^{(2)}(z_1)(x_1 - c)^2 + t\frac{1}{2}f^{(2)}(z_2)(x_2 - c)^2 \geq f(c).$$

since  $f^{(2)}(z_1)$  and  $f^{(2)}(z_2)$  both are non-negative. Thus,  $f$  is convex.  $\square$

**Corollary 1.24.** *Let  $p > 0$ . The function  $f(x) := x^p$  is convex on  $(0, \infty)$  if and only if  $p \geq 1$ .*

*Proof.* Note that  $f^{(2)}(x) = p(p-1)x^{p-2}$  for all  $x > 0$ . Then the result follows immediately from Proposition 1.23.  $\square$

**Proposition 1.25. Netwon's Method:** *Let  $f$  be a continuous real-valued function defined on  $[a, b]$  with  $f(a) < 0 < f(b)$  and  $f(z) = 0$  for some  $z \in (a, b)$ . Assume that  $f$  is a  $C^2$  function on  $(a, b)$  and  $f'(x) \neq 0$  for all  $x \in (a, b)$ . Then there is  $\delta > 0$  with  $J := [z - \delta, z + \delta] \subseteq [a, b]$  which have the following property:*

*if we fix any  $x_1 \in J$  and let*

$$(1.1) \quad x_{n+1} := x_n - \frac{f(x_n)}{f'(x_n)}$$

*for  $n = 1, 2, \dots$ , then we have  $z = \lim x_n$ .*

*Proof.* We first choose  $r > 0$  such that  $[z - r, z + r] \subseteq (a, b)$ . We fix any point  $x_1 \in (z - r, z + r)$  with  $x_1 \neq z$ . Then by the Lagrange Remainder Theorem, there is a point  $\xi$  between  $z$  and  $x_1$  such that

$$0 = f(z) = f(x_1) + f'(x_1)(z - x_1) + \frac{1}{2}f^{(2)}(\xi)(z - x_1)^2.$$

This, together with Eq 1.1 above, we have

$$x_2 - x_1 = -\frac{f(x_1)}{f'(x_1)} = z - x_1 + \frac{f^{(2)}(\xi)}{2f'(x_1)}(z - x_1)^2.$$

Therefore, we have

$$(1.2) \quad x_2 - z = \frac{f^{(2)}(\xi)}{2f'(x_1)}(z - x_1)^2.$$

Note that the functions  $f'(x)$  and  $f^{(2)}(x)$  are continuous on  $[z - r, z + r]$  and  $f'(x) \neq 0$ , hence, there is  $M > 0$  such that  $|\frac{f^{(2)}(u)}{2f'(v)}| \leq M$  for all  $u, v \in [z - r, z + r]$ . Then the Eq 1.2 implies that

$$(1.3) \quad |x_2 - z| = \left| \frac{f^{(2)}(\xi)}{2f'(x_1)} (z - x_1)^2 \right| \leq M(z - x_1)^2.$$

Choose  $\delta > 0$  such that  $M\delta < 1$  and  $J := [z - \delta, z + \delta] \subseteq (z - r, z + r)$ . Note that Now we take any  $x_1 \in J$ . Eq 1.3 implies that  $|x_2 - z| \leq M \cdot |z - x_1|^2 \leq (M\delta) \cdot |x_1 - z| < \delta$ . By using Eq 1.1 inductively, we have a sequence  $(x_n)$  in  $J$  such that

$$|x_{n+1} - z| \leq M \cdot |z - x_n|^2 \leq (M\delta) \cdot |x_n - z|$$

for all  $n = 1, 2, \dots$ . Therefore, we have

$$|x_{n+1} - z| \leq (M\delta)^n \cdot |x_1 - z|$$

for all  $n = 1, 2, \dots$ , thus,  $\lim x_n = z$ . The proof is complete.  $\square$



## 2. RIEMANN INTEGRABLE FUNCTIONS

We will use the following notation throughout this chapter.

- (i): All functions  $f, g, h, \dots$  are bounded real valued functions defined on  $[a, b]$  and  $m \leq f \leq M$  on  $[a, b]$ .
- (ii): Let  $P : a = x_0 < x_1 < \dots < x_n = b$  denote a partition on  $[a, b]$ ; Put  $\Delta x_i = x_i - x_{i-1}$  and  $\|P\| = \max \Delta x_i$ .
- (iii):  $M_i(f, P) := \sup\{f(x) : x \in [x_{i-1}, x_i]\}$ ;  $m_i(f, P) := \inf\{f(x) : x \in [x_{i-1}, x_i]\}$ .  
Set  $\omega_i(f, P) = M_i(f, P) - m_i(f, P)$ .
- (iv): (the *upper sum* of  $f$ ):  $U(f, P) := \sum M_i(f, P)\Delta x_i$   
(the *lower sum* of  $f$ ):  $L(f, P) := \sum m_i(f, P)\Delta x_i$ .

**Remark 2.1.** *It is clear that for any partition on  $[a, b]$ , we always have*

- (i)  $m(b-a) \leq L(f, P) \leq U(f, P) \leq M(b-a)$ .
- (ii)  $L(-f, P) = -U(f, P)$  and  $U(-f, P) = -L(f, P)$ .

The following lemma is the critical step in this section.

**Lemma 2.2.** *Let  $P$  and  $Q$  be the partitions on  $[a, b]$ . We have the following assertions.*

- (i) *If  $P \subseteq Q$ , then  $L(f, P) \leq L(f, Q) \leq U(f, Q) \leq U(f, P)$ .*
- (ii) *We always have  $L(f, P) \leq U(f, Q)$ .*

*Proof.* For Part (i), we first claim that  $L(f, P) \leq L(f, Q)$  if  $P \subseteq Q$ . By using the induction on  $l := \#Q - \#P$ , it suffices to show that  $L(f, P) \leq L(f, Q)$  as  $l = 1$ . Let  $P : a = x_0 < x_1 < \dots < x_n = b$  and  $Q = P \cup \{c\}$ . Then  $c \in (x_{s-1}, x_s)$  for some  $s$ . Notice that we have

$$m_s(f, P) \leq \min\{m_s(f, Q), m_{s+1}(f, Q)\}.$$

So, we have

$$m_s(f, P)(x_s - x_{s-1}) \leq m_s(f, Q)(c - x_{s-1}) + m_{s+1}(f, Q)(x_s - c).$$

This gives the following inequality as desired.

$$(2.1) \quad L(f, Q) - L(f, P) = m_s(f, Q)(c - x_{s-1}) + m_{s+1}(f, Q)(x_s - c) - m_s(f, P)(x_s - x_{s-1}) \geq 0.$$

Now by considering  $-f$  in the Inequality 2.1 above, we see that  $U(f, Q) \leq U(f, P)$ .

For Part (ii), let  $P$  and  $Q$  be any pair of partitions on  $[a, b]$ . Notice that  $P \cup Q$  is also a partition on  $[a, b]$  with  $P \subseteq P \cup Q$  and  $Q \subseteq P \cup Q$ . So, Part (i) implies that

$$L(f, P) \leq L(f, P \cup Q) \leq U(f, P \cup Q) \leq U(f, Q).$$

The proof is complete. □

The following notion plays an important role in this chapter.

**Definition 2.3.** *Let  $f$  be a bounded function on  $[a, b]$ . The upper integral (resp. lower integral) of  $f$  over  $[a, b]$ , write  $\overline{\int_a^b} f$  (resp.  $\underline{\int_a^b} f$ ), is defined by*

$$\overline{\int_a^b} f = \inf\{U(f, P) : P \text{ is a partation on } [a, b]\}.$$

(resp.

$$\int_a^b f = \sup\{L(f, P) : P \text{ is a partation on } [a, b]\}.)$$

Notice that the upper integral and lower integral of  $f$  must exist by Remark 2.1.

**Remark 2.4. Appendix:** We call a partially set  $(I, \leq)$  a *directed set* if for each pair of elements  $i_1$  and  $i_2$  in  $I$ , there is  $i_3 \in I$  such that  $i_1 \leq i_3$  and  $i_2 \leq i_3$ .

A *net* in  $\mathbb{R}$  is a real-valued function  $f$  defined on a directed set  $I$ , write  $f = (x_i)_{i \in I}$ , where  $x_i := f(i)$  for  $i \in I$ .

We say that a net  $(x_i)$  converges to a point  $L \in \mathbb{R}$  (call a limit of  $(x_i)$ ) if for any  $\varepsilon > 0$ , there is  $i_0 \in I$  such that  $|x_i - L| < \varepsilon$  for all  $i \geq i_0$ .

Using the similar argument as in the sequence case, a limit of  $(x_i)$  is unique if it exists and we write  $\lim_i x_i$  for its limits.

**Example 2.5. Appendix:** Using the notation given as before, let

$$I := \{P : P \text{ is a partation on } [a, b]\}.$$

We say that  $P_1 \leq P_2$  for  $P_1, P_2 \in I$  if  $P_1 \subseteq P_2$ . Clearly,  $I$  is a directed set with this order. If we put  $u_P := U(f, P)$ , then we have

$$\lim_P u_P = \int_a^b f.$$

In fact, let  $\varepsilon > 0$ . Then by the definition of an upper integral, there is  $P_0 \in I$  such that

$$\int_a^b f \leq U(f, P_0) \leq \int_a^b f + \varepsilon.$$

Lemma 2.2 tells us that whenever  $P \in I$  with  $P \geq P_0$ , we have  $U(f, P) \leq U(f, P_0)$ . Thus we have  $|u_P - \int_a^b f| < \varepsilon$  whenever  $P \geq P_0$  as desired.

**Proposition 2.6.** *Let  $f$  and  $g$  both are bounded functions on  $[a, b]$ . With the notation as above, we always have*

(i)

$$\int_a^b f \leq \int_a^b f.$$

(ii)  $\int_a^b (-f) = -\int_a^b f.$

(iii)

$$\int_a^b f + \int_a^b g \leq \int_a^b (f + g) \leq \int_a^b (f + g) \leq \int_a^b f + \int_a^b g.$$

*Proof.* Part (i) follows from Lemma 2.2 at once.

Part (ii) is clearly obtained by  $L(-f, P) = -U(f, P)$ .

For proving the inequality  $\int_a^b f + \int_a^b g \leq \int_a^b (f + g) \leq$  first. It is clear that we have  $L(f, P) + L(g, P) \leq L(f + g, P)$  for all partitions  $P$  on  $[a, b]$ . Now let  $P_1$  and  $P_2$  be any partition on  $[a, b]$ . Then by Lemma 2.2, we have

$$L(f, P_1) + L(g, P_2) \leq L(f, P_1 \cup P_2) + L(g, P_1 \cup P_2) \leq L(f + g, P_1 \cup P_2) \leq \int_a^b (f + g).$$

So, we have

$$(2.2) \quad \int_a^b f + \int_a^b g \leq \int_a^b (f + g).$$

As before, we consider  $-f$  and  $-g$  in the Inequality 2.2, we get  $\overline{\int_a^b (f + g)} \leq \overline{\int_a^b f} + \overline{\int_a^b g}$  as desired.  $\square$

The following example shows the strict inequality in Proposition 2.6 (iii) may hold in general.

**Example 2.7.** Define a function  $f, g : [0, 1] \rightarrow \mathbb{R}$  by

$$f(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \cap \mathbb{Q}; \\ -1 & \text{otherwise.} \end{cases}$$

and

$$g(x) = \begin{cases} -1 & \text{if } x \in [0, 1] \cap \mathbb{Q}; \\ 1 & \text{otherwise.} \end{cases}$$

Then it is easy to see that  $f + g \equiv 0$  and

$$\int_0^1 f = \int_0^1 g = 1 \quad \text{and} \quad \int_0^1 f = \int_0^1 g = -1.$$

So, we have

$$-2 = \int_a^b f + \int_a^b g < \int_a^b (f + g) = 0 = \overline{\int_a^b (f + g)} < \overline{\int_a^b f} + \overline{\int_a^b g} = 2.$$

We can now reaching the main definition in this chapter.

**Definition 2.8.** Let  $f$  be a bounded function on  $[a, b]$ . We say that  $f$  is Riemann integrable over  $[a, b]$  if  $\overline{\int_a^b f} = \underline{\int_a^b f}$ . In this case, we write  $\int_a^b f$  for this common value and it is called the Riemann integral of  $f$  over  $[a, b]$ .

Also, write  $R[a, b]$  for the class of Riemann integrable functions on  $[a, b]$ .

**Proposition 2.9.** With the notation as above,  $R[a, b]$  is a vector space over  $\mathbb{R}$  and the integral

$$\int_a^b : f \in R[a, b] \mapsto \int_a^b f \in \mathbb{R}$$

defines a linear functional, that is,  $\alpha f + \beta g \in R[a, b]$  and  $\int_a^b (\alpha f + \beta g) = \alpha \int_a^b f + \beta \int_a^b g$  for all  $f, g \in R[a, b]$  and  $\alpha, \beta \in \mathbb{R}$ .

*Proof.* Let  $f, g \in R[a, b]$  and  $\alpha, \beta \in \mathbb{R}$ . Notice that if  $\alpha \geq 0$ , it is clear that  $\overline{\int_a^b \alpha f} = \alpha \overline{\int_a^b f} = \alpha \int_a^b f = \alpha \underline{\int_a^b f} = \underline{\int_a^b \alpha f}$ . Also, if  $\alpha < 0$ , we have  $\overline{\int_a^b \alpha f} = \alpha \underline{\int_a^b f} = \alpha \int_a^b f = \alpha \overline{\int_a^b f} = \underline{\int_a^b \alpha f}$ . Therefore, we have  $\int_a^b \alpha f = \alpha \int_a^b f$  for all  $\alpha \in \mathbb{R}$ . For showing  $f + g \in R[a, b]$  and  $\int_a^b (f + g) = \int_a^b f + \int_a^b g$ , these will follow from Proposition 2.6 (iii) at once. The proof is finished.  $\square$

The following result is the important characterization of a Riemann integrable function. Before showing this, we will use the following notation in the rest of this chapter.

For a partition  $P : a = x_0 < x_1 < \cdots < x_n = b$  and  $1 \leq i \leq n$ , put

$$\omega_i(f, P) := \sup\{|f(x) - f(x')| : x, x' \in [x_{i-1}, x_i]\}.$$

It is easy to see that  $U(f, P) - L(f, P) = \sum_{i=1}^n \omega_i(f, P)\Delta x_i$ .

**Theorem 2.10.** *Let  $f$  be a bounded function on  $[a, b]$ . Then  $f \in R[a, b]$  if and only if for all  $\varepsilon > 0$ , there is a partition  $P : a = x_0 < \cdots < x_n = b$  on  $[a, b]$  such that*

$$(2.3) \quad 0 \leq U(f, P) - L(f, P) = \sum_{i=1}^n \omega_i(f, P)\Delta x_i < \varepsilon.$$

*Proof.* Suppose that  $f \in R[a, b]$ . Let  $\varepsilon > 0$ . Then by the definition of the upper integral and lower integral of  $f$ , we can find the partitions  $P$  and  $Q$  such that  $U(f, P) < \overline{\int_a^b} f + \varepsilon$  and  $\underline{\int_a^b} f - \varepsilon < L(f, Q)$ . By considering the partition  $P \cup Q$ , we see that

$$\underline{\int_a^b} f - \varepsilon < L(f, Q) \leq L(f, P \cup Q) \leq U(f, P \cup Q) \leq U(f, P) < \overline{\int_a^b} f + \varepsilon.$$

Since  $\int_a^b f = \overline{\int_a^b} f = \underline{\int_a^b} f$ , we have  $0 \leq U(f, P \cup Q) - L(f, P \cup Q) < 2\varepsilon$ . So, the partition  $P \cup Q$  is as desired.

Conversely, let  $\varepsilon > 0$ , assume that the Inequality 2.3 above holds for some partition  $P$ . Notice that we have

$$L(f, P) \leq \underline{\int_a^b} f \leq \overline{\int_a^b} f \leq U(f, P).$$

So, we have  $0 \leq \overline{\int_a^b} f - \underline{\int_a^b} f < \varepsilon$  for all  $\varepsilon > 0$ . The proof is finished.  $\square$

**Remark 2.11.** *Theorem 2.10 tells us that a bounded function  $f$  is Riemann integrable over  $[a, b]$  if and only if the “size” of the discontinuous set of  $f$  is arbitrary small. See the Appendix 3 below for details.*

**Example 2.12.** *Let  $f : [0, 1] \rightarrow \mathbb{R}$  be the function defined by*

$$f(x) = \begin{cases} \frac{1}{p} & \text{if } x = \frac{q}{p}, \text{ where } p, q \text{ are relatively prime positive integers;} \\ 0 & \text{otherwise.} \end{cases}$$

*Then  $f \in R[0, 1]$ .*

*(Notice that the set of all discontinuous points of  $f$ , say  $D$ , is just the set of all  $(0, 1] \cap \mathbb{Q}$ . Since the set  $(0, 1] \cap \mathbb{Q}$  is countable, we can write  $(0, 1] \cap \mathbb{Q} = \{z_1, z_2, \dots\}$ . So, if we let  $m(D)$  be the “size” of the set  $D$ , then  $m(D) = m(\bigcup_{i=1}^{\infty} \{z_i\}) = \sum_{i=1}^{\infty} m(\{z_i\}) = 0$ , in here, you may think that the size of each set  $\{z_i\}$  is 0. )*

*Proof.* Let  $\varepsilon > 0$ . By Theorem 2.10, it aims to find a partition  $P$  on  $[0, 1]$  such that

$$U(f, P) - L(f, P) < \varepsilon.$$

Notice that for  $x \in [0, 1]$  such that  $f(x) \geq \varepsilon$  if and only if  $x = q/p$  for a pair of relatively prime positive integers  $p, q$  with  $\frac{1}{p} \geq \varepsilon$ . Since  $1 \leq q \leq p$ , there are only finitely many pairs of relatively prime positive integers  $p$  and  $q$  such that  $f(\frac{q}{p}) \geq \varepsilon$ . So, if we let  $S := \{x \in [0, 1] : f(x) \geq \varepsilon\}$ , then  $S$  is a finite subset

of  $[0, 1]$ . Let  $L$  be the number of the elements in  $S$ . Then, for any partition  $P : a = x_0 < \cdots < x_n = 1$ , we have

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i = \left( \sum_{i: [x_{i-1}, x_i] \cap S = \emptyset} + \sum_{i: [x_{i-1}, x_i] \cap S \neq \emptyset} \right) \omega_i(f, P) \Delta x_i.$$

Notice that if  $[x_{i-1}, x_i] \cap S = \emptyset$ , then we have  $\omega_i(f, P) \leq \varepsilon$  and thus,

$$\sum_{i: [x_{i-1}, x_i] \cap S = \emptyset} \omega_i(f, P) \Delta x_i \leq \varepsilon \sum_{i: [x_{i-1}, x_i] \cap S = \emptyset} \Delta x_i \leq \varepsilon(1 - 0).$$

On the other hand, since there are at most  $2L$  sub-intervals  $[x_{i-1}, x_i]$  such that  $[x_{i-1}, x_i] \cap S \neq \emptyset$  and  $\omega_i(f, P) \leq 1$  for all  $i = 1, \dots, n$ , so, we have

$$\sum_{i: [x_{i-1}, x_i] \cap S \neq \emptyset} \omega_i(f, P) \Delta x_i \leq 1 \cdot \sum_{i: [x_{i-1}, x_i] \cap S \neq \emptyset} \Delta x_i \leq 2L \|P\|.$$

We can now conclude that for any partition  $P$ , we have

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i \leq \varepsilon + 2L \|P\|.$$

So, if we take a partition  $P$  with  $\|P\| < \varepsilon/(2L)$ , then we have  $\sum_{i=1}^n \omega_i(f, P) \Delta x_i \leq 2\varepsilon$ .

The proof is finished.  $\square$

**Proposition 2.13.** *Let  $f$  be a function defined on  $[a, b]$ . If  $f$  is either monotone or continuous on  $[a, b]$ , then  $f \in R[a, b]$ .*

*Proof.* We first show the case of  $f$  being monotone. We may assume that  $f$  is monotone increasing. Notice that for any partition  $P : a = x_0 < \cdots < x_n = b$ , we have  $\omega_i(f, P) = f(x_i) - f(x_{i-1})$ . So, if  $\|P\| < \varepsilon$ , we have

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i = \sum_{i=1}^n (f(x_i) - f(x_{i-1})) \Delta x_i < \|P\| \sum_{i=1}^n (f(x_i) - f(x_{i-1})) = \|P\| (f(b) - f(a)) < \varepsilon (f(b) - f(a)).$$

Therefore,  $f \in R[a, b]$  if  $f$  is monotone.

Suppose that  $f$  is continuous on  $[a, b]$ . Then  $f$  is uniform continuous on  $[a, b]$ . Then for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $|f(x) - f(x')| < \varepsilon$  as  $x, x' \in [a, b]$  with  $|x - x'| < \delta$ . So, if we choose a partition  $P$  with  $\|P\| < \delta$ , then  $\omega_i(f, P) < \varepsilon$  for all  $i$ . This implies that

$$\sum_{i=1}^n \omega_i(f, P) \Delta x_i \leq \varepsilon \sum_{i=1}^n \Delta x_i = \varepsilon(b - a).$$

The proof is complete.  $\square$

**Proposition 2.14.** *We have the following assertions.*

(i) *If  $f, g \in R[a, b]$  with  $f \leq g$ , then  $\int_a^b f \leq \int_a^b g$ .*

(ii) *If  $f \in R[a, b]$ , then the absolute valued function  $|f| \in R[a, b]$ . In this case, we have  $|\int_a^b f| \leq \int_a^b |f|$ .*

*Proof.* For Part (i), it is clear that we have the inequality  $U(f, P) \leq U(g, P)$  for any partition  $P$ . So, we have  $\int_a^b f = \overline{\int_a^b f} \leq \overline{\int_a^b g} = \int_a^b g$ .

For Part (ii), the integrability of  $|f|$  follows immediately from Theorem 2.10 and the simple inequality  $||f|(x') - |f|(x'')| \leq |f(x') - f(x'')|$  for all  $x', x'' \in [a, b]$ . Thus, we have  $U(|f|, P) - L(|f|, P) \leq$

$U(f, P) - L(f, P)$  for any partition  $P$  on  $[a, b]$ .

Finally, since we have  $-f \leq |f| \leq f$ , by Part (i), we have  $|\int_a^b f| \leq \int_a^b |f|$  at once.  $\square$

**Proposition 2.15.** *Let  $a < c < b$ . We have  $f \in R[a, b]$  if and only if the restrictions  $f|_{[a, c]} \in R[a, c]$  and  $f|_{[c, b]} \in R[c, b]$ . In this case we have*

$$(2.4) \quad \int_a^b f = \int_a^c f + \int_c^b f.$$

*Proof.* Let  $f_1 := f|_{[a, c]}$  and  $f_2 := f|_{[c, b]}$ .

It is clear that we always have

$$U(f_1, P_1) - L(f_1, P_1) + U(f_2, P_2) - L(f_2, P_2) = U(f, P) - L(f, P)$$

for any partition  $P_1$  on  $[a, c]$  and  $P_2$  on  $[c, b]$  with  $P = P_1 \cup P_2$ .

From this, we can show the sufficient condition at once.

For showing the necessary condition, since  $f \in R[a, b]$ , for any  $\varepsilon > 0$ , there is a partition  $Q$  on  $[a, b]$  such that  $U(f, Q) - L(f, Q) < \varepsilon$  by Theorem 2.10. Notice that there are partitions  $P_1$  and  $P_2$  on  $[a, c]$  and  $[c, b]$  respectively such that  $P := Q \cup \{c\} = P_1 \cup P_2$ . Thus, we have

$$U(f_1, P_1) - L(f_1, P_1) + U(f_2, P_2) - L(f_2, P_2) = U(f, P) - L(f, P) \leq U(f, Q) - L(f, Q) < \varepsilon.$$

So, we have  $f_1 \in R[a, c]$  and  $f_2 \in R[c, b]$ .

It remains to show the Equation 2.4 above. Notice that for any partition  $P_1$  on  $[a, c]$  and  $P_2$  on  $[c, b]$ , we have

$$L(f_1, P_1) + L(f_2, P_2) = L(f, P_1 \cup P_2) \leq \int_a^b f = \int_a^c f + \int_c^b f.$$

So, we have  $\int_a^c f + \int_c^b f \leq \int_a^b f$ . Then the inverse inequality can be obtained at once by considering the function  $-f$ . Then the result is obtained by using Theorem 2.10.  $\square$

**Proposition 2.16.** *Let  $f$  and  $g$  be Riemann integrable functions defined on  $[a, b]$ . Then the pointwise product function  $f \cdot g \in R[a, b]$ .*

*Proof.* We first show that the square function  $f^2$  is Riemann integrable. In fact, if we let  $M = \sup\{|f(x)| : x \in [a, b]\}$ , then we have  $\omega_k(f^2, P) \leq 2M\omega_k(f, P)$  for any partition  $P : a = x_0 < \dots < x_n = b$  because we always have  $|f^2(x) - f^2(x')| \leq 2M|f(x) - f(x')|$  for all  $x, x' \in [a, b]$ . Then by Theorem 2.10, the square function  $f^2 \in R[a, b]$ .

This, together with the identity  $f \cdot g = \frac{1}{2}((f+g)^2 - f^2 - g^2)$ . The result follows.  $\square$

**Remark 2.17.** *In the proof of Proposition 2.16, we have shown that if  $f \in R[a, b]$ , then so is its square function  $f^2$ . However, the converse does not hold. For example, if we consider  $f(x) = 1$  for  $x \in \mathbb{Q} \cap [0, 1]$  and  $f(x) = -1$  for  $x \in \mathbb{Q}^c \cap [0, 1]$ , then  $f \notin R[0, 1]$  but  $f^2 \equiv 1$  on  $[0, 1]$ .*

**Proposition 2.18.** *Assume that  $f : [a, b] \rightarrow [c, d]$  is integrable and  $g : [c, d] \rightarrow \mathbb{R}$  is continuous. Then the composition  $g \circ f \in R[a, b]$ .*

*Proof.* Let  $\varepsilon > 0$ . Note that  $g$  is uniformly continuous on  $[c, d]$  because  $g$  is continuous on  $[c, d]$ . Then there is  $\delta > 0$  such that  $|g(y) - g(y')| < \varepsilon$  whenever  $y, y' \in [c, d]$  with  $|y - y'| < \delta$ . On the other hand, since  $f \in R[a, b]$ , there is a partition  $P$  on  $[a, b]$  such that  $\sum \omega_k(f, P)\Delta x_k < \varepsilon\delta$ . Hence, we have

$$\delta \sum_{k: \omega_k(f, P) \geq \delta} \Delta x_k \leq \delta \sum_{k: \omega_k(f, P) \geq \delta} \omega_k(f, P)\Delta x_k < \varepsilon\delta.$$

This implies that

$$\sum_{k:\omega_k(f,P)\geq\delta} \Delta x_k < \varepsilon.$$

On the other hand, by the choice of  $\delta$ , we see that  $\omega_k(g \circ f, P) < \varepsilon$  whenever  $\omega_k(f, P) < \delta$ . Therefore, we can conclude that

$$\sum_k \omega_k(g \circ f, P) \Delta x_k = \sum_{k:\omega_k(f,P)<\delta} \omega_k(g \circ f, P) \Delta x_k + \sum_{k:\omega_k(f,P)\geq\delta} \omega_k(g \circ f, P) \Delta x_k < \varepsilon(b-a) + 2M\varepsilon$$

where  $M := \sup |f(x)|$ . The proof is complete.  $\square$

**Remark 2.19.** *The composition of integrable functions need not be integrable. For example, if we put  $f$  is given as in Example 2.12 and  $g(x) = x$  for  $x = 1/n, n = 1, 2, \dots$ ; otherwise  $g(x) = 0$ . Then  $f, g \in R[0, 1]$  but  $g \circ f \notin R[0, 1]$ .*

**Proposition 2.20. (Mean Value Theorem for Integrals)**

Let  $f$  and  $g$  be the functions defined on  $[a, b]$ . Assume that  $f$  is continuous and  $g$  is a non-negative Riemann integrable function. Then, there is a point  $\xi \in (a, b)$  such that

$$(2.5) \quad \int_a^b f(x)g(x)dx = f(\xi) \int_a^b g(x)dx.$$

In particular, there is a point  $\xi$  in  $(a, b)$  such that  $f(\xi) = \frac{1}{b-a} \int_a^b f(x)dx$ .

*Proof.* By the continuity of  $f$  on  $[a, b]$ , there exist two points  $x_1$  and  $x_2$  in  $[a, b]$  such that

$$f(x_1) = m := \min f(x); \text{ and } f(x_2) = M := \max f(x).$$

We may assume that  $a \leq x_1 < x_2 \leq b$ . From this, since  $g \geq 0$ , we have

$$mg(x) \leq f(x)g(x) \leq Mg(x)$$

for all  $x \in [a, b]$ . From this and Proposition 2.16 above, we have

$$m \int_a^b g \leq \int_a^b fg \leq M \int_a^b g.$$

So, if  $\int_a^b g = 0$ , then the result follows at once.

We may now suppose that  $\int_a^b g > 0$ . The above inequality shows that

$$m = f(x_1) \leq \frac{\int_a^b fg}{\int_a^b g} \leq f(x_2) = M.$$

Therefore, there is a point  $\xi \in [x_1, x_2] \subseteq [a, b]$  so that the Equation 2.5 holds by using the Intermediate Value Theorem for the function  $f$ . Thus, it remains to show that such element  $\xi$  can be chosen in  $(a, b)$ .

Let  $a \leq x_1 < x_2 \leq b$  be as above.

If  $x_1$  and  $x_2$  can be found so that  $a < x_1 < x_2 < b$ , then the result is proved immediately since  $\xi \in [x_1, x_2] \subset (a, b)$  in this case.

Now suppose that  $x_1$  or  $x_2$  does not exist in  $(a, b)$ , i.e.,  $m = f(a) < f(x)$  for all  $x \in (a, b]$  or  $f(x) < f(b) = M$  for all  $x \in [a, b)$ .

**Claim 1:** If  $f(a) < f(x)$  for all  $x \in (a, b]$ , then  $\int_a^b fg > f(a) \int_a^b g$  and hence,  $\xi \in (a, x_2] \subseteq (a, b]$ .

For showing **Claim1**, put  $h(x) := f(x) - f(a)$  for  $x \in [a, b]$ . Then  $h$  is continuous on  $[a, b]$  and  $h > 0$  on  $(a, b]$ . This implies that  $\int_c^d h > 0$  for any subinterval  $[c, d] \subseteq [a, b]$ . (**Why?**)

On the other hand, since  $\int_a^b g = \int_a^b g > 0$ , there is a partition  $P : a = x_0 < \cdots < x_n = b$  so that  $L(g, P) > 0$ . This implies that  $m_k(g, P) > 0$  for some sub-interval  $[x_{k-1}, x_k]$ . Therefore, we have

$$\int_a^b hg \geq \int_{x_{k-1}}^{x_k} hg \geq m_k(g, P) \int_{x_{k-1}}^{x_k} h > 0.$$

Hence, we have  $\int_a^b fg > f(a) \int_a^b g$ . **Claim 1** follows.

Similarly, one can show that if  $f(x) < f(b) = M$  for all  $x \in [a, b)$ , then we have  $\int_a^b fg < f(b) \int_a^b g$ . This, together with **Claim 1** give us that such  $\xi$  can be found in  $(a, b)$ . The proof is finished.  $\square$

**Example 2.21.** We have  $\lim_n \int_0^{\pi/2} \sin^n x dx = 0$ . To see this, for any  $0 < \varepsilon < \pi/2$  and for each  $n = 1, 2, \dots$ , the Mean value theorem gives a point  $\xi_n \in (0, \frac{\pi}{2} - \varepsilon)$  such that

$$\begin{aligned} 0 < \int_0^{\pi/2} \sin^n x dx &= \left( \int_0^{\frac{\pi}{2}-\varepsilon} + \int_{\frac{\pi}{2}-\varepsilon}^{\pi/2} \right) \sin^n x dx \\ &\leq \sin^{n-1} \xi_n \int_0^{\frac{\pi}{2}-\varepsilon} \sin x dx + \int_{\frac{\pi}{2}-\varepsilon}^{\pi/2} \sin^n x dx \\ &< \sin^{n-1} \left( \frac{\pi}{2} - \varepsilon \right) + \varepsilon. \end{aligned}$$

Taking  $n \rightarrow \infty$ , we have  $\overline{\lim}_n \int_0^{\pi/2} \sin^n x dx = 0$ . The proof is finished.

Now if  $f \in R[a, b]$ , then by Proposition 2.15, we can define a function  $F : [a, b] \rightarrow \mathbb{R}$  by

$$(2.6) \quad F(c) = \begin{cases} 0 & \text{if } c = a \\ \int_a^c f & \text{if } a < c \leq b. \end{cases}$$

**Theorem 2.22. Fundamental Theorem of Calculus:** *With the notation as above, assume that  $f \in R[a, b]$ , we have the following assertion.*

- (i) *If there is a continuous function  $F$  on  $[a, b]$  which is differentiable on  $(a, b)$  with  $F' = f$ , then  $\int_a^b f = F(b) - F(a)$ . In this case,  $F$  is called an indefinite integral of  $f$ . (**note:** if  $F_1$  and  $F_2$  both are the indefinite integrals of  $f$ , then by the Mean Value Theorem, we have  $F_2 = F_1 + \text{constant}$ ).*
- (ii) *The function  $F$  defined as in Eq. 2.6 above is continuous on  $[a, b]$ . Furthermore, if  $f$  is continuous on  $[a, b]$ , then  $F'$  exists on  $(a, b)$  and  $F' = f$  on  $(a, b)$ .*

*Proof.* For Part (i), notice that for any partition  $P : a = x_0 < \cdots < x_n = b$ , then by the Mean Value Theorem, for each  $[x_{i-1}, x_i]$ , there is  $\xi_i \in (x_{i-1}, x_i)$  such that  $F(x_i) - F(x_{i-1}) = F'(\xi_i) \Delta x_i = f(\xi_i) \Delta x_i$ . So, we have

$$L(f, P) \leq \sum f(\xi_i) \Delta x_i = \sum F(x_i) - F(x_{i-1}) = F(b) - F(a) \leq U(f, P)$$

for all partitions  $P$  on  $[a, b]$ . This gives

$$\int_a^b f = \int_a^b f \leq F(b) - F(a) \leq \overline{\int_a^b f} = \int_a^b f$$

as desired.

For showing the continuity of  $F$  in Part (ii), let  $a < c < x < b$ . If  $|f| \leq M$  on  $[a, b]$ , then we have  $|F(x) - F(c)| = \left| \int_c^x f \right| \leq M(x - c)$ . So,  $\lim_{x \rightarrow c^+} F(x) = F(c)$ . Similarly, we also have  $\lim_{x \rightarrow c^-} F(x) =$



$F(c)$ . Thus  $F$  is continuous on  $[a, b]$ .

Now assume that  $f$  is continuous on  $[a, b]$ . Notice that for any  $t > 0$  with  $a < c < c + t < b$ , we have

$$\inf_{x \in [c, c+t]} f(x) \leq \frac{1}{t}(F(c+t) - F(c)) = \frac{1}{t} \int_c^{c+t} f \leq \sup_{x \in [c, c+t]} f(x).$$

Since  $f$  is continuous at  $c$ , we see that  $\lim_{t \rightarrow 0^+} \frac{1}{t}(F(c+t) - F(c)) = f(c)$ . Similarly, we have  $\lim_{t \rightarrow 0^-} \frac{1}{t}(F(c+t) - F(c)) = f(c)$ . So, we have  $F'(c) = f(c)$  as desired. The proof is finished.  $\square$

**Definition 2.23.** For each function  $f$  on  $[a, b]$  and a partition  $P : a = x_0 < \dots < x_n = b$ , we call  $R(f, P, \{\xi_i\}) := \sum_{i=1}^n f(\xi_i) \Delta x_i$ , where  $\xi_i \in [x_{i-1}, x_i]$ , the Riemann sum of  $f$  over  $[a, b]$ .

We say that the Riemann sum  $R(f, P, \{\xi_i\})$  converges to a number  $A$  as  $\|P\| \rightarrow 0$ , write  $A = \lim_{\|P\| \rightarrow 0} R(f, P, \{\xi_i\})$ , if for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that

$$|A - R(f, P, \{\xi_i\})| < \varepsilon$$

whenever  $\|P\| < \delta$  and for any  $\xi_i \in [x_{i-1}, x_i]$ .

**Proposition 2.24.** Let  $f$  be a function defined on  $[a, b]$ . If the limit  $\lim_{\|P\| \rightarrow 0} R(f, P, \{\xi_i\}) = A$  exists, then  $f$  is automatically bounded.

*Proof.* Suppose that  $f$  is unbounded. Then by the assumption, there exists a partition  $P : a = x_0 < \dots < x_n = b$  such that  $|\sum_{k=1}^n f(\xi_k) \Delta x_k| < 1 + |A|$  for any  $\xi_k \in [x_{k-1}, x_k]$ . Since  $f$  is unbounded, we may assume that  $f$  is unbounded on  $[a, x_1]$ . In particular, we choose  $\xi_k = x_k$  for  $k = 2, \dots, n$ . Also, we can choose  $\xi_1 \in [a, x_1]$  such that

$$|f(\xi_1)| \Delta x_1 < 1 + |A| + \left| \sum_{k=2}^n f(x_k) \Delta x_k \right|.$$

It leads to a contradiction because we have  $1 + |A| > |f(\xi_1)| \Delta x_1 - \left| \sum_{k=2}^n f(x_k) \Delta x_k \right|$ . The proof is finished.  $\square$

**Lemma 2.25.**  $f \in R[a, b]$  if and only if for any  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $U(f, P) - L(f, P) < \varepsilon$  whenever  $\|P\| < \delta$ .

*Proof.* The converse follows from Theorem 2.10.

Assume that  $f$  is integrable over  $[a, b]$ . Let  $\varepsilon > 0$ . Then there is a partition  $Q : a = y_0 < \dots < y_l = b$  on  $[a, b]$  such that  $U(f, Q) - L(f, Q) < \varepsilon$ . Now take  $0 < \delta < \varepsilon/l$ . Suppose that  $P : a = x_0 < \dots < x_n = b$  with  $\|P\| < \delta$ . Then we have

$$U(f, P) - L(f, P) = I + II$$

where

$$I = \sum_{i: Q \cap [x_{i-1}, x_i] = \emptyset} \omega_i(f, P) \Delta x_i;$$

and

$$II = \sum_{i: Q \cap [x_{i-1}, x_i] \neq \emptyset} \omega_i(f, P) \Delta x_i$$

Notice that we have

$$I \leq U(f, Q) - L(f, Q) < \varepsilon$$

and

$$II \leq (M - m) \sum_{i: Q \cap [x_{i-1}, x_i] \neq \emptyset} \Delta x_i \leq (M - m) \cdot 2l \cdot \frac{\varepsilon}{l} = 2(M - m)\varepsilon.$$

The proof is finished. □

**Theorem 2.26.**  $f \in R[a, b]$  if and only if the Riemann sum  $R(f, P, \{\xi_i\})$  is convergent. In this case,  $R(f, P, \{\xi_i\})$  converges to  $\int_a^b f(x)dx$  as  $\|P\| \rightarrow 0$ .

*Proof.* For the proof ( $\Rightarrow$ ): we first note that we always have

$$L(f, P) \leq R(f, P, \{\xi_i\}) \leq U(f, P)$$

and

$$L(f, P) \leq \int_a^b f(x)dx \leq U(f, P)$$

for any partition  $P$  and  $\xi_i \in [x_{i-1}, x_i]$ .

Now let  $\varepsilon > 0$ . Lemma 2.25 gives  $\delta > 0$  such that  $U(f, P) - L(f, P) < \varepsilon$  as  $\|P\| < \delta$ . Then we have

$$\left| \int_a^b f(x)dx - R(f, P, \{\xi_i\}) \right| < \varepsilon$$

as  $\|P\| < \delta$  and  $\xi_i \in [x_{i-1}, x_i]$ . The necessary part is proved and  $R(f, P, \{\xi_i\})$  converges to  $\int_a^b f(x)dx$ . For ( $\Leftarrow$ ): assume that there is a number  $A$  such that for any  $\varepsilon > 0$ , there is  $\delta > 0$ , we have

$$A - \varepsilon < R(f, P, \{\xi_i\}) < A + \varepsilon$$

for any partition  $P$  with  $\|P\| < \delta$  and  $\xi_i \in [x_{i-1}, x_i]$ .

Note that  $f$  is automatically bounded in this case by Proposition 2.24.

Now fix a partition  $P$  with  $\|P\| < \delta$ . Then for each  $[x_{i-1}, x_i]$ , choose  $\xi_i \in [x_{i-1}, x_i]$  such that  $M_i(f, P) - \varepsilon \leq f(\xi_i)$ . This implies that we have

$$U(f, P) - \varepsilon(b - a) \leq R(f, P, \{\xi_i\}) < A + \varepsilon.$$

Thus, we have shown that for any  $\varepsilon > 0$ , there is a partition  $\mathcal{P}$  such that

$$(2.7) \quad \int_a^b f(x)dx \leq U(f, P) \leq A + \varepsilon(1 + b - a).$$

By considering  $-f$ , note that the Riemann sum of  $-f$  will converge to  $-A$ . The inequality 2.7 will imply that for any  $\varepsilon > 0$ , there is a partition  $P$  such that

$$A - \varepsilon(1 + b - a) \leq \int_a^b f(x)dx \leq \int_a^b f(x)dx \leq A + \varepsilon(1 + b - a).$$

The proof is complete. □

**Theorem 2.27.** Let  $f \in R[c, d]$  and let  $\phi : [a, b] \rightarrow [c, d]$  be a strictly increasing function with  $\phi(a) = c$  and  $\phi(b) = d$ . Assume that  $\phi$  is a  $C^1$  function and  $\phi'$  can be extended to a strictly continuous function on  $[a, b]$ . Then  $f \circ \phi \in R[a, b]$ , moreover, we have

$$\int_c^d f(x)dx = \int_a^b f(\phi(t))\phi'(t)dt.$$

*Proof.* Let  $A = \int_c^d f(x)dx$ . By using Theorem 2.26, we need to show that for all  $\varepsilon > 0$ , there is  $\delta > 0$  such that

$$\left| A - \sum f(\phi(\xi_k))\phi'(\xi_k)\Delta t_k \right| < \varepsilon$$

for all  $\xi_k \in [t_{k-1}, t_k]$  whenever  $Q : a = t_0 < \dots < t_m = b$  with  $\|Q\| < \delta$ .

Now let  $\varepsilon > 0$ . Then by Lemma 2.25 and Theorem 2.26, there is  $\delta_1 > 0$  such that

$$(2.8) \quad |A - \sum f(\eta_k) \Delta x_k| < \varepsilon$$

and

$$(2.9) \quad \sum \omega_k(f, P) \Delta x_k < \varepsilon$$

for all  $\eta_k \in [x_{k-1}, x_k]$  whenever  $P : c = x_0 < \dots < x_m = d$  with  $\|P\| < \delta_1$ .

Now put  $x = \phi(t)$  for  $t \in [a, b]$ .

Note that there is  $\delta > 0$  such that  $|\phi(t) - \phi(t')| < \delta_1$  and  $|\phi'(t) - \phi'(t')| < \varepsilon$  for all  $t, t'$  in  $[a, b]$  with  $|t - t'| < \delta$ .

Now let  $Q : a = t_0 < \dots < t_m = b$  with  $\|Q\| < \delta$ . If we put  $x_k = \phi(t_k)$ , then  $P : c = x_0 < \dots < x_m = d$  is a partition on  $[c, d]$  with  $\|P\| < \delta_1$  because  $\phi$  is strictly increasing.

Note that the Mean Value Theorem implies that for each  $[t_{k-1}, t_k]$ , there is  $\xi_k^* \in (t_{k-1}, t_k)$  such that

$$\Delta x_k = \phi(t_k) - \phi(t_{k-1}) = \phi'(\xi_k^*) \Delta t_k.$$

This yields that

$$(2.10) \quad |\Delta x_k - \phi'(\xi_k) \Delta t_k| < \varepsilon \Delta t_k$$

for any  $\xi_k \in [t_{k-1}, t_k]$  for all  $k = 1, \dots, m$  because of the choice of  $\delta$ .

Now for any  $\xi_k \in [t_{k-1}, t_k]$ , we have

$$(2.11) \quad \begin{aligned} |A - \sum f(\phi(\xi_k)) \phi'(\xi_k) \Delta t_k| &\leq |A - \sum f(\phi(\xi_k^*)) \phi'(\xi_k^*) \Delta t_k| \\ &+ | \sum f(\phi(\xi_k^*)) \phi'(\xi_k^*) \Delta t_k - \sum f(\phi(\xi_k^*)) \phi'(\xi_k) \Delta t_k | \\ &+ | \sum f(\phi(\xi_k^*)) \phi'(\xi_k) \Delta t_k - \sum f(\phi(\xi_k)) \phi'(\xi_k) \Delta t_k | \end{aligned}$$

Notice that inequality 2.8 implies that

$$|A - \sum f(\phi(\xi_k^*)) \phi'(\xi_k^*) \Delta t_k| = |A - \sum f(\phi(\xi_k^*)) \Delta x_k| < \varepsilon.$$

Moreover, since we have  $|\phi'(\xi_k^*) - \phi'(\xi_k)| < \varepsilon$  for all  $k = 1, \dots, m$ , we have

$$| \sum f(\phi(\xi_k^*)) \phi'(\xi_k^*) \Delta t_k - \sum f(\phi(\xi_k^*)) \phi'(\xi_k) \Delta t_k | \leq M(b-a)\varepsilon$$

where  $|f(x)| \leq M$  for all  $x \in [c, d]$ .

On the other hand, by using inequality 2.10 we have

$$|\phi'(\xi_k) \Delta t_k| \leq \Delta x_k + \varepsilon \Delta t_k$$

for all  $k$ . This, together with inequality 2.9 imply that

$$\begin{aligned} &| \sum f(\phi(\xi_k^*)) \phi'(\xi_k) \Delta t_k - \sum f(\phi(\xi_k)) \phi'(\xi_k) \Delta t_k | \\ &\leq \sum \omega_k(f, P) |\phi'(\xi_k) \Delta t_k| \quad (\because \phi(\xi_k^*), \phi(\xi_k) \in [x_{k-1}, x_k]) \\ &\leq \sum \omega_k(f, P) (\Delta x_k + \varepsilon \Delta t_k) \\ &\leq \varepsilon + 2M(b-a)\varepsilon. \end{aligned}$$

Finally by inequality 2.11, we have

$$|A - \sum f(\phi(\xi_k)) \phi'(\xi_k) \Delta t_k| \leq \varepsilon + M(b-a)\varepsilon + \varepsilon + 2M(b-a)\varepsilon.$$

Finally, we have to show that  $f \circ \phi \in R[a, b]$ . To see this, we have shown that the function  $f \circ \phi(t) \phi'(t) \in R[0, 1]$  by above. Since  $\phi' > 0$  is continuous on  $[a, b]$ ,  $\frac{1}{\phi'}$  is continuous on  $[a, b]$  and thus  $\frac{1}{\phi'} \in R[a, b]$ .

This implies that the function  $f \circ \phi = \frac{1}{\phi'} (f \circ \phi \cdot \phi') \in R[0, 1]$  as desired. The proof is complete.  $\square$

**Lemma 2.28.** *Let  $g$  be a convex function defined on  $[a, b]$ . Then for  $a < c < x < d < b$ , we have*

$$\frac{g(x) - g(c)}{x - c} \leq \frac{g(d) - g(c)}{d - x}.$$

*Proof.* Let  $\ell(x)$  be the straight line between the points  $(c, g(c))$  and  $(d, g(d))$ . Then we have  $g(x) \leq \ell(x)$  for all  $x \in [c, d]$  by the convexity. This implies the following that we desired.

$$\frac{g(x) - g(c)}{x - c} \leq \frac{\ell(x) - \ell(c)}{x - c} = \frac{\ell(d) - \ell(x)}{d - x} \leq \frac{g(d) - g(c)}{d - x}.$$

□

**Proposition 2.29. (Jensen's inequality):** *Let  $g : [a', b'] \rightarrow \mathbb{R}$  be a convex function and  $f \in R([0, 1])$  such that  $f([0, 1]) \subseteq [a, b] \subseteq [a', b']$  and  $g \circ f \in R([0, 1])$ . Then we have*

$$g\left(\int_0^1 f(x)dx\right) \leq \int_0^1 (g \circ f)(x)dx.$$

*Proof.* Notice that if we let  $c := \int_0^1 f$ , then  $c \in [a, b]$  and hence,  $g(c)$  is defined. Let  $s := \sup\{\frac{g(c)-g(x)}{c-x} : a' < x < c\}$ . Then by Lemma 2.28, we have  $g(c) + s(f(x) - c) \leq (g \circ f)(x)$  for all  $x \in [0, 1]$ . This gives

$$g(c) = g(c) + s \int_0^1 (f(x) - c)dx \leq \int_0^1 (g \circ f)(x)dx.$$

The proof is complete. □

**Example 2.30.** *Let  $a_1, \dots, a_n$  be any real numbers. Let  $p > 1$ . Then we have*

$$\left(\frac{|a_1| + \dots + |a_n|}{n}\right)^p \leq \frac{1}{n} \sum_{k=1}^n |a_k|^p.$$

*To see this, , the results obtained by applying the Jensen's inequality for the convex function  $g(x) = x^p$  for  $x \geq 0$  and  $f(t) := |a_k|$  for  $t \in [(k-1)/n, k/n]$  for  $k = 1, \dots, n$ .*

**Definition 2.31.** *Let  $-\infty < a < b < \infty$ .*

(i) *Let  $f$  be a function defined on  $[a, \infty)$ . Assume that the restriction  $f|_{[a, T]}$  is integrable over*

*$[a, T]$  for all  $T > a$ . Put  $\int_a^\infty f := \lim_{T \rightarrow \infty} \int_a^T f$  if this limit exists.*

*Similarly, we can define  $\int_{-\infty}^b f$  if  $f$  is defined on  $(-\infty, b]$ .*

(ii) *If  $f$  is defined on  $(a, b]$  and  $f|_{[c, b]} \in R[c, b]$  for all  $a < c < b$ . Put  $\int_a^b f := \lim_{c \rightarrow a^+} \int_c^b f$  if it exists.*

*Similarly, we can define  $\int_a^b f$  if  $f$  is defined on  $[a, b)$ .*

(iii) *As  $f$  is defined on  $\mathbb{R}$ , if  $\int_0^\infty f$  and  $\int_{-\infty}^0 f$  both exist, then we put  $\int_{-\infty}^\infty f = \int_{-\infty}^0 f + \int_0^\infty f$ .*

*In the cases above, we call the resulting limits the improper Riemann integrals of  $f$  and say that the integrals are convergent.*

Clearly, the Cauchy criterion will imply the following immediately.

**Proposition 2.32.** *Let  $f : [a, \infty) \rightarrow \mathbb{R}$  be a function given as in Definition 2.31.*

(i) *The improper integral  $\int_a^\infty f$  exists if and only if for any  $\varepsilon > 0$ , there is  $M > 0$  such that  $|\int_A^B f| < \varepsilon$  whenever  $M < A < B$ .*

(ii) Let  $g$  be a non-negative function defined on  $[a, \infty)$  such that  $|f| \leq g$  on  $[a, \infty)$ . If  $\int_a^\infty g$  is convergent, then so is  $\int_a^\infty f$ .

(iii) Suppose that  $0 \leq g \leq f$  on  $[a, \infty)$ . If  $\int_a^\infty g$  is divergent, then so is  $\int_a^\infty f$ .

Similar assertion holds when  $f$  is defined on  $(a, b]$ .

**Remark 2.33.** By using the Cauchy Theorem, it is clear that if  $\int_a^\infty |f|$  is convergent, then so is the integral  $\int_a^\infty f$ . However, the converse does not hold. It is quite different from the case when  $f$  is defined on  $[a, b]$ .

For example, if  $f(x) = \frac{(-1)^{n-1}}{n}$  as  $n \in [n-1, n)$ ,  $n = 1, 2, \dots$ , then  $\int_a^\infty f$  is convergent (it will be shown in the last chapter) but  $\int_a^\infty |f|$  is divergent.

**Example 2.34.** Define (formally) an improper integral  $\Gamma(s)$  (called the  $\Gamma$ -function) as follows:

$$\Gamma(s) := \int_0^\infty x^{s-1} e^{-x} dx$$

for  $s \in \mathbb{R}$ . Then  $\Gamma(s)$  is convergent if and only if  $s > 0$ .

*Proof.* Put  $I(s) := \int_0^1 x^{s-1} e^{-x} dx$  and  $II(s) := \int_1^\infty x^{s-1} e^{-x} dx$ . We first claim that the integral  $II(s)$  is convergent for all  $s \in \mathbb{R}$ .

In fact, if we fix  $s \in \mathbb{R}$ , then we have

$$\lim_{x \rightarrow \infty} \frac{x^{s-1}}{e^{x/2}} = 0.$$

So there is  $M > 1$  such that  $\frac{x^{s-1}}{e^{x/2}} \leq 1$  for all  $x \geq M$ . Thus we have

$$0 \leq \int_M^\infty x^{s-1} e^{-x} dx \leq \int_M^\infty e^{-x/2} dx < \infty.$$

Therefore we need to show that the integral  $I(s)$  is convergent if and only if  $s > 0$ .

Note that for  $0 < \eta < 1$ , we have

$$0 \leq \int_\eta^1 x^{s-1} e^{-x} dx \leq \int_\eta^1 x^{s-1} dx = \begin{cases} \frac{1}{s}(1 - \eta^s) & \text{if } s - 1 \neq -1; \\ -\ln \eta & \text{otherwise.} \end{cases}$$

Thus the integral  $I(s) = \lim_{\eta \rightarrow 0^+} \int_\eta^1 x^{s-1} e^{-x} dx$  is convergent if  $s > 0$ .

Conversely, we also have

$$\int_\eta^1 x^{s-1} e^{-x} dx \geq e^{-1} \int_\eta^1 x^{s-1} dx = \begin{cases} \frac{e^{-1}}{s}(1 - \eta^s) & \text{if } s - 1 \neq -1; \\ -e^{-1} \ln \eta & \text{otherwise.} \end{cases}$$

So if  $s \leq 0$ , then  $\int_\eta^1 x^{s-1} e^{-x} dx$  is divergent as  $\eta \rightarrow 0^+$ . The result follows.  $\square$

### 3. APPENDIX: LEBESGUE INTEGRABILITY THEOREM

Throughout this section, let  $f$  be a  $\mathbb{R}$ -valued function defined on  $[a, b]$  and let  $M := \sup |f(x)|$ .

**Definition 3.1.** A subset  $A$  of  $\mathbb{R}$  is said to have measure zero (or null set) if for every  $\varepsilon > 0$ , there is a sequence of open intervals,  $(a_n, b_n)$  such that  $A \subseteq \bigcup (a_n, b_n)$  and  $\sum (b_n - a_n) < \varepsilon$ .

Clearly we have the following assertion.

**Lemma 3.2.** *If  $(A_n)$  is a sequence of null sets, then so is  $\bigcup A_n$ . Consequently, all countable sets are null sets.*

From now on, we use the following notation in the rest of this section.

- (1) For each subset  $A$  of  $\mathbb{R}$ , put  $\omega(f, A) := \sup\{|f(x) - f(x')| : x, x' \in A\}$ .
- (2) For  $c \in [a, b]$ , put  $\omega(f, c) := \inf\{\omega(f, B(c, r)) : r > 0\}$ , where  $B(c, r) := (c - r, c + r)$ .

The following is easy shown directly from the definition.

**Lemma 3.3.** *The function  $f$  is continuous at  $c \in [a, b]$  if and only if  $\omega(f, c) = 0$ .*

**Theorem 3.4. Lebesgue integrability theorem:** *Retains the notation as above. Let  $D := \{c \in [a, b] : f \text{ is discontinuous at } c\}$ . Then  $f \in R[a, b]$  if and only if  $D$  has measure zero.*

*Proof.* For each positive integer  $n$ , let  $D_n := \{x \in [a, b] : \omega(f, x) \geq \frac{1}{n}\}$ . Then we have  $D = \bigcup_{n=1}^{\infty} D_n$ .

For  $(\Rightarrow)$ , assume that  $f \in R[a, b]$ . Then by Lemma 3.2, it suffices to show that each  $D_n$  is a null set. Fix a positive integer  $m$  such that  $D_m \neq \emptyset$ . Now let  $\varepsilon > 0$ . Since  $f \in R[a, b]$ , there is a partition  $P : a = x_0 < \cdots < x_n = b$  such that  $\sum \omega_k(f, P) \Delta x_k < \frac{\varepsilon}{m}$ . Notice that  $c \in D_m$  if and only if  $\omega(f, B(c, \delta)) \geq \frac{1}{m}$  for all  $\delta > 0$ , where  $B(c, \delta) := (c - \delta, c + \delta)$ . Thus, if  $[x_{k-1}, x_k] \cap D_m \neq \emptyset$ , then  $\omega_k(f, P) \geq \frac{1}{m}$ . This implies that

$$\begin{aligned} \frac{\varepsilon}{m} &> \sum_{k=1}^n \omega_k(f, P) \Delta x_k \\ &\geq \sum_{k: [x_{k-1}, x_k] \cap D_m \neq \emptyset} \omega_k(f, P) \Delta x_k \\ &\geq \frac{1}{m} \sum_{k: [x_{k-1}, x_k] \cap D_m \neq \emptyset} \Delta x_k. \end{aligned}$$

Therefore, we have  $D_m \subseteq \bigcup_{k: [x_{k-1}, x_k] \cap D_m \neq \emptyset} [x_{k-1}, x_k]$  and

$$\sum_{k: [x_{k-1}, x_k] \cap D_m \neq \emptyset} \Delta x_k < \varepsilon.$$

Thus,  $D_m$  is a null set for each positive integer  $m$  as desired.

Now for showing  $(\Leftarrow)$ , assume that the set  $D$  of all discontinuous points of  $f$  is a null set.

We first claim that each  $D_m$  is a closed set. To see this, note that a point  $c \in D_m$  if and only if  $\omega(f, B(c, r)) \geq \frac{1}{m}$  for all  $r > 0$  if and only if for all  $\eta > 0$  and for all  $r > 0$ , there are points  $x', x'' \in B(c, r)$  such that  $|f(x') - f(x'')| > \frac{1}{m} - \eta$ . Now let  $(c_n)$  be a sequence in  $D_m$  converging to a point  $c$ . Let  $r > 0$  and  $\eta > 0$ . Then there is  $c_N$  such that  $|c_N - c| < \frac{r}{2}$ . Since  $c_N \in D_m$ , there are  $x', x'' \in B(c_N, \frac{r}{2})$  such that  $|f(x') - f(x'')| > \frac{1}{m} - \eta$ . Since  $x', x'' \in B(c_N, \frac{r}{2})$ ,  $x', x'' \in B(c, r)$ . Thus,  $c \in D_m$  is as desired. This shows that  $D_m$  is a closed subset of  $[a, b]$ , and hence it is compact.

Let  $\varepsilon > 0$  and let  $m$  be a positive integer such that  $1/m < \varepsilon$ . By the assumption  $D = \bigcup_{l=1}^{\infty} D_l$  is a null set and so is the set  $D_m$ . Then there is a sequence of open intervals, say  $\{(a_j, b_j)\}$ , such that  $D_m \subseteq \bigcup (a_j, b_j)$  and  $\sum (b_j - a_j) < \varepsilon$ . Since  $D_m$  is compact, there are finitely many  $(a_j, b_j)$ 's for  $j = 1, \dots, K$  such that  $D_m \subseteq \bigcup_{j=1}^K (a_j, b_j)$ . Note that we may assume that the sequence  $a_1 < b_1 < a_2 <$

$b_2 < \cdots < a_K < b_K$ . Choose a partition  $Q := \{a_j, b_j : j = 1, \dots, K\} \cup \{a, b\}$  on  $[a, b]$  and rewrite  $Q$  as  $a = x_0 < \cdots < x_n = b$ . Let  $J = (a_1, b_1) \cup \cdots \cup (a_K, b_K)$ .

Put  $I := \{j : [x_{j-1}, x_j] \cap J = \emptyset\}$  and  $II := \{j : [x_{j-1}, x_j] \cap J \neq \emptyset\}$ .

Note that if  $j \in I$ , then  $\omega(f, x) < \frac{1}{m}$  for all  $x \in [x_{j-1}, x_j]$ . Hence, for each  $x \in [x_{j-1}, x_j]$ , there is  $\delta_x > 0$  such that  $\omega(f, B(x, \delta_x)) < \frac{1}{m}$ . Then by the compactness of  $[x_{j-1}, x_j]$ , there is a partition  $P'_j : x_{j-1} = x'_0 < \cdots < x'_l = x_j$  on  $[x_{j-1}, x_j]$  such that  $\omega_{j'}(f, P'_j) < \frac{1}{m}$  for all  $j' = 1, \dots, l$ . Thus, we have  $\sum_{j'} \omega_{j'}(f, P'_j) \Delta x_{j'} < \frac{1}{m} (x_j - x_{j-1}) < \varepsilon (x_{j-1} - x_j)$  whenever  $j \in I$ .

On the other hand, if  $j \in II$ , then  $[x_{j-1}, x_j] \cap J \neq \emptyset$ . Since  $\sum_{j=1}^K (b_j - a_j) < \varepsilon$ , we see that  $\sum_{j \in II} \omega_j(f, Q) \Delta x_j < 2M\varepsilon$ .

Now put  $P := Q \cup \bigcup_{j \in I} P'_j : a = y_0 < \cdots < y_N = b$ . From the above argument, we have shown that

$\sum_{i=1}^N \omega_i(f, P) \Delta y_i < \varepsilon(b - a) + 2M\varepsilon$ . Thus  $f \in R[a, b]$ . The proof is complete.  $\square$

## 4. SOME RESULTS OF SEQUENCES OF FUNCTIONS

**Proposition 4.1.** *Let  $f_n : (a, b) \rightarrow \mathbb{R}$  be a sequence of functions. Assume that it satisfies the following conditions:*

- (i) :  $f_n(x)$  point-wise converges to a function  $f(x)$  on  $(a, b)$ ;
- (ii) : each  $f_n$  is a  $C^1$  function on  $(a, b)$ ;
- (iii) :  $f'_n \rightarrow g$  uniformly on  $(a, b)$ .

Then  $f$  is a  $C^1$ -function on  $(a, b)$  with  $f' = g$ .

*Proof.* Fix  $c \in (a, b)$ . Then for each  $x$  with  $c < x < b$  (similarly, we can prove it in the same way as  $a < x < c$ ), the Fundamental Theorem of Calculus implies that

$$f_n(x) = \int_c^x f'_n(t)dt + f_n(c).$$

Since  $f'_n \rightarrow g$  uniformly on  $(a, b)$ , we see that

$$\int_c^x f'_n(t)dt \rightarrow \int_c^x g(t)dt.$$

This gives

$$(4.1) \quad f(x) = \int_c^x g(t)dt + f(c).$$

for all  $x \in (c, b)$ . Similarly, we have  $f(x) = \int_c^x g(t)dt + f(c)$  for all  $x \in (a, b)$ .

On the other hand,  $g$  is continuous on  $(a, b)$  since each  $f'_n$  is continuous and  $f'_n \rightarrow g$  uniformly on  $(a, b)$ . Equation 4.1 will tell us that  $f'$  exists and  $f' = g$  on  $(a, b)$ . The proof is finished.  $\square$

**Proposition 4.2.** *Let  $(f_n)$  be a sequence of differentiable functions defined on  $(a, b)$ . Assume that*

- (i): there is a point  $c \in (a, b)$  such that  $\lim f_n(c)$  exists;
- (ii):  $f'_n$  converges uniformly to a function  $g$  on  $(a, b)$ .

Then

- (a):  $f_n$  converges uniformly to a function  $f$  on  $(a, b)$ ;
- (b):  $f$  is differentiable on  $(a, b)$  and  $f' = g$ .

*Proof.* For Part (a), we will make use the Cauchy theorem.

Let  $\varepsilon > 0$ . Then by the assumptions (i) and (ii), there is a positive integer  $N$  such that

$$|f_m(c) - f_n(c)| < \varepsilon \quad \text{and} \quad |f'_m(x) - f'_n(x)| < \varepsilon$$

for all  $m, n \geq N$  and for all  $x \in (a, b)$ . Now fix  $c < x < b$  and  $m, n \geq N$ . To apply the Mean Value Theorem for  $f_m - f_n$  on  $(c, x)$ , then there is a point  $\xi$  between  $c$  and  $x$  such that

$$(4.2) \quad f_m(x) - f_n(x) = f_m(c) - f_n(c) + (f'_m(\xi) - f'_n(\xi))(x - c).$$

This implies that

$$|f_m(x) - f_n(x)| \leq |f_m(c) - f_n(c)| + |f'_m(\xi) - f'_n(\xi)||x - c| < \varepsilon + (b - a)\varepsilon$$

for all  $m, n \geq N$  and for all  $x \in (c, b)$ . Similarly, when  $x \in (a, c)$ , we also have

$$|f_m(x) - f_n(x)| < \varepsilon + (b - a)\varepsilon.$$

So Part (a) follows.

Let  $f$  be the uniform limit of  $(f_n)$  on  $(a, b)$

For Part (b), we fix  $u \in (a, b)$ . We are going to show

$$\lim_{x \rightarrow u} \frac{f(x) - f(u)}{x - u} = g(u).$$



Let  $\varepsilon > 0$ . Since  $(f'_n)$  is uniformly convergent on  $(a, b)$ , there is  $N \in \mathbb{N}$  such that

$$(4.3) \quad |f'_m(x) - f'_n(x)| < \varepsilon$$

for all  $m, n \geq N$  and for all  $x \in (a, b)$

Note that for all  $m \geq N$  and  $x \in (a, b) \setminus \{u\}$ , applying the Mean value Theorem for  $f_m - f_N$  as before, we have

$$\frac{f_m(x) - f_N(x)}{x - u} = \frac{f_m(u) - f_N(u)}{x - u} + (f'_m(\xi) - f'_N(\xi))$$

for some  $\xi$  between  $u$  and  $x$ .

So Eq.4.3 implies that

$$(4.4) \quad \left| \frac{f_m(x) - f_m(u)}{x - u} - \frac{f_N(x) - f_N(u)}{x - u} \right| \leq \varepsilon$$

for all  $m \geq N$  and for all  $x \in (a, b)$  with  $x \neq u$ .

Taking  $m \rightarrow \infty$  in Eq.4.4, we have

$$\left| \frac{f(x) - f(u)}{x - u} - \frac{f_N(x) - f_N(u)}{x - u} \right| \leq \varepsilon.$$

Hence we have

$$\begin{aligned} \left| \frac{f(x) - f(u)}{x - u} - f'_N(u) \right| &\leq \left| \frac{f(x) - f(u)}{x - u} - \frac{f_N(x) - f_N(u)}{x - u} \right| + \left| \frac{f_N(x) - f_N(u)}{x - u} - f'_N(u) \right| \\ &\leq \varepsilon + \left| \frac{f_N(x) - f_N(u)}{x - u} - f'_N(u) \right|. \end{aligned}$$

So if we can take  $0 < \delta$  such that  $\left| \frac{f_N(x) - f_N(u)}{x - u} - f'_N(u) \right| < \varepsilon$  for  $0 < |x - u| < \delta$ , then we have

$$(4.5) \quad \left| \frac{f(x) - f(u)}{x - u} - f'_N(u) \right| \leq 2\varepsilon$$

for  $0 < |x - u| < \delta$ . On the other hand, by the choice of  $N$ , we have  $|f'_m(y) - f'_N(y)| < \varepsilon$  for all  $y \in (a, b)$  and  $m \geq N$ . So we have  $|g(u) - f'_N(u)| \leq \varepsilon$ . This together with Eq.4.5 give

$$\left| \frac{f(x) - f(u)}{x - u} - g(u) \right| \leq 3\varepsilon$$

as  $0 < |x - u| < \delta$ , that is we have

$$\lim_{x \rightarrow u} \frac{f(x) - f(u)}{x - u} = g(u).$$

The proof is finished. □

**Remark 4.3.** *The uniform convergence assumption of  $(f'_n)$  in the Propositions above is essential.*

**Example 4.4.** *Let  $f_n(x) := \frac{x}{1+n^2x^2}$  for  $x \in (-1, 1)$ . Then we have*

$$g(x) := \lim_n f'_n(x) := \lim_n \frac{1 - n^2x^2}{(1 + n^2x^2)^2} = \begin{cases} 0 & \text{if } x \neq 0; \\ 1 & \text{if } x = 0. \end{cases}$$

*On the other hand,  $f_n \rightarrow 0$  uniformly on  $(-1, 1)$ . In fact, if  $f'_n(1/n) = 0$  for all  $n = 1, 2, \dots$ , then  $f_n$  attains the maximal value  $f_n(1/n) = \frac{1}{2n}$  at  $x = 1/n$  for each  $n = 1, \dots$  and hence,  $f_n \rightarrow 0$  uniformly on  $(-1, 1)$ .*

*So Propositions 4.1 and 4.2 does not hold. Note that  $(f'_n)$  does not converge uniformly to  $g$  on  $(-1, 1)$ .*

**Proposition 4.5. (Dini's Theorem):** Let  $A$  be a compact subset of  $\mathbb{R}$  and  $f_n : A \rightarrow \mathbb{R}$  be a sequence of continuous functions defined on  $A$ . Suppose that

- (i) for each  $x \in A$ , we have  $f_n(x) \leq f_{n+1}(x)$  for all  $n = 1, 2, \dots$ ;
- (ii) the pointwise limit  $f(x) := \lim_n f_n(x)$  exists for all  $x \in A$ ;
- (iii)  $f$  is continuous on  $A$ .

Then  $f_n$  converges to  $f$  uniformly on  $A$ .

*Proof.* Let  $g_n := f - f_n$  defined on  $A$ . Then each  $g_n$  is continuous and  $g_n(x) \downarrow 0$  pointwise on  $A$ . It suffices to show that  $g_n$  converges to 0 uniformly on  $A$ .

**Method I:** Suppose not. Then there is  $\varepsilon > 0$  such that for all positive integer  $N$ , we have

$$(4.6) \quad g_n(x_n) \geq \varepsilon.$$

for some  $n \geq N$  and some  $x_n \in A$ . From this, by passing to a subsequence we may assume that  $g_n(x_n) \geq \varepsilon$  for all  $n = 1, 2, \dots$ . Then by using the compactness of  $A$ , there is a convergent subsequence  $(x_{n_k})$  of  $(x_n)$  in  $A$ . Let  $z := \lim_k x_{n_k} \in A$ . Since  $g_{n_k}(z) \downarrow 0$  as  $k \rightarrow \infty$ . So, there is a positive integer  $K$  such that  $0 \leq g_{n_K}(z) < \varepsilon/2$ . Since  $g_{n_K}$  is continuous at  $z$  and  $\lim_i x_{n_i} = z$ , we have  $\lim_i g_{n_K}(x_{n_i}) = g_{n_K}(z)$ . So, we can choose  $i$  large enough such that  $i > K$

$$g_{n_i}(x_{n_i}) \leq g_{n_K}(x_{n_i}) < \varepsilon/2$$

because  $g_m(x_{n_i}) \downarrow 0$  as  $m \rightarrow \infty$ . This contradicts to the Inequality 4.6.

**Method II:** Let  $\varepsilon > 0$ . Fix  $x \in A$ . Since  $g_n(x) \downarrow 0$ , there is  $N(x) \in \mathbb{N}$  such that  $0 \leq g_n(x) < \varepsilon$  for all  $n \geq N(x)$ . Since  $g_{N(x)}$  is continuous, there is  $\delta(x) > 0$  such that  $g_{N(x)}(y) < \varepsilon$  for all  $y \in A$  with  $|x - y| < \delta(x)$ . If we put  $J_x := (x - \delta(x), x + \delta(x))$ , then  $A \subseteq \bigcup_{x \in A} J_x$ . Then by the compactness of  $A$ , there are finitely many  $x_1, \dots, x_m$  in  $A$  such that  $A \subseteq J_{x_1} \cup \dots \cup J_{x_m}$ . Put  $N := \max(N(x_1), \dots, N(x_m))$ . Now if  $y \in A$ , then  $y \in J(x_i)$  for some  $1 \leq i \leq m$ . This implies that

$$g_n(y) \leq g_{N(x_i)}(y) < \varepsilon$$

for all  $n \geq N \geq N(x_i)$ . □

## 5. ABSOLUTELY CONVERGENT SERIES

Throughout this section, let  $(a_n)$  be a sequence of complex numbers.

**Definition 5.1.** We say that a series  $\sum_{n=1}^{\infty} a_n$  is absolutely convergent if  $\sum_{n=1}^{\infty} |a_n| < \infty$ .

Also a convergent series  $\sum_{n=1}^{\infty} a_n$  is said to be conditionally convergent if it is not absolute convergent.

**Example 5.2. Important Example :** The series  $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^\alpha}$  is conditionally convergent when

$$0 < \alpha \leq 1.$$

This example shows us that a convergent improper integral may fail to the absolute convergence or square integrable property.

For instance, if we consider the function  $f : [1, \infty) \rightarrow \mathbb{R}$  given by

$$f(x) = \frac{(-1)^{n+1}}{n^\alpha} \quad \text{if } n \leq x < n + 1.$$

If  $\alpha = 1/2$ , then  $\int_1^{\infty} f(x)dx$  is convergent but it is neither absolutely convergent nor square integrable.

**Notation 5.3.** Let  $\sigma : \{1, 2, \dots\} \rightarrow \{1, 2, \dots\}$  be a bijection. A formal series  $\sum_{n=1}^{\infty} a_{\sigma(n)}$  is called an rearrangement of  $\sum_{n=1}^{\infty} a_n$ .

**Example 5.4.** In this example, we are going to show that there is an rearrangement of the series  $\sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i}$  is divergent although the original series is convergent. In fact, it is conditionally convergent.

We first notice that the series  $\sum_i \frac{1}{2i-1}$  diverges to infinity. Thus for each  $M > 0$ , there is a positive integer  $N$  such that

$$\sum_{i=1}^n \frac{1}{2i-1} \geq M \quad \dots\dots\dots (*)$$

for all  $n \geq N$ . Then there is  $N_1 \in \mathbb{N}$  such that

$$\sum_{i=1}^{N_1} \frac{1}{2i-1} - \frac{1}{2} > 1.$$

By using (\*) again, there is a positive integer  $N_2$  with  $N_1 < N_2$  such that

$$\sum_{i=1}^{N_1} \frac{1}{2i-1} - \frac{1}{2} + \sum_{N_1 < i \leq N_2} \frac{1}{2i-1} - \frac{1}{4} > 2.$$

To repeat the same procedure, we can find a positive integers subsequence  $(N_k)$  such that

$$\sum_{i=1}^{N_1} \frac{1}{2i-1} - \frac{1}{2} + \sum_{N_1 < i \leq N_2} \frac{1}{2i-1} - \frac{1}{4} + \dots\dots\dots - \sum_{N_{k-1} < i \leq N_k} \frac{1}{2i-1} - \frac{1}{2k} > k$$

for all positive integers  $k$ . So if we let  $a_n = \frac{(-1)^{n+1}}{n}$ , then one can find a bijection  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that the series  $\sum_{i=1}^{\infty} a_{\sigma(i)}$  is an rearrangement of the series  $\sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i}$  and diverges to infinity. The proof is finished.

**Theorem 5.5.** Let  $\sum_{n=1}^{\infty} a_n$  be an absolutely convergent series. Then for any rearrangement  $\sum_{n=1}^{\infty} a_{\sigma(n)}$

is also absolutely convergent. Moreover, we have  $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} a_{\sigma(n)}$ .

*Proof.* Let  $\sigma : \{1, 2, \dots\} \rightarrow \{1, 2, \dots\}$  be a bijection as before.

We first claim that  $\sum_n a_{\sigma(n)}$  is also absolutely convergent.

Let  $\varepsilon > 0$ . Since  $\sum_n |a_n| < \infty$ , there is a positive integer  $N$  such that

$$|a_{N+1}| + \dots\dots\dots + |a_{N+p}| < \varepsilon \quad \dots\dots\dots (*)$$

for all  $p = 1, 2, \dots$ . Notice that since  $\sigma$  is a bijection, we can find a positive integer  $M$  such that  $M > \max\{j : 1 \leq \sigma(j) \leq N\}$ . Then  $\sigma(i) \geq N$  if  $i \geq M$ . This together with (\*) imply that if  $i \geq M$  and  $p \in \mathbb{N}$ , we have

$$|a_{\sigma(i+1)}| + \dots\dots\dots |a_{\sigma(i+p)}| < \varepsilon.$$

Thus the series  $\sum_n a_{\sigma(n)}$  is absolutely convergent by the Cauchy criteria.

Finally we claim that  $\sum_n a_n = \sum_n a_{\sigma(n)}$ . Put  $l = \sum_n a_n$  and  $l' = \sum_n a_{\sigma(n)}$ . Now let  $\varepsilon > 0$ . Then there is  $N \in \mathbb{N}$  such that

$$|l - \sum_{n=1}^N a_n| < \varepsilon \quad \text{and} \quad |a_{N+1}| + \dots + |a_{N+p}| < \varepsilon \dots \dots \dots (**)$$

for all  $p \in \mathbb{N}$ . Now choose a positive integer  $M$  large enough so that  $\{1, \dots, N\} \subseteq \{\sigma(1), \dots, \sigma(M)\}$  and  $|l' - \sum_{i=1}^M a_{\sigma(i)}| < \varepsilon$ . Notice that since we have  $\{1, \dots, N\} \subseteq \{\sigma(1), \dots, \sigma(M)\}$ , the condition  $(**)$  gives

$$|\sum_{n=1}^N a_n - \sum_{i=1}^M a_{\sigma(i)}| \leq \sum_{N < i < \infty} |a_i| \leq \varepsilon.$$

We can now conclude that

$$|l - l'| \leq |l - \sum_{n=1}^N a_n| + |\sum_{n=1}^N a_n - \sum_{i=1}^M a_{\sigma(i)}| + |\sum_{i=1}^M a_{\sigma(i)} - l'| \leq 3\varepsilon.$$

The proof is complete. □

### 6. POWER SERIES

Throughout this section, let

$$f(x) = \sum_{i=0}^{\infty} a_i x^i \quad \dots \dots \dots (*)$$

denote a formal power series, where  $a_i \in \mathbb{R}$ .

**Lemma 6.1.** *Suppose that there is  $c \in \mathbb{R}$  with  $c \neq 0$  such that  $f(c)$  is convergent. Then*

- (i) :  $f(x)$  is absolutely convergent for all  $x$  with  $|x| < |c|$ .
- (ii) :  $f$  converges uniformly on  $[-\eta, \eta]$  for any  $0 < \eta < |c|$ .

*Proof.* For Part (i), note that since  $f(c)$  is convergent, then  $\lim a_n c^n = 0$ . So there is a positive integer  $N$  such that  $|a_n c^n| \leq 1$  for all  $n \geq N$ . Now if we fix  $|x| < |c|$ , then  $|x/c| < 1$ . Therefore, we have

$$\sum_{n=1}^{\infty} |a_n| |x^n| \leq \sum_{n=1}^{N-1} |a_n| |x^n| + \sum_{n \geq N} |a_n c^n| |x/c|^n \leq \sum_{n=1}^{N-1} |a_n| |x^n| + \sum_{n \geq N} |x/c|^n < \infty.$$

So Part (i) follows.

Now for Part (ii), if we fix  $0 < \eta < |c|$ , then  $|a_n x^n| \leq |a_n \eta^n|$  for all  $n$  and for all  $x \in [-\eta, \eta]$ . On the other hand, we have  $\sum_n |a_n \eta^n| < \infty$  by Part (i). So  $f$  converges uniformly on  $[-\eta, \eta]$  by the  $M$ -test. The proof is finished. □

**Remark 6.2.** *In Lemma 6.9(ii), notice that if  $f(c)$  is convergent, it does not imply  $f$  converges uniformly on  $[-c, c]$  in general.*

*For example,  $f(x) := 1 + \sum_{n=1}^{\infty} \frac{x^n}{n}$ . Then  $f(-1)$  is convergent but  $f(1)$  is divergent.*

**Definition 6.3.** *Call the set  $\text{dom } f := \{x \in \mathbb{R} : f(x) \text{ is convergent}\}$  the domain of convergence of  $f$  for convenience. Let  $0 \leq r := \sup\{|c| : c \in \text{dom } f\} \leq \infty$ . Then  $r$  is called the radius of convergence of  $f$ .*

**Remark 6.4.** Notice that by Lemma 6.9, then the domain of convergence of  $f$  must be the interval with the end points  $\pm r$  if  $0 < r < \infty$ .

When  $r = 0$ , then  $\text{dom } f = \{0\}$ .

Finally, if  $r = \infty$ , then  $\text{dom } f = \mathbb{R}$ .

**Example 6.5.** If  $f(x) = \sum_{n=0}^{\infty} n!x^n$ , then  $r = (0)$ . In fact, notice that if we fix a non-zero number  $x$  and consider  $\lim_n |(n+1)!x^{n+1}|/|n!x^n| = \infty$ , then by the ratio test  $f(x)$  must be divergent for any  $x \neq 0$ . So  $r = 0$  and  $\text{dom } f = (0)$ .

**Example 6.6.** Let  $f(x) = 1 + \sum_{n=1}^{\infty} x^n/n^n$ . Notice that we have  $\lim_n |x^n/n^n|^{1/n} = 0$  for all  $x$ . So the root test implies that  $f(x)$  is convergent for all  $x$  and then  $r = \infty$  and  $\text{dom } f = \mathbb{R}$ .

**Example 6.7.** Let  $f(x) = 1 + \sum_{n=1}^{\infty} x^n/n$ . Then  $\lim_n |x^{n+1}/(n+1)| \cdot |n/x^n| = |x|$  for all  $x \neq 0$ . So by the ration test, we see that if  $|x| < 1$ , then  $f(x)$  is convergent and if  $|x| > 1$ , then  $f(x)$  is divergent. So  $r = 1$ . Also, it is known that  $f(1)$  is divergent but  $f(-1)$  is convergent. Therefore, we have  $\text{dom } f = [-1, 1)$ .

**Example 6.8.** Let  $f(x) = \sum x^n/n^2$ . Then by using the same argument of Example 6.7, we have  $r = 1$ . On the other hand, it is known that  $f(\pm 1)$  both are convergent. So  $\text{dom } f = [-1, 1]$ .

**Lemma 6.9.** With the notation as above, if  $r > 0$ , then  $f$  converges uniformly on  $(-\eta, \eta)$  for any  $0 < \eta < r$ .

*Proof.* It follows from Lemma 6.1 at once. □

**Remark 6.10.** Note that the Example 6.7 shows us that  $f$  may not converge uniformly on  $(-r, r)$ . In fact let  $f$  be defined as in Example 6.7. Then  $f$  does not converges on  $(-1, 1)$ . In fact, if we let  $s_n(x) = \sum_{k=0}^{\infty} a_k x^k$ , then for any positive integer  $n$  and  $0 < x < 1$ , we have

$$|s_{2n}(x) - s_n(x)| = \frac{x^{n+1}}{n+1} + \cdots + \frac{x^n}{2n}.$$

From this we see that if  $n$  is fixed, then  $|s_{2n}(x) - s_n(x)| \rightarrow 1/2$  as  $x \rightarrow 1-$ . So for each  $n$ , we can find  $0 < x < 1$  such that  $|s_{2n}(x) - s_n(x)| > \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$ . Thus  $f$  does not converges uniformly on  $(-1, 1)$  by the Cauchy Theorem.

**Proposition 6.11.** With the notation as above, let  $\ell = \overline{\lim} |a_n|^{1/n}$  or  $\lim \frac{|a_{n+1}|}{|a_n|}$  provided it exists.

Then

$$r = \begin{cases} \frac{1}{\ell} & \text{if } 0 < \ell < \infty; \\ 0 & \text{if } \ell = \infty; \\ \infty & \text{if } \ell = 0. \end{cases}$$

**Proposition 6.12.** With the notation as above if  $0 < r \leq \infty$ , then  $f \in C^\infty(-r, r)$ . Moreover, the  $k$ -derivatives  $f^{(k)}(x) = \sum_{n \geq k} a_k n(n-1)(n-2) \cdots (n-k+1)x^{n-k}$  for all  $x \in (-r, r)$ .

*Proof.* Fix  $c \in (-r, r)$ . By Lemma 6.9, one can choose  $0 < \eta < r$  such that  $c \in (-\eta, \eta)$  and  $f$  converges uniformly on  $(-\eta, \eta)$ .

It needs to show that the  $k$ -derivatives  $f^{(k)}(c)$  exists for all  $k \geq 0$ . Consider the case  $k = 1$  first.

If we consider the series  $\sum_{n=0}^{\infty} (a_n x^n)' = \sum_{n=1}^{\infty} n a_n x^{n-1}$ , then it also has the same radius  $r$  because  $\lim_n |n a_n|^{1/n} = \lim_n |a_n|^{1/n}$ . This implies that the series  $\sum_{n=1}^{\infty} n a_n x^{n-1}$  converges uniformly

on  $(-\eta, \eta)$ . Therefore, the restriction  $f|_{(-\eta, \eta)}$  is differentiable. In particular,  $f'(c)$  exists and  $f'(c) = \sum_{n=1}^{\infty} na_n c^{n-1}$ .  
So the result can be shown inductively on  $k$ .  $\square$

**Proposition 6.13.** *With the notation as above, suppose that  $r > 0$ . Then we have*

$$\int_0^x f(t)dt = \sum_{n=0}^{\infty} \int_0^x a_n t^n dt = \sum_{n=0}^{\infty} \frac{1}{n+1} a_n x^{n+1}$$

for all  $x \in (-r, r)$ .

*Proof.* Fix  $0 < x < r$ . Then by Lemma 6.9  $f$  converges uniformly on  $[0, x]$ . Since each term  $a_n t^n$  is continuous, the result follows.  $\square$

**Theorem 6.14. (Abel) :** *With the notation as above, suppose that  $0 < r$  and  $f(r)$  (or  $f(-r)$ ) exists. Then  $f$  is continuous at  $x = r$  (resp.  $x = -r$ ), that is  $\lim_{x \rightarrow r^-} f(x) = f(r)$ .*

*Proof.* Note that by considering  $f(-x)$ , it suffices to show that the case  $x = r$  holds. Assume  $r = 1$ .

Notice that if  $f$  converges uniformly on  $[0, 1]$ , then  $f$  is continuous at  $x = 1$  as desired. Let  $\varepsilon > 0$ . Since  $f(1)$  is convergent, then there is a positive integer such that

$$|a_{n+1} + \dots + a_{n+p}| < \varepsilon$$

for  $n \geq N$  and for all  $p = 1, 2, \dots$ . Note that for  $n \geq N$ ;  $p = 1, 2, \dots$  and  $x \in [0, 1]$ , we have

$$\begin{aligned} s_{n+p}(x) - s_n(x) &= a_{n+1}x^{n+1} + a_{n+2}x^{n+1} + a_{n+3}x^{n+1} + \dots + a_{n+p}x^{n+1} \\ &\quad + a_{n+2}(x^{n+2} - x^{n+1}) + a_{n+3}(x^{n+2} - x^{n+1}) + \dots + a_{n+p}(x^{n+2} - x^{n+1}) \\ (6.1) \quad &\quad + a_{n+3}(x^{n+3} - x^{n+2}) + \dots + a_{n+p}(x^{n+3} - x^{n+2}) \\ &\quad \vdots \\ &\quad + a_{n+p}(x^{n+p} - x^{n+p-1}). \end{aligned}$$

Since  $x \in [0, 1]$ ,  $|x^{n+k+1} - x^{n+k}| = x^{n+k} - x^{n+k+1}$ . So the Eq.6.1 implies that

$$|s_{n+p}(x) - s_n(x)| \leq \varepsilon(x_{n+1} + (x^{n+1} - x^{n+2}) + (x^{n+2} - x^{n+3}) + \dots + (x^{n+p-1} - x^{n+p})) = \varepsilon(2x^{n+1} - x^{n+p}) \leq 2\varepsilon.$$

So  $f$  converges uniformly on  $[0, 1]$  as desired.

Finally for the general case, we consider  $g(x) := f(rx) = \sum_n a_n r^n x^n$ . Note that  $\lim_n |a_n r^n|^{1/n} = 1$  and  $g(1) = f(r)$ . Then by the case above, we have shown that

$$f(r) = g(1) = \lim_{x \rightarrow 1^-} g(x) = \lim_{x \rightarrow r^-} f(x).$$

The proof is finished.  $\square$

**Remark 6.15.** *In Remark 6.10, we have seen that  $f$  may not converges uniformly on  $(-r, r)$ . However, in the proof of Abel's Theorem above, we have shown that if  $f(\pm r)$  both exist, then  $f$  converges uniformly on  $[-r, r]$  in this case.*

7. REAL ANALYTIC FUNCTIONS

**Proposition 7.1.** *Let  $f \in C^\infty(a, b)$  and  $c \in (a, b)$ . Then for any  $x \in (a, b) \setminus \{c\}$  and for any  $n \in \mathbb{N}$ , there is  $\xi = \xi(x, n)$  between  $c$  and  $x$  such that*

$$f(x) = \sum_{k=0}^n \frac{f^{(k)}(c)}{k!} (x - c)^k + \int_c^x \frac{f^{(n+1)}(t)}{n!} (x - t)^n dt$$

Call  $\sum_{k=0}^\infty \frac{f^{(k)}(c)}{k!} (x - c)^k$  (may not be convergent) the Taylor series of  $f$  at  $c$ .

*Proof.* It is easy to prove by induction on  $n$  and the integration by part. □

**Definition 7.2.** *A real-valued function  $f$  defined on  $(a, b)$  is said to be real analytic if for each  $c \in (a, b)$ , one can find  $\delta > 0$  and a power series  $\sum_{k=0}^\infty a_k(x - c)^k$  such that*

$$f(x) = \sum_{k=0}^\infty a_k(x - c)^k \quad \dots\dots\dots (*)$$

for all  $x \in (c - \delta, c + \delta) \subseteq (a, b)$ .

**Remark 7.3.**

(i) : *Concerning about the definition of a real analytic function  $f$ , the expression (\*) above is uniquely determined by  $f$ , that is, each coefficient  $a_k$ 's is uniquely determined by  $f$ . In fact, by Proposition 6.12, we have seen that  $f \in C^\infty(a, b)$  and*

$$a_k = \frac{f^{(k)}(c)}{k!} \quad \dots\dots\dots (**)$$

for all  $k = 0, 1, 2, \dots$

(ii) : *Although every real analytic function is  $C^\infty$ , the following example shows that the converse does not hold.*

Define a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0; \\ 0 & \text{if } x = 0. \end{cases}$$

*One can directly check that  $f \in C^\infty(\mathbb{R})$  and  $f^{(k)}(0) = 0$  for all  $k = 0, 1, 2, \dots$ . So if  $f$  is real analytic, then there is  $\delta > 0$  such that  $a_k = 0$  for all  $k$  by the Eq.(\*\*) above and hence  $f(x) \equiv 0$  for all  $x \in (-\delta, \delta)$ . It is absurd.*

(iii) **Interesting Fact** : *Let  $D$  be an open disc in  $\mathbb{C}$ . A complex analytic function  $f$  on  $D$  is similarly defined as in the real case. However, we always have:  $f$  is complex analytic if and only if it is  $C^\infty$ .*

**Proposition 7.4.** *Suppose that  $f(x) := \sum_{k=0}^\infty a_k(x - c)^k$  is convergent on some open interval  $I$  centered at  $c$ , that is  $I = (c - r, c + r)$  for some  $r > 0$ . Then  $f$  is analytic on  $I$ .*

*Proof.* We first note that  $f \in C^\infty(I)$ . By considering the translation  $x - c$ , we may assume that  $c = 0$ . Now fix  $z \in I$ . Now choose  $\delta > 0$  such that  $(z - \delta, z + \delta) \subseteq I$ . We are going to show that

$$f(x) = \sum_{j=0}^\infty \frac{f^{(j)}(z)}{j!} (x - z)^j.$$

for all  $x \in (z - \delta, z + \delta)$ .

Notice that  $f(x)$  is absolutely convergent on  $I$ . This implies that

$$\begin{aligned} f(x) &= \sum_{k=0}^{\infty} a_k (x - z + z)^k \\ &= \sum_{k=0}^{\infty} a_k \sum_{j=0}^k \frac{k(k-1)\cdots(k-j+1)}{j!} (x-z)^j z^{k-j} \\ &= \sum_{j=0}^{\infty} \left( \sum_{k \geq j} k(k-1)\cdots(k-j+1) a_k z^{k-j} \right) \frac{(x-z)^j}{j!} \\ &= \sum_{j=0}^{\infty} \frac{f^{(j)}(z)}{j!} (x-z)^j \end{aligned}$$

for all  $x \in (z - \delta, z + \delta)$ . The proof is finished.  $\square$

**Example 7.5.** Let  $\alpha \in \mathbb{R}$ . Recall that  $(1+x)^\alpha$  is defined by  $e^{\alpha \ln(1+x)}$  for  $x > -1$ .

Now for each  $k \in \mathbb{N}$ , put

$$\binom{\alpha}{k} = \begin{cases} \frac{\alpha(\alpha-1)\cdots(\alpha-k+1)}{k!} & \text{if } k \neq 0; \\ 1 & \text{if } k = 0. \end{cases}$$

Then

$$f(x) := (1+x)^\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} x^k$$

whenever  $|x| < 1$ .

Consequently,  $f(x)$  is analytic on  $(-1, 1)$ .

*Proof.* Notice that  $f^{(k)}(x) = \alpha(\alpha-1)\cdots(\alpha-k+1)(1+x)^{\alpha-k}$  for  $|x| < 1$ .

Fix  $|x| < 1$ . Then by Proposition 7.1, for each positive integer  $n$  we have

$$f(x) = \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{k!} x^k + \int_0^x \frac{f^{(n)}(t)}{(n-1)!} (x-t)^{n-1} dt$$

So by the mean value theorem for integrals, for each positive integer  $n$ , there is  $\xi_n$  between 0 and  $x$  such that

$$\int_0^x \frac{f^{(n)}(t)}{(n-1)!} (x-t)^{n-1} dt = \frac{f^{(n)}(\xi_n)}{(n-1)!} (x-\xi_n)^{n-1} x$$

Now write  $\xi_n = \eta_n x$  for some  $0 < \eta_n < 1$  and  $R_n(x) := \frac{f^{(n)}(\xi_n)}{(n-1)!} (x-\xi_n)^{n-1} x$ . Then

$$R_n(x) = (\alpha-n+1) \binom{\alpha}{n-1} (1+\eta_n x)^{\alpha-n} (x-\eta_n x)^{n-1} x = (\alpha-n+1) \binom{\alpha}{n-1} x^n (1+\eta_n x)^{\alpha-1} \left( \frac{1-\eta_n}{1+\eta_n x} \right)^{n-1}.$$

We need to show that  $R_n(x) \rightarrow 0$  as  $n \rightarrow \infty$ , that is the Taylor series of  $f$  centered at 0 converges to

$f$ . By the Ratio Test, it is easy to see that the series  $\sum_{k=0}^{\infty} (\alpha-k+1) \binom{\alpha}{k} y^k$  is convergent as  $|y| < 1$ .

This tells us that  $\lim_n |(\alpha-n+1) \binom{\alpha}{n} x^n| = 0$ .

On the other hand, note that we always have  $0 < 1 - \eta_n < 1 + \eta_n x$  for all  $n$  because  $x > -1$ . Thus, we



can now conclude that  $R_n(x) \rightarrow 0$  as  $|x| < 1$ . The proof is finished. Finally the last assertion follows from Proposition 7.4 at once. The proof is complete.  $\square$

(Chi-Wai Leung) DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG  
*E-mail address:* cwleung@math.cuhk.edu.hk