

**MATH 2028 Honours Advanced Calculus II**  
**2021-22 Term 1**  
**Problem Set 8**

*due on Nov 15, 2021 (Monday) at 11:59PM*

**Instructions:** You are allowed to discuss with your classmates or seek help from the TAs but you are required to write/type up your own solutions. You can either type up your assignment or scan a copy of your written assignment into ONE PDF file and submit through Gradescope on/before the due date. Please remember to write down your name and student ID. **No late homework will be accepted.**

**Notations:** All surfaces are contained inside  $\mathbb{R}^3$  with rectangular coordinates  $(x, y, z)$ . We use  $U$  to denote a bounded open subset of  $\mathbb{R}^2$ .

**Problems to hand in**

1. Find the area of the portion of the cylinder  $x^2 + y^2 = 2y$  lying inside the sphere  $x^2 + y^2 + z^2 = 4$ .
2. Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$ . Calculate  $\int_S x^2 d\sigma$ . (*Hint: make use of the symmetry*)
3. Find the flux of the vector field  $F(x, y, z) = (x^2, y^2, z^2)$  outward across the given surface  $S$  (all oriented with outward pointing normal pointing away from the origin, unless otherwise specified):
  - (a)  $S$  is the sphere of radius  $a$  centered at the origin.
  - (b)  $S$  is the upper hemisphere of radius  $a$  centered at the origin.
  - (c)  $S$  is the cone  $z = \sqrt{x^2 + y^2}$ ,  $0 < z < 1$ , with outward pointing normal having a negative  $z$ -component.
  - (d)  $S$  is the cylinder  $x^2 + y^2 = a^2$ ,  $0 \leq z \leq h$ .
  - (e)  $S$  is the cylinder  $x^2 + y^2 = a^2$ ,  $0 \leq z \leq h$ , along with the disks  $x^2 + y^2 \leq a^2$ ,  $z = 0$  and  $z = h$ .
4. Calculate the flux of the vector field  $F(x, y, z) = (xz, yz, x^2 + y^2)$  outward across the surface of the paraboloid  $S$  given by  $z = 4 - x^2 - y^2$ ,  $z \geq 0$  (with outward pointing normal having positive  $z$ -component).

**Suggested Exercises**

1. Let  $S$  be the portion of the plane  $x + 2y + 2z = 4$  lying in the first octant of  $\mathbb{R}^3$ , oriented with outward normal pointing upward. Find
  - (a) the area of  $S$ ,
  - (b)  $\int_S (x - y + 3z) d\sigma$ ,
  - (c)  $\int_S F \cdot \vec{n} d\sigma$  where  $F(x, y, z) = (x, y, z)$ .
2. Find the area of the portion of the cylinder  $x^2 + y^2 = a^2$  lying above the  $xy$ -plane and below the plane  $z = y$ .
3. Find the area of the portion of the cone  $z = \sqrt{2(x^2 + y^2)}$  lying beneath the plane  $y + z = 1$ .

4. Let  $S$  be the portion of the helicoid given by the parametrization  $g(u, v) : (0, 1) \times (0, 2\pi) \rightarrow \mathbb{R}^3$  by

$$g(u, v) = (u \cos v, u \sin v, v).$$

Suppose  $S$  is oriented by the upward pointing unit normal  $\vec{n}$ . Compute  $\int_S F \cdot \vec{n} \, d\sigma$  where  $F(x, y, z) = (0, x, 0)$ .

5. Compute  $\int_S F \cdot \vec{n} \, d\sigma$  where  $F(x, y, z) = (x, y, z)$  for each of the following surfaces in  $\mathbb{R}^3$  (all oriented with the outward pointing unit normal pointing away from the origin):

(a) the sphere of radius  $a$  centered at the origin,

(b) the cylinder  $x^2 + y^2 = a^2$ ,  $-h \leq z \leq h$ ,

(c) the cylinder  $x^2 + y^2 = a^2$ ,  $-h \leq z \leq h$ , together with the two disks  $x^2 + y^2 \leq a^2$ ,  $z = \pm h$ ,

(d) the cube with vertices at  $(\pm 1, \pm 1, \pm 1)$ .

6. Repeat the question above for the vector field  $F(x, y, z) = (x^2 + y^2 + z^2)^{-3/2}(x, y, z)$ .

7. Prove that the area of a graphical surface  $S$  given by  $z = f(x, y)$ , where  $f : U \rightarrow \mathbb{R}$  is a  $C^1$  function, is given by

$$\text{Area}(S) = \iint_U \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} \, dA.$$

### Challenging Exercises

1. Prove or give a counterexample: If  $S$  is an orientable surface, then there are exactly two possible orientations on  $S$ .

2. Let  $\alpha, \beta, f : [0, 1] \rightarrow \mathbb{R}$  be  $C^1$  functions with  $f(t) > 0$  for all  $t \in [0, 1]$ . Suppose that  $S$  is a surface in  $\mathbb{R}^3$  whose intersection with the plane  $z = t$  is the circle

$$(x - \alpha(t))^2 + (y - \beta(t))^2 = (f(t))^2, \quad z = t$$

for each  $t \in [0, 1]$  and is empty for  $t \notin [0, 1]$ .

(a) Set up an integral for the area of  $S$ .

(b) Evaluate the integral in (a) when  $\alpha$  and  $\beta$  are constant functions and  $f(t) = (1 + t)^{1/2}$ .

(c) What form does the integral take when  $f$  is constant and  $\alpha(t) = 0$  and  $\beta(t) = at$  where  $a$  is a constant?