


"Uniform" Continuity (§ 5.4 in textbook)

Recall: Let $f: A \rightarrow \mathbb{R}$.

- f is cts at $c \in A$ $\Leftrightarrow \forall \epsilon > 0, \exists \delta = \delta(\epsilon) > 0$ s.t.
 $|f(x) - f(c)| < \epsilon \quad \forall |x - c| < \delta$
- f is cts on A $\stackrel{\text{def}}{\Leftrightarrow}$ f is cts at EVERY $c \in A$
 $\Leftrightarrow \forall c \in A, \forall \epsilon > 0, \exists \delta = \delta(\epsilon, c) > 0,$
s.t. $|f(x) - f(c)| < \epsilon \quad \forall |x - c| < \delta$

Caution: The choice of δ depends on BOTH ϵ AND c .

Example: $f: (0, \infty) \rightarrow \mathbb{R} \quad f(x) := \frac{1}{x}$ cts on $(0, \infty)$



FOR THE SAME $\epsilon > 0$


If $c \approx 0$, then we need to choose a much smaller δ s.t.

$$|f(x) - f(c)| < \epsilon \quad \forall |x - c| < \delta$$

Idea: This function is NOT "uniformly" cts

$\therefore \delta$ is NOT "uniform" in c

Example: $f: (0, \infty) \rightarrow \mathbb{R}$ $f(x) := x$ cts on $(0, \infty)$



FOR THE SAME $\epsilon > 0$

You can choose ONE $\delta > 0$
s.t. it works for ALL $c \in A$

$$|f(x) - f(c)| < \epsilon \quad \forall |x - c| < \delta$$

Idea: This function is "uniformly" cts.

Def: $f: A \rightarrow \mathbb{R}$ is **uniformly continuous** (on A)

iff $\forall \epsilon > 0, \exists \delta = \delta(\epsilon) > 0$ s.t.

$$|f(u) - f(v)| < \epsilon \quad \forall u, v \in A, |u - v| < \delta$$

Remark: (1) uniform cts \Rightarrow cts on A (\because take $v = c \in A$)

(2) Uniform continuity is a "global" concept. It does NOT make sense to talk about uniform continuity at one point $c \in A$.

Q: How to see if $f: A \rightarrow \mathbb{R}$ is uniformly cts (on A)?

We first begin with a "non-uniform continuity" criteria.

Prop: $f: A \rightarrow \mathbb{R}$ is NOT uniformly continuous

$\Leftrightarrow \exists \epsilon_0 > 0$ st $\forall \delta > 0, \exists u_\delta, v_\delta \in A$

st $|u_\delta - v_\delta| < \delta$ BUT $|f(u_\delta) - f(v_\delta)| \geq \epsilon_0$

$\Leftrightarrow \exists \epsilon_0 > 0$ and seq. $(u_n), (v_n)$ in A

st $|u_n - v_n| < \frac{1}{n}$ BUT $|f(u_n) - f(v_n)| \geq \epsilon_0 \quad \forall n \in \mathbb{N}$

Proof: Take negation of defⁿ and choose $\delta = \frac{1}{n}$. _____ ◻

Example: Show that $f: (0, \infty) \rightarrow \mathbb{R}, f(x) = \frac{1}{x}$, is NOT uniformly continuous on $(0, \infty)$.

Proof: Take $(u_n) := (\frac{1}{n})$ and $(v_n) := (\frac{1}{n+1})$ in $(0, \infty)$.

THEN, $|u_n - v_n| = |\frac{1}{n} - \frac{1}{n+1}| = \frac{1}{n(n+1)} < \frac{1}{n} \quad \forall n \in \mathbb{N}$

BUT $|f(u_n) - f(v_n)| = |n - (n+1)| = 1 \geq \epsilon_0 := \frac{1}{2} > 0$.

By Prop, f is NOT uniformly cts on $(0, \infty)$. _____ ◻

Exercise: Show that $f: [a, \infty) \rightarrow \mathbb{R}, f(x) = \frac{1}{x}$ is uniformly cts on $[a, \infty)$ for any fixed $a > 0$.

Idea: We can say more about "uniform continuity" of $f: A \rightarrow \mathbb{R}$ if A is an interval.

{ Uniform Continuity Thm
Continuous Extension Thm

Uniform Continuity Thm

$f: [a, b] \rightarrow \mathbb{R}$ - closed & bdd interval.
cts on $[a, b]$ \Rightarrow f is uniformly cts on $[a, b]$.

Proof: Argue by contradiction. Suppose NOT, i.e. f is NOT uniformly cts. Then, by non-uniform continuity criteria, $\exists \epsilon_0 > 0$ and seq. $(u_n), (v_n)$ in $[a, b]$

(*) \dots [s.t. $|u_n - v_n| < \frac{1}{n}$ BUT $|f(u_n) - f(v_n)| \geq \epsilon_0 \quad \forall n \in \mathbb{N}$]

By Bolzano-Weierstrass Thm, since (u_n) is bdd.

$\Rightarrow \exists$ subseq. (u_{n_k}) of (u_n) s.t.

$$\lim_{k \rightarrow \infty} (u_{n_k}) = x^* \in [a, b]$$

Claim: $\lim_{k \rightarrow \infty} (v_{n_k}) = x^*$

Pf: $|u_{n_k} - v_{n_k}| < \frac{1}{n_k} \quad \Rightarrow \quad \lim_{k \rightarrow \infty} (v_{n_k}) = x^* \quad \text{by limit thm.}$
 $\forall k \in \mathbb{N}$

By continuity of f at $x^* \in [a, b]$.

$$0 < \epsilon_0 \stackrel{(*)}{\leq} \lim_{k \rightarrow \infty} |f(u_{n_k}) - f(v_{n_k})| = |f(x^*) - f(x^*)| = 0$$


Contradiction!

Q: When can we extend a cts $f: (a, b) \rightarrow \mathbb{R}$ to a cts function $\bar{f}: [a, b] \rightarrow \mathbb{R}$?

(s.t. $\bar{f}(x) = f(x) \quad \forall x \in (a, b)$.)


Examples: Yes!

Extend $f(x) = x$ on $x \in (0, 1)$
to $\bar{f}(x) = x$ on $x \in [0, 1]$



No!

$f(x) = \frac{1}{x}$ on $x \in (0, 1)$
 \leadsto \nexists cts extension \bar{f} to $[0, 1]$




Continuous Extension Thm

If $f: (a, b) \rightarrow \mathbb{R}$ is uniformly cts (on (a, b))
then \exists an "extension" $\bar{f}: [a, b] \rightarrow \mathbb{R}$ s.t.

- (i) $\bar{f}(x) = f(x) \quad \forall x \in [a, b]$
- (ii) \bar{f} is cts on $[a, b]$

Remarks: (a) \bar{f} is uniformly cts by Uniform Continuity Thm
(b) Such an extension \bar{f} is unique.

Example:



$y = f(x) = \sin \frac{1}{x}$ remain bdd on $(0, 1)$

BUT not unif. cts on $(0, 1)$
[Ex: Prove this.]

We will use the following lemma in the proof.

Lemma: Let $f: A \rightarrow \mathbb{R}$ be uniformly cts. THEN.

$$\begin{array}{ccc} (x_n) \text{ Cauchy seq} & \Rightarrow & (f(x_n)) \text{ Cauchy seq.} \\ \text{in } A & & \text{in } \mathbb{R} \end{array}$$

Proof of Lemma: Let $\varepsilon > 0$.

By uniform continuity of f , $\exists \delta = \delta(\varepsilon) > 0$ s.t.

$$(\#) \dots \left[|f(u) - f(v)| < \varepsilon \quad \text{whenever } u, v \in A \text{ st } |u - v| < \delta \right]$$

Let (x_n) be a Cauchy seq in A . By ε -H def²,

for this $\delta > 0$ above, $\exists H = H(\delta) \in \mathbb{N}$ st

$$|x_m - x_n| < \delta \quad \forall n, m \geq H$$

$$\text{By } (\#), |f(x_m) - f(x_n)| < \varepsilon \quad \forall n, m \geq H$$

So, $(f(x_n))$ is Cauchy.

_____ \square