

MMAT 5340 Assignment #3

Please submit your assignment online on Blackboard

Due at 12 p.m. on Wednesday, October 6, 2021

- Let $X : \Omega \rightarrow \mathbb{R}$ be a random variable such that $X \equiv 0$, i.e. for any $\omega \in \Omega$, $X(\omega) = 0$. Prove that $\sigma(X) = \{\phi, \Omega\}$.
 - Let $\mathcal{G} := \{\phi, \Omega\}$, and $X : \Omega \rightarrow \mathbb{R}$ be \mathcal{G} -measurable. Prove that $X \equiv c$ for some constant $c \in \mathbb{R}$.
- Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathbb{F} = (\mathcal{F}_n)_{n \geq 0}$ be a filtration. Given a \mathbb{F} -predictable process $(H_n)_{n \geq 0}$, which is uniformly bounded, and a \mathbb{F} -martingale $(X_n)_{n \geq 0}$, we define a process $(V_n)_{n \geq 0}$ by

$$V_0 := 0, \quad V_n := \sum_{k=1}^n H_k (X_k - X_{k-1}).$$

Prove that $(V_n)_{n \geq 0}$ is still a \mathbb{F} -martingale.

- Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathbb{F} = (\mathcal{F}_n)_{n \geq 0}$ be a filtration and $(X_n)_{n \geq 0}$ be a \mathbb{F} -submartingale, we define

$$\Delta A_n := \mathbb{E}[X_n | \mathcal{F}_{n-1}] - X_{n-1}, \quad \Delta M_n := X_n - \mathbb{E}[X_n | \mathcal{F}_{n-1}], \quad \forall n \geq 1,$$

and

$$A_0 = M_0 = 0, \quad A_n := \sum_{k=1}^n \Delta A_k, \quad M_n := \sum_{k=1}^n \Delta M_k.$$

- Prove that $(M_n)_{n \geq 0}$ is a \mathbb{F} -martingale, and that $(A_n)_{n \geq 0}$ is an increasing \mathbb{F} -predictable process.
- Prove that $(X_n)_{n \geq 0}$ has the decomposition

$$X_n = X_0 + M_n + A_n, \quad \forall n \geq 0. \tag{1}$$

- (c) Let $(A_n^1)_{n \geq 0}$ and $(A_n^2)_{n \geq 0}$ be two \mathbb{F} -predictable processes such that $A_0^1 = A_0^2 = 0$. Prove that if $(A_n^1 - A_n^2)_{n \geq 0}$ is a \mathbb{F} -martingale, then $A_n^1 = A_n^2$, a.s. for each $n \geq 1$.
- (d) Deduce that the decomposition (1) is unique, i.e. if one has

$$X_n = X_0 + \widetilde{M}_n + \widetilde{A}_n, \quad \forall n \geq 0.$$

for some \mathbb{F} -martingale $(\widetilde{M}_n)_{n \geq 0}$ and increasing \mathbb{F} -predictable process $(\widetilde{A}_n)_{n \geq 0}$ such that $\widetilde{M}_0 = \widetilde{A}_0 = 0$, then $A_n = \widetilde{A}_n$ and $M_n = \widetilde{M}_n$, a.s. for each $n \geq 1$.