

**MMAT 5340 Assignment #9**  
**Please submit your assignment online on Blackboard**  
**Due at 12 p.m. on Wednesday, 1st Dec, 2021**

1. Suppose we have a box, and  $N$  balls in it. Initially, some of these balls are black and the rests are white. Now we repeatedly apply the following procedure:

- Randomly choose one of the  $N$  balls with equal probability and take it out.
- If the chosen ball is black, we put a white ball into the box.
- If the chosen ball is white, we put a black ball into the box.

Let  $X_n$  be the number of black balls in the box after repeating the above procedure for independently  $n$  times. So we know  $X = (X_n)_{n \geq 0}$  is a Markov chain with state space  $S = \{0, 1, \dots, N\}$  and the transition matrix  $P$ , which is given by

$$P(x, y) = \begin{cases} 1 - \frac{x}{N}, & y = x + 1 \\ \frac{x}{N}, & y = x - 1 \\ 0, & \text{otherwise.} \end{cases} \quad (1)$$

- (a) Prove that the Markov chain  $X$  is irreducible.

By the theorem proved in class, there exists a stationary distribution

$$\mu = (\mu(0), \mu(1), \dots, \mu(N)).$$

- (b) Recall that the stationary distribution  $\mu$  satisfies  $\mu^\top P = \mu^\top$ , we obtain  $N + 1$  linear equations  $\mu(n) = \sum_{k=0}^N \mu(k) P(k, n)$ , for  $n = 0, 1, \dots, N$ .

Please simplify these equations for the transition matrix defined by (1).

(For example, for  $n = 0$ , the linear equation is written as  $\mu(1)/N = \mu(0)$ .)

- (c) Prove that  $\mu(2) = \frac{N(N-1)}{2} \mu(0)$  and  $\mu(x) = \binom{N}{x} \mu(0)$  for all  $x \in S$ .
- (d) Compute the stationary distribution  $\mu$ .

2. Consider the simple random walk  $X = (X_n)_{n \geq 0}$  with state space  $\mathbb{Z}$  (the set of all integers) and transition matrix  $P$ , which is given by

$$P(i, j) = \begin{cases} 1/2, & j = i + 1 \text{ or } j = i - 1 \\ 0, & \text{otherwise.} \end{cases}$$

If  $\pi$  is a stationary distribution of  $X$ , then

- (a) Prove that  $\frac{\pi(x-1) + \pi(x+1)}{2} = \pi(x)$  for all  $x \in \mathbb{Z}$ .
- (b) Let  $u(x) = \pi(x) - \pi(x - 1)$  for  $x \in \mathbb{Z}$  and prove that  $u(x) = C$  for some constant  $C$  for any  $x \in \mathbb{Z}$ .
- (c) Prove that  $\pi(x) = ax + b$  for some constant  $a, b$ .

(d) Prove that  $X$  does not have a stationary distribution.

2

3. Consider a Markov chain  $X = (X_n)_{n \geq 0}$  with state space  $\mathbb{N}$  (the set of all nonnegative integers) and transition matrix  $P$ , which is given by

$$P(j, k) = \begin{cases} 1, & k = j - 1, j \geq 1, \\ 0, & k \neq j - 1, j \geq 1, \\ \nu(k), & k \in \mathbb{N}, j = 0. \end{cases}$$

where  $\nu = \{\nu(n)\}_{n \geq 0}$  is a probability measure on  $\mathbb{N}$ , i.e.  $\nu(n) \geq 0$  for all  $n \in \mathbb{N}$ , and  $\sum_{n=0}^{\infty} \nu(n) = 1$ .

- (a) Prove that  $X$  is irreducible if and only if  $\nu(\{n, n + 1, \dots\}) > 0$  for any  $n \in \mathbb{N}$ , where  $\nu(\{n, n + 1, \dots\}) = \sum_{k=n}^{\infty} \nu(k)$ .
- (b) Prove that 0 is recurrent.
- (c) Prove that the measure defined by  $\mu(n) = \nu(\{n, n + 1, \dots\}), n \in \mathbb{N}$  is stationary, i.e.  $\mu^\top P = \mu^\top$ .