

Suggested Solution to Assignment 4

Exercise 4.1

2. The solution to this problem satisfies the following PDE

$$\begin{aligned} u_t &= ku_{xx}, \quad (0 < x < l, 0 < t < \infty) \\ u(0, t) &= u(l, t) = 0, \\ u(x, 0) &= 1. \end{aligned}$$

Following the process in Page 85 of the textbook, we have

$$u(x, t) = \sum_{n=1}^{\infty} A_n e^{-(\frac{n\pi}{l})^2 kt} \sin \frac{n\pi x}{l},$$

and the initial condition implies

$$1 = \sum_{n=1}^{\infty} A_n \sin \frac{n\pi x}{l}.$$

By the assumption, we have $A_n = \frac{4}{n\pi}$ for odd n and $A_n = 0$ for even ones. Then

$$u(x, t) = \sum_{k=1}^{\infty} \frac{4}{(2k-1)\pi} e^{-(\frac{(2k-1)\pi}{l})^2 kt} \sin \frac{(2k-1)\pi x}{l}. \quad \square$$

4. Let $u(x, t) = T(t)X(x)$, we have

$$\frac{T'' + rT'}{c^2T} = \frac{X''}{X} = -\lambda.$$

Hence,

$$\lambda_n = \left(\frac{n\pi}{l}\right)^2, \quad X(x) = \sin \frac{n\pi x}{l}, \quad n = 1, 2, \dots.$$

Since $0 < r < 2\pi c/l$, we get

$$T_n(t) = [A_n \cos(\sqrt{-\Delta_n}t/2) + B_n \sin(\sqrt{-\Delta_n}t/2)]e^{-rt/2}, \quad n = 1, 2, \dots,$$

where $\Delta_n = r^2 - (2n\pi c/l)^2$ relative to the equation

$$\lambda^2 + r\lambda + \left(\frac{n\pi c}{l}\right)^2 = 0$$

Therefore ,

$$u(x, t) = \sum_{n=1}^{\infty} [A_n \cos(\sqrt{-\Delta_n}t/2) + B_n \sin(\sqrt{-\Delta_n}t/2)]e^{-rt/2} \sin \frac{n\pi x}{l}. \quad \square$$

5. Let $u(x, t) = T(t)X(x)$, we have

$$\frac{T'' + rT'}{c^2T} = \frac{X''}{X} = -\lambda.$$

Hence,

$$\lambda_n = \left(\frac{n\pi}{l}\right)^2, \quad X(x) = \sin \frac{n\pi x}{l}, \quad n = 1, 2, \dots.$$

When $n = 1$, since $2\pi c/l < r < 4\pi c/l$,

$$T_1(t) = A_1 e^{\lambda_1^+ t} + B_1 e^{\lambda_1^- t},$$

where $\lambda_1^\pm = \frac{-r \pm \sqrt{r^2 - (\frac{2\pi c}{l})^2}}{2}$ are the roots of the equation $\lambda^2 + r\lambda + (\frac{\pi c}{l})^2 = 0$.

When $n \geq 2$,

$$T_n(t) = [A_n \cos(\sqrt{-\Delta_n}t/2) + B_n \sin(\sqrt{-\Delta_n}t/2)]e^{-rt/2}, n = 1, 2, \dots,$$

where $\Delta_n = r^2 - (2n\pi c/l)^2$ relative to the equation $\lambda^2 + r\lambda + (\frac{n\pi c}{l})^2 = 0$.

Therefore ,

$$u(x, t) = [A_1 e^{\lambda_1^+ t} + B_1 e^{\lambda_1^- t}] \sin \frac{\pi x}{l} + \sum_{n=2}^{\infty} [A_n \cos(\sqrt{-\Delta_n}t/2) + B_n \sin(\sqrt{-\Delta_n}t/2)] e^{-rt/2} \sin \frac{n\pi x}{l}. \quad \square$$

6. Let $u(x, t) = T(t)X(x)$, we have

$$\frac{tT' - 2T}{T} = \frac{X''}{X} = -\lambda,$$

$$\lambda_n = n^2, X(x) = \sin nx, n = 1, 2, \dots.$$

The initial condition implies

$$tT' - 2T = -\lambda T, T(0) = 0.$$

Therefore,

$$u(x, t) = ct \sin x, \text{ for any constant } c,$$

are solutions. So uniqueness is false for this equation! \square

Exercise 4.2

1. Let $u(x, t) = T(t)X(x)$, we have

$$\frac{T'}{kT} = \frac{X''}{X} = -\lambda.$$

The initial condition implies

$$-X'' = \lambda X, X(0) = X'(l) = 0.$$

So by solving the above DE, the eigenvalues are $[(\frac{n + \frac{1}{2}}{l})\pi]^2$, the eigenfunctions are $X_n(x) = \sin \frac{(n + \frac{1}{2})\pi x}{l}$ for $n = 0, 1, 2, \dots$, and the solution is

$$u(x, t) = \sum_{n=0}^{\infty} e^{-[(\frac{n + \frac{1}{2}}{l})\pi]^2 kt} \sin \frac{(n + \frac{1}{2})\pi x}{l}. \quad \square$$

2. (a) This can be proved as above. Here we give another proof. Since $X'(0) = 0$, the we can use even expansion, this is, $X(-x) = X(x)$ for $-l \leq x \leq 0$, then X satisfies

$$-X'' = \lambda X, X(-l) = X(l) = 0.$$

Hence,

$$\lambda_n = [(n + \frac{1}{2})\pi]^2/l^2, X_n(x) = \cos[(n + \frac{1}{2})\pi x/l], n = 0, 1, 2, \dots.$$

(b) Having known the eigenvalues, it is easy to get the solution

$$u(x, t) = \sum_{n=0}^{\infty} \left[A_n \cos \frac{(n + \frac{1}{2})\pi ct}{l} + B_n \sin \frac{(n + \frac{1}{2})\pi ct}{l} \right] \cos \frac{(n + \frac{1}{2})\pi x}{l}. \quad \square$$

4. We just show how to solve the eigenvalue problem under the periodic boundary conditions; As before, let $u(x, t) = T(t)X(x)$,

$$\frac{T'}{kT} = \frac{X''}{X} = -\lambda$$

Solving $T' = -\lambda kT$ gives $T = Ae^{-\lambda kT}$. The general solutions of $X'' + \lambda X = 0$ are $X = Ce^{\gamma x} + De^{-\gamma x}$, where let λ is a complex number and γ is either one of the two roots of $-\lambda$; the other one is $-\gamma$. The boundary conditions yield

$$Ce^{-\gamma l} + De^{\gamma l} = Ce^{\gamma l} + De^{-\gamma l}, \gamma(Ce^{-\gamma l} - De^{\gamma l}) = \gamma(Ce^{\gamma l} - De^{-\gamma l}).$$

Hence $e^{2\gamma l} = 1$ and then

$$\gamma = \pm n\pi i/l, \lambda = -\gamma^2 = (n\pi/l)^2, n = 0, 1, 2, \dots$$

$$X_n(x) = \begin{cases} \frac{1}{2}A_0 & n = 0 \\ A_n \cos \frac{n\pi x}{l} + B_n \sin \frac{n\pi x}{l}, T = e^{-(n\pi/l)^2 kt} & n = 1, 2, \dots \end{cases}$$

Therefore, the concentration is

$$u(x, t) = \frac{1}{2}A_0 + \sum_{n=0}^{\infty} \left(A_n \cos \frac{n\pi x}{l} + B_n \sin \frac{n\pi x}{l} \right) e^{-(n\pi/l)^2 kt}. \quad \square$$

Exercise 4.3

1. Firstly, let's look for the positive eigenvalues $\lambda = \beta^2 > 0$. As usual, the general solution of the ODE is

$$X(x) = C \cos \beta x + D \sin \beta x.$$

The boundary conditions imply

$$C = 0, D\beta \cos(\beta l) + aD \sin(\beta l) = 0.$$

Hence, $\tan(\beta l) = -\frac{\beta}{a}$. The graph is omitted.

Secondly, let's look for the zero eigenvalue, i.e., $X(x) = Ax + B$, by the boundary conditions, $al + 1 = 0$. Hence, $\lambda = 0$ is an eigenvalue if and only if $al + 1 = 0$.

Thirdly, let's look for the negative eigenvalues $\lambda = -\gamma^2 < 0$. As usual, the solution of the ODE is

$$X(x) = C \cosh(\gamma x) + D \sinh(\gamma x).$$

Then the boundary conditions imply

$$C = 0, D\gamma \cosh(\gamma l) + aD \sinh(\gamma l) = 0.$$

Hence, $\tanh(\gamma l) = -\frac{\gamma}{a}$. The graph is omitted. \square

2. (a) If $\lambda = 0$, then $X(x) = Ax + B$. The boundary conditions imply

$$A - a_0B = 0, A + a_l(A l + B) = 0.$$

These two equalities are equivalent to

$$a_0 + a_l = -a_0 a_l l.$$

Hence, $\lambda = 0$ is an eigenvalue if and only if $a_0 + a_l = -a_0 a_l l$.

(b) By (a), we have $X(x) = B(a_0x + 1)$, here B is constant. \square

3. If $\lambda = -\gamma^2 < 0$, we have

$$X(x) = C \cosh \gamma x + D \sinh \gamma x.$$

Hence,

$$X'(x) = C\gamma \sinh \gamma x + D\gamma \cosh \gamma x,$$

and the boundary conditions imply

$$D\gamma - a_0C = 0,$$

$$C\gamma \sinh \gamma l + D\gamma \cosh \gamma l + a_l[C \cosh \gamma l + D \sinh \gamma l] = 0.$$

Therefore, the eigenvalues satisfy

$$\tanh \gamma l = -\frac{(a_0 + a_l)\gamma}{\gamma^2 + a_0a_l},$$

and the corresponding eigenfunctions are

$$X(x) = C \cosh \gamma x + \frac{a_0}{\gamma}C \sinh \gamma x,$$

where C is a constant. \square

4. It is easily known that the rational curve $y = -\frac{(a_0+a_l)\gamma}{\gamma^2+a_0a_l}$ has a single maximum at $\gamma = \sqrt{a_0a_l}$ and is monotone in the two intervals $(0, \sqrt{a_0a_l})$ and $(\sqrt{a_0a_l}, \infty)$. Furthermore,

$$\max_{\gamma \in [0, \infty)} y(\gamma) = -\frac{a_0 + a_l}{2\sqrt{a_0a_l}} \geq 1, \quad \lim_{\gamma \rightarrow \infty} y(\gamma) = 0, \quad \text{for } y'(0) = -\frac{a_0 + a_l}{a_0a_l}.$$

Note that $\tanh \gamma l$ is monotone in $[0, \infty)$,

$$\tanh \gamma l < 1 \text{ when } \gamma \in [0, \infty), \quad \lim_{\gamma \rightarrow \infty} \tanh \gamma l = 1, \quad \text{and } (\tanh \gamma l)'|_{\gamma=0} = l > -\frac{a_0 + a_l}{a_0a_l}.$$

Therefore, the rational curve $y = -\frac{(a_0+a_l)\gamma}{\gamma^2+a_0a_l}$ and the curve $y = \tanh \gamma l$ intersect at two points, that is, there are two negative eigenvalue. \square

5. When $\lambda = \beta^2 > 0$, β satisfies (10), i.e.

$$\tan \beta l = \frac{(a_0 + a_l)\beta}{\beta^2 - a_0a_l}.$$

Since $y = \tan \beta l$ is monotonically increasing when $\beta \in ((n - \frac{1}{2})\pi/l, (n + \frac{1}{2})\pi/l)$ ($n = 0, 1, 2, \dots$) and

$$\lim_{\beta \rightarrow (n-\frac{1}{2})\pi/l} \tan \beta l = -\infty, \quad \lim_{\beta \rightarrow (n+\frac{1}{2})\pi/l} \tan \beta l = \infty,$$

while $y = \frac{(a_0+a_l)\beta}{\beta^2-a_0a_l}$ is negative, monotonically increasing when $\beta \in (\sqrt{a_0a_l}, \infty)$ and

$$\lim_{\beta \rightarrow \infty} \frac{(a_0 + a_l)\beta}{\beta^2 - a_0a_l} = 0,$$

the two curves intersects at infinite many points, that is, there are an infinite many number of positive eigenvalues. The graph is similiar to the Figure 1 in Section 4.3 in the textbook but $y = \frac{(a_0+a_l)\beta}{\beta^2-a_0a_l}$ is positive first and then negative now. \square




Figure 1:

6. (a) If $a > 0$, the case turns out to be case 1 in Section 4.3 and thus there are no negative eigenvalues; if $a = 0$, the case turns out to be the Neumann boundary condition problem and thus there are no negative eigenvalues, either; if $-2/l \leq a < 0$, we have $(\tanh \gamma l)'|_{\gamma=0} = l \leq -\frac{a_0 + a_l}{a_0 a_l} = -\frac{2}{a}$, using the same way as Exercise 4.3.4 above, we conclude that there is only one negative eigenvalue; if $a < -2/l$, we have $(\tanh \gamma l)'|_{\gamma=0} = l > -\frac{a_0 + a_l}{a_0 a_l} = -\frac{2}{a}$ and thus there are two negative eigenvalues.
- (b) Exercise 4.3.2 implies that $\lambda = 0$ is an eigenvalue if and only if $a_0 + a_l = -a_0 a_l l$, i.e., $a = 0$ or $a = -2/l$. \square
7. Under the condition $a_0 = a_l = a$, the eigenvalue satisfies

$$\lambda = \beta^2, \quad \tan \beta l = \frac{2a\beta}{\beta^2 - a^2}.$$

Hence, when $a \rightarrow \infty$ and $\frac{n\pi}{l} < \beta_n < \frac{(n+1)\pi}{l}$, $\frac{2a\beta}{\beta^2 - a^2}$ is negative and tends to 0. So Figure 1 in Section 4.3 implies

$$\lim_{a \rightarrow \infty} \left\{ \beta_n(a) - \frac{(n+1)\pi}{l} \right\} = 0. \quad \square$$

8. (a)-(c) Please see Figure 1. (d) $|a_0| = |a_l| = \infty$.
9. (a) If $\lambda = 0$, then $X(x) = ax + b$ for some constants a and b . Then the boundary conditions imply $a + b = 0$. Therefore, $X_0(x) = ax - a$ for some nonzero constant a .
- (b) If $\lambda = \beta^2$, then $X(x) = A \cos \beta x + B \sin \beta x$. Then the boundary conditions imply

$$A + B\beta = 0, \quad A \cos \beta + B \sin \beta = 0.$$

Since A, B can not both be 0, we have $\beta = \tan \beta$.

(c) omit.

(d) If $\lambda = -\gamma^2$, then $X(x) = Ae^{\gamma x} + Be^{-\gamma x}$ and

$$A + B + A\gamma - B\gamma = 0, \quad Ae^{\gamma} + Be^{-\gamma} = 0.$$

Then we find the coefficient matrix $\begin{pmatrix} 1 + \gamma & 1 - \gamma \\ e^{2\gamma} & e^{-\gamma} \end{pmatrix}$ is always nonsingular (since $e^{\gamma} > \frac{1+\gamma}{1-\gamma}$ when $\gamma > 0$, verify by yourself!), then $a = b = 0$. So we conclude that there is not any negative eigenvalue. \square

10. Let $u(x, t) = X(x)T(t)$, by the summary on Page 97, we can have

$$u(x, t) = \sum_{n=1}^{\infty} (C_n \cos \beta_n ct + D_n \sin \beta_n ct) \left(\cos \beta_n x + \frac{a_0}{\beta_n} \sin \beta_n x \right) + (C_0 \cosh \gamma ct + D_0 \sinh \gamma ct) \left(\cosh \gamma x + \frac{a_0}{\gamma} \sinh \gamma x \right),$$

where γ is determined by the intersection point of $\tanh \gamma l = -\frac{(a_0 + a_l)\gamma}{\gamma^2 + a_0 a_l}$, and the initial conditions are

$$\begin{aligned} \phi(x) &= \sum_{n=1}^{\infty} C_n \left(\cos \beta_n x + \frac{a_0}{\beta_n} \sin \beta_n x \right) + C_0 \left(\cosh \gamma x + \frac{a_0}{\gamma} \sinh \gamma x \right), \\ \psi(x) &= \sum_{n=1}^{\infty} D_n \beta_n c \left(\cos \beta_n x + \frac{a_0}{\beta_n} \sin \beta_n x \right) + D_0 \gamma c \left(\cosh \gamma x + \frac{a_0}{\gamma} \sinh \gamma x \right). \quad \square \end{aligned}$$

11. (a) By the wave equation,

$$\begin{aligned} \frac{dE}{dt} &= \int_0^l \left[\frac{1}{c^2} u_t u_{tt} + u_x u_{xt} \right] dx \\ &= \int_0^l [u_t u_{xx} + u_x u_{xt}] \\ &= (u_t u_x) \Big|_0^l = u_t(l, t) u_x(l, t) - u_t(0, t) u_x(0, t). \end{aligned}$$

The Dirichlet boundary conditions $u(l, t) = u(0, t) = 0$ imply $u_t(l, t) = u_x(l, t) = 0$. Hence, $\frac{dE}{dt} \equiv 0$.

(b) Same as above. Omit here.

(c) By the computation in (a) and the Robin boundary conditions, we can get that

$$\frac{dE_R}{dt} = u_t u_x \Big|_0^l + a_l u_t(l, t) u(l, t) + a_0 u_t(0, t) u_x(0, t) \equiv 0. \quad \square$$

12. (a) Let $\lambda = 0$, we have $v(x) = Ax + B$. Since $v(x) = Ax + B$ satisfy the boundary conditions for any A and B , $\lambda = 0$ is a double eigenvalue.

(b) Let $\lambda = \beta^2 > 0$ and suppose $\beta > 0$, we have $v(x) = C \cos \beta x + D \sin \beta x$. Then boundary conditions imply

$$D\beta = -C\beta \sin \beta l + D\beta \cos \beta l = \frac{C \cos \beta l + D \sin \beta l - C}{l}.$$

Therefore, eigenvalues $\lambda > 0$ satisfies the equation

$$\lambda = \beta^2, \quad \sin \beta l (-\sin \beta l + \beta l) = (1 - \cos \beta l)^2.$$

(c) Let $\gamma = \frac{1}{2} l \sqrt{\lambda}$, then γ is a root of the following equation

$$\gamma \sin \gamma \cos \gamma = \sin^2 \gamma.$$

(d) By (c), we have $\sin \gamma = 0$ or $\gamma = \tan \gamma$. So the positive eigenvalues are $\frac{4n^2\pi^2}{l^2}$ and $4\gamma_n^2/l^2$ where $\gamma_n = \tan \gamma_n \in (n\pi - \pi, n\pi - \frac{\pi}{2})$ for $n = 1, 2, \dots$. The graph is omitted here.

(e) By (a) and (d), for $\lambda = 0$, the eigenfunctions are 1 and x ; for $\lambda = \frac{4n^2\pi^2}{l^2}$, $n = 1, 2, \dots$, the eigenfunctions are $\cos(\frac{2n\pi x}{l})$; for $\lambda = \frac{4\gamma_n^2}{l^2}$, where $\gamma_n = \tan \gamma_n \in (n\pi - \pi, n\pi - \frac{1}{2}\pi)$, $n = 1, 2, \dots$, the eigenfunctions are

$$\gamma_n \cos \frac{2\gamma_n x}{l} - \sin \frac{2\gamma_n x}{l}.$$

(f) From above, we have

$$u(x, t) = A + Bx + \sum_{n=1}^{\infty} C_n e^{-\frac{4\gamma_n^2}{l^2} kt} \left[\gamma_n \cos \frac{2\gamma_n x}{l} - \sin \frac{2\gamma_n x}{l} \right] + \sum_{n=1}^{\infty} D_n e^{-\frac{4n^2 \pi^2}{l^2} kt} \cos \frac{2n\pi x}{l}.$$

(g) By (f), we have $\lim_{t \rightarrow \infty} u(x, t) = A + Bx$ since $\lim_{t \rightarrow \infty} e^{-\lambda kt} = 0$. \square

13. (a) By Example 2 in Section 1.3, we know that the string should satisfy $u_{tt} = c^2 u_{xx}$. By the arguments in Sections 1.4 and Newton's laws of motion, we obtain the boundary conditions $u(0, t) = 0$ and $u_{tt}(l, t) = -k u_x(l, t)$.

(b) Let $u(x, t) = X(x)T(t)$, then

$$\frac{X''(x)}{X(x)} = \frac{T''(t)}{c^2 T(t)} = -\lambda.$$

$u(0, t) = 0$ implies $X(0) = 0$ and $c^2 u_{xx}(l, t) = u_{tt}(l, t) = -k u_x(l, t)$ implies $c^2 X''(l) = -k X'(l)$. The eigenvalue problem is $X''(x) = -\lambda X(x)$ with the boundary conditions $X(0) = 0$ and $c^2 X''(l) = -k X'(l)$.

(c) Let $\lambda = \beta^2 > 0$. Then $X(x) = A \cos(\beta x) + B \sin(\beta x)$. By boundary conditions, we obtain $A = 0$ and $k \cos(\beta l) = c^2 \beta \sin(\beta l)$. Thus, $\tan(\beta l) = \frac{k}{c^2 \beta}$, which has solutions $n\pi/l < \beta_n < (n+1/2)\pi/l$, $n = 0, 1, 2, \dots$, and the corresponding eigenfunctions are $X_n(x) = \sin(\beta_n x)$.

15. Let $\lambda = \beta^2$, then

$$X(x) = A \cos \frac{\beta \rho_1 x}{\kappa_1} + B \sin \frac{\beta \rho_1 x}{\kappa_1}, \quad 0 < x < a;$$

$$X(x) = C \cos \frac{\beta \rho_2 x}{\kappa_2} + D \sin \frac{\beta \rho_2 x}{\kappa_2}, \quad a < x < l.$$

Hence, the boundary conditions imply

$$A = 0; \quad C \cos \frac{\beta \rho_2 l}{\kappa_2} + D \sin \frac{\beta \rho_2 l}{\kappa_2} = 0;$$

$$A \cos \frac{\beta \rho_1 a}{\kappa_1} + B \sin \frac{\beta \rho_1 a}{\kappa_1} = C \cos \frac{\beta \rho_2 a}{\kappa_2} + D \sin \frac{\beta \rho_2 a}{\kappa_2};$$

$$-A \frac{\beta \rho_1}{\kappa_1} \sin \frac{\beta \rho_1 a}{\kappa_1} + B \frac{\beta \rho_1}{\kappa_1} \cos \frac{\beta \rho_1 a}{\kappa_1} = -C \frac{\beta \rho_2}{\kappa_2} \sin \frac{\beta \rho_2 a}{\kappa_2} + D \frac{\beta \rho_1}{\kappa_1} \cos \frac{\beta \rho_2 a}{\kappa_2}.$$

Hence, when the eigenvalue is positive, i.e. $\lambda = \beta^2 > 0$, β satisfies

$$\frac{\rho_1}{\kappa_1} \cot \frac{\beta \rho_1 a}{\kappa_1} + \frac{\rho_2}{\kappa_2} \cot \frac{\beta \rho_2 (l - a)}{\kappa_2} = 0.$$

Let $\lambda = 0$, then the boundary conditions imply

$$X(x) = \begin{cases} Ax & 0 < x < a; \\ B(x - l) & a < x < l. \end{cases}$$

Since $X(x)$ should be differentiable at $x = a$, such A and B can not exist except $A = B = 0$.

Let $\lambda = -\gamma^2 < 0$, then

$$X(x) = A \cosh \frac{\beta \rho_1 x}{\kappa_1} + B \sinh \frac{\beta \rho_1 x}{\kappa_1}, \quad 0 < x < a;$$

$$X(x) = C \cosh \frac{\beta \rho_2 x}{\kappa_2} + D \sinh \frac{\beta \rho_2 x}{\kappa_2}, \quad a < x < l.$$

Hence, the boundary conditions imply

$$A = 0; \quad C \cosh \frac{\beta \rho_2 l}{\kappa_2} + D \sinh \frac{\beta \rho_2 l}{\kappa_2} = 0;$$

$$A \cosh \frac{\beta \rho_1 a}{\kappa_1} + B \sinh \frac{\beta \rho_1 a}{\kappa_1} = C \cosh \frac{\beta \rho_2 a}{\kappa_2} + D \sinh \frac{\beta \rho_2 a}{\kappa_2};$$

$$A \frac{\beta \rho_1}{\kappa_1} \sinh \frac{\beta \rho_1 a}{\kappa_1} + B \frac{\beta \rho_1}{\kappa_1} \cosh \frac{\beta \rho_1 a}{\kappa_1} = C \frac{\beta \rho_2}{\kappa_2} \sinh \frac{\beta \rho_2 a}{\kappa_2} + D \frac{\beta \rho_1}{\kappa_1} \cosh \frac{\beta \rho_2 a}{\kappa_2}.$$

Hence, when the eigenvalue is negative, i.e. $\lambda = \beta^2 > 0$, β satisfies

$$\frac{\rho_1}{\kappa_1} \coth \frac{\beta \rho_1 a}{\kappa_1} + \frac{\rho_2}{\kappa_2} \coth \frac{\beta \rho_2 (l - a)}{\kappa_2} = 0.$$

However, since the left handside is always positive. Therefore, there is no negative eigenvalues. \square

16. Let $\lambda = \beta^4 > 0$ where $\beta > 0$, and $X(x) = A \cosh \beta x + B \sinh \beta x + C \cos \beta x + D \sin \beta x$. By the boundary conditions

$$\beta_n = \frac{n\pi}{l}, \quad \lambda_n = \left(\frac{n\pi}{l}\right)^4, \quad X_n(x) = \sin \frac{n\pi x}{l}, \quad n = 1, 2, \dots$$

The details are as the following exercise. \square