

## Chapter 3: Continuity

**Learning Objectives:**

- (1) Explore the concept of continuity and examine the continuity of several functions.
- (2) Investigate the intermediate value property.

**3.1 Continuity**

**Definition 3.1.1.** A function  $f$  is **continuous** at  $x = x_0$  if  $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ . It means all three of these conditions are satisfied:

1.  $f(x_0)$  is defined.
2.  $\lim_{x \rightarrow x_0} f(x)$  exists.
3. They are equal.

If some of (1)-(3) are not satisfied, then  $f(x)$  is **discontinuous** at  $x_0$ .

If  $f(x)$  is continuous at every point in the domain,  $f(x)$  is called a **continuous function**.

Informally, a function  $f(x)$  is continuous at  $x = x_0$  if the curve of  $f(x)$  does not break up at  $x_0$ . A continuous function is one whose graph has no holes or gaps.

**Example 3.1.1.** Show that  $f(x) = x^3 - 1$  is continuous at  $x = 1$ .


*Solution.*

$$f(1) = 0.$$

$$\lim_{x \rightarrow 1} f(x) = 1^3 - 1 = 0 = f(1)$$

(i.e., limit exists and is equal to  $f(1)$ .)





**Example 3.1.2.** Decide whether the function

$$f(x) = \begin{cases} x^3 - 1, & x \neq 1, \\ 1, & x = 1. \end{cases}$$

is continuous at  $x = 1$ .

*Solution.* Since

$$\lim_{x \rightarrow 1} f(x) = 0 \neq f(1),$$

$f(x)$  is discontinuous at  $x = 1$ . ■

**Example 3.1.3.** Discuss the continuity of  $f(x) = \frac{1}{x}$ .

*Solution.*  $f(x)$  is defined everywhere except at  $x = 0$ , and  $\lim_{x \rightarrow c} \frac{1}{x} = \frac{1}{c} \forall c \neq 0$  by the first propositions of Chapter 2. So  $f(x)$  is continuous for all  $x \neq 0$ . ■

**Example 3.1.4.** Piecewise linear functions (e.g. step functions, the ceil/floor function,  $f(x) = |x|$ ); piecewise continuous functions.

**Proposition 3.1.1. (Properties of continuity)**

1. Suppose  $f(x)$  and  $g(x)$  are continuous at  $x = x_0$ . It follows from Proposition 2 in Chapter 2 that:
  - (a)  $f(x) + g(x)$ ,  $f(x) - g(x)$ ,  $f(x)g(x)$  are continuous at  $x = x_0$ .
  - (b) If  $g(x_0) \neq 0$ , then  $\frac{f(x)}{g(x)}$  is continuous at  $x = x_0$ .
2. It follows from Proposition 3 in Chapter 2 that: If  $g(x)$  is continuous at  $x = x_0$  and  $f(x)$  is continuous at  $x = g(x_0)$ . Then  $(f \circ g)(x)$ , i.e.,  $f(g(x))$  is continuous at  $x = x_0$ . In fact  $\lim_{x \rightarrow x_0} f(g(x)) = \lim_{u \rightarrow g(x_0)} f(u) = f(g(x_0))$ .
3.  $x^a$ ,  $a^x$ ,  $\log_a x$  and trig functions are all continuous functions in the domain. As a consequence, their  $+$ ,  $-$ ,  $\times$ ,  $\div$ ,  $\circ$  are all continuous in the domain.

**Example 3.1.5.**

1. If  $p(x)$  and  $q(x)$  are polynomials, then

$$\lim_{x \rightarrow c} p(x) = p(c)$$

and

$$\lim_{x \rightarrow c} \frac{p(x)}{q(x)} = \frac{p(c)}{q(c)} \text{ if } q(c) \neq 0.$$

So a polynomial or a rational function is continuous wherever it is defined (i.e.  $q(c) \neq 0$ ).

2.  $f(x) = \frac{x-1}{x+1}$  is continuous at  $x = 2$ .
3.  $f(x) = \frac{x^2-1}{x+1}$  is defined everywhere except at  $x = -1$ , so it is continuous everywhere except at  $x \neq -1$ .
4.  $g(x) = \ln \sqrt{x^2+1}$  is continuous on  $\mathbb{R}$ .

**Example 3.1.6.** Discuss the continuity of the piecewise function:

$$f(x) = \begin{cases} x + 1 & \text{if } x \leq 1, \\ 2x^2 & \text{if } x > 1. \end{cases}$$

*Solution.* For  $x < 1$ ,  $f(x) = x + 1$  is continuous on  $(-\infty, 1)$ ;

For  $x > 1$ ,  $f(x) = 2x^2$  is continuous on  $(1, +\infty)$ ;

At  $x = 1$ ,  $f(1) = 1 + 1 = 2$ .

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x + 1) = 1 + 1 = 2.$$


$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} 2x^2 = 2 \cdot 1^2 = 2.$$

Because the left hand limit and the right hand limit exist and equal. So  $\lim_{x \rightarrow 1} f(x) = 2 = f(1)$ . Therefore  $f(x)$  is continuous at all  $x$ . ■


**Example 3.1.7.**

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

$|x|$  is a continuous everywhere and  $\lim_{x \rightarrow 0} |x| = 0$ .



**Example 3.1.8.** (Discontinuity)



**Example 3.1.9.** For what value of  $A$  such that the following function is continuous at all  $x$ ?

$$f(x) = \begin{cases} x^2 + x - 1 & \text{if } x \leq 0, \\ x + A & \text{if } x > 0. \end{cases}$$

*Solution.* Because  $x^2 + x - 1$  and  $x + A$  are polynomials, they are continuous everywhere except possibly at  $x = 0$ . Also  $f(0) = 0^2 + 0 - 1 = -1$ .

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (x^2 + x - 1) = -1$$

and

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x + A) = A.$$

For  $\lim_{x \rightarrow 0} f(x)$  to exist, the left hand limit and the right hand limit must be equal. So we must have  $A = -1$ . In which case

$$\lim_{x \rightarrow 0} f(x) = -1 = f(0).$$

This means that  $f(x)$  is continuous for all  $x$  only when  $A = -1$ . ■

**Proposition 3.1.2.**  $f(x)$  is continuous at  $x = c$  if and only if

$$\lim_{h \rightarrow 0} f(c + h) = f(c).$$

*Proof.* Let  $h = x - c$ . Then  $h \rightarrow 0$  as  $x \rightarrow c$ .

$$\lim_{x \rightarrow c} f(x) = \lim_{h \rightarrow 0} f(c + h).$$

□

*Exercise 3.1.1.*

1. Show that  $\sqrt[3]{x^3 + 1}$  is a continuous function.
2. Show that  $\left| \frac{x + 1}{x - 1} \right|$  is a continuous function on  $\mathbb{R} \setminus \{1\}$ .
3. Let

$$f(x) = \begin{cases} x^2 - 1, & x \leq 0, \\ x + a, & x > 0. \end{cases}$$

Find  $a$  such that  $f(x)$  is continuous at 0. (Ans:  $a = -1$ )

**Example 3.1.10** (Using continuity to compute limits).  $\lim_{x \rightarrow \infty} \sin\left(\frac{1}{x}\right) = ?$

## 3.2 Continuity on $[a, b]$

**Definition 3.2.1.** Let  $f : (a, b) \rightarrow \mathbb{R}$  be a function. Then  $f$  is said to be continuous on  $(a, b)$  if it is continuous at every point on  $(a, b)$ .

Next, let's assume  $f : [a, b] \rightarrow \mathbb{R}$  be a function. What's the meaning of  $f$  being continuous at one of the end point  $a$ ?  $\lim_{x \rightarrow a} f(x)$  does not make sense because  $f$  is not defined on  $x < a$ . So to define the continuity at  $x = a$ , we only concern about the value  $x > a$ . Similarly, to discuss about the continuity at  $x = b$ , we only concern about the value  $x < b$ .

**Definition 3.2.2.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a function. Then  $f$  is said to be continuous at  $a$  if

$$\lim_{x \rightarrow a^+} f(x) = f(a).$$

$f$  is said to be continuous at  $b$  if

$$\lim_{x \rightarrow b^-} f(x) = f(b).$$

Then  $f$  is said to be a **continuous function on  $[a, b]$**  if  $f$  is continuous on  $a \leq x \leq b$ .


**Example 3.2.1.** Discuss the continuity of the function  $f : [0, 1] \rightarrow \mathbb{R}$  defined by

$$f(x) = \begin{cases} \frac{x-1}{x} & \text{if } x \in (0, 1], \\ 0 & \text{if } x = 0. \end{cases}$$

*Solution.*  $f(x)$  is continuous on  $(0, 1)$ .  $f(x)$  is also continuous at  $x = 1$ , but  $\lim_{x \rightarrow 0^+} f(x)$  does not exist. So  $f$  is not continuous at  $x = 0$ . ■

**Theorem 3.2.1 (Intermediate Value Theorem or Intermediate Value Property).** Suppose  $f$  is a continuous function on  $[a, b]$  and  $K$  is a number between  $f(a)$  and  $f(b)$ . Then there exist a number  $c$ , between  $a$  and  $b$ , such that  $f(c) = K$ .

Geometrically, the Intermediate Value Theorem says that any horizontal line  $y = y_0$  crossing the  $y$ -axis between the numbers  $f(a)$  and  $f(b)$  will cross the curve  $y = f(x)$  at least once over the interval  $[a, b]$ .



### Application: Root Finding

If  $f(x)$  is continuous on  $[a, b]$ ,  $f(a)$  and  $f(b)$  change sign, then, there exists at least one root of the function, that is, exists at least one  $c \in (a, b)$ , such that  $f(c) = 0$ .

**Example 3.2.2.** Show that  $f(x) = x^5 - x + 1$  has a root.

*Solution.* Aim: find  $a, b$ , such that  $f(a), f(b)$  change sign. Since

$$f(-2) = -29, \quad f(0) = 1,$$

and  $f$  is continuous on  $[-2, 0]$ . By Intermediate value theorem, there exists  $c \in (-2, 0)$ , such that  $f(c) = 0$ .

■

*Remark.* Although we don't know how to find the root, we know a root exists.

**Example 3.2.3.** 1. All odd functions have a root.

2. All polynomials of odd degrees have a root.


*Exercise 3.2.1.* Show that  $2^x = \frac{1}{x^2}$  has a solution.

**Theorem 3.2.2** (Extreme Value Theorem). If  $f(x)$  is *continuous on  $[a, b]$* , then  $f$  must attain an *absolute maximum* and *absolute minimum*, that is, there exist  $c, d$  in  $[a, b]$  such that


$$f(c) \leq f(x) \leq f(d),$$

for all  $x \in [a, b]$ .

**Example 3.2.4.** Absolute extreme for  $f(x) = x^3 - 21x^2 + 135x - 170$  for various closed intervals.



(A)  $[a, b] = [2, 12]$



(B)  $[a, b] = [4, 10]$

*Exercise 3.2.2 (Hard!).* Derive the extreme value theorem from the intermediate value theorem.

*Remark.* Caveat: The intermediate value theorem and the extreme value theorem only work on *finite* and *closed* intervals! E.g. Consider the previous example on  $\mathbb{R}$ , and  $\frac{1}{x}$  on  $\mathbb{R}^+$  or on  $(0, 1)$ .

**Question:** How to find the absolute maximum and minimum?

Ans: (for “good” functions) **Differentiation!**