

## 1. Definition. ('Standard base' for a 'vector space of matrices'.)

For each positive integer  $p, q$ , and for each  $i = 1, \dots, p, j = 1, \dots, q$ , we define the  $(p \times q)$ -matrix  $E_{i,j}^{p,q}$  to be the  $(p \times q)$ -matrix whose  $(i, j)$ -th entry is 1 and whose other entries are all 0.

There are altogether  $pq$  matrices  $E_{i,j}^{p,q}$  as  $i, j$  vary.


They are collectively referred to as the 'standard base' for the vector space of  $(p \times q)$ -matrices.

## 2. Examples. ('Standard base' for various 'vector spaces of matrices'.)

$$\begin{aligned} \text{(a) } E_{1,1}^{2,3} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, & E_{1,2}^{2,3} &= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, & E_{1,3}^{2,3} &= \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \\ E_{2,1}^{2,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, & E_{2,2}^{2,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, & E_{2,3}^{2,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

1. **Definition.** ('Standard base' for a 'vector space of matrices'.)

For each positive integer  $p, q$ , and for each  $i = 1, \dots, p, j = 1, \dots, q$ , we define the  $(p \times q)$ -matrix  $E_{i,j}^{p,q}$  to be the  $(p \times q)$ -matrix whose  $(i, j)$ -th entry is 1 and whose other entries are all 0.



There are altogether  $pq$  matrices  $E_{i,j}^{p,q}$  as  $i, j$  vary.

They are collectively referred to as the 'standard base' for the vector space of  $(p \times q)$ -matrices.

2. **Examples.** ('Standard base' for various 'vector spaces of matrices'.)

$$(a) \quad E_{1,1}^{2,3} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad E_{1,2}^{2,3} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad E_{1,3}^{2,3} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix},$$
$$E_{2,1}^{2,3} = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad E_{2,2}^{2,3} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \quad E_{2,3}^{2,3} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$\begin{aligned}
\text{(b) } E_{1,1}^{3,3} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, & E_{1,2}^{3,3} &= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, & E_{1,3}^{3,3} &= \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \\
E_{2,1}^{3,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, & E_{2,2}^{3,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, & E_{2,3}^{2,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \\
E_{3,1}^{3,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, & E_{3,2}^{3,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, & E_{3,3}^{3,3} &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.
\end{aligned}$$

### 3. Lemma (1).

Let  $p, q$  be positive integers. Suppose  $s, t$  are integers between 1 and  $p$ .

Let  $A$  be a  $(p \times q)$ -matrix, whose  $(i, j)$ -th entry is denoted by  $a_{ij}$ .

Then  $E_{s,t}^{p,p} A$  is the  $(p \times q)$ -matrix whose  $s$ -th row is

$$\left[ a_{t1} \ a_{t2} \ \cdots \ a_{tq} \right],$$

and whose every other entry is 0.

**Remark.** In plain words, multiplying  $E_{s,t}^{p,p}$  to  $A$  from the left results in simultaneously ‘putting’ the  $t$ -th row of  $A$  into its  $s$ -th row and setting to ‘zero’ all other rows of  $A$ .

**Proof.** For convenience, denote the  $(g, h)$ -th entry of  $E_{s,t}^{p,p}$  by  $\varepsilon_{gh}$

For each  $k = 1, 2, \dots, q$ , the  $(s, k)$ -th entry of  $E_{s,t}^{p,p} A$  is the product of the  $s$ -th row of  $E_{s,t}^{p,p}$  and the  $k$ -th column of  $A$ , and therefore is given by

$$\varepsilon_{s1}a_{1k} + \varepsilon_{s2}a_{2k} + \cdots + \varepsilon_{sp}a_{pk} = a_{tk}.$$

Hence the  $s$ -th row of  $E_{s,t}^{p,p} A$  is  $\left[ a_{t1} \ a_{t2} \ \cdots \ a_{tq} \right]$ .

Whenever  $g \neq s$ , we have  $\varepsilon_{gh} = 0$  for each  $h$ . Then, no matter which  $k$  is, the  $(g, k)$ -th entry of  $E_{s,t}^{p,p} A$  is a sum of  $p$  copies of 0’s, and hence is 0.

### 3. Lemma (1).


Let  $p, q$  be positive integers. Suppose  $s, t$  are integers between 1 and  $p$ .

Let  $A$  be a  $(p \times q)$ -matrix, whose  $(i, j)$ -th entry is denoted by  $a_{ij}$ .

Then  $E_{s,t}^{p,p} A$  is the  $(p \times q)$ -matrix whose  $s$ -th row is

$$[a_{t1} \ a_{t2} \ \cdots \ a_{tq}],$$

and whose every other entry is 0.



**Remark.** In plain words, multiplying  $E_{s,t}^{p,p}$  to  $A$  from the left results in simultaneously 'putting' the  $t$ -th row of  $A$  into its  $s$ -th row and setting to 'zero' all other rows of  $A$ .

**Proof.** For convenience, denote the  $(g, h)$ -th entry of  $E_{s,t}^{p,p}$  by  $\varepsilon_{gh}$

For each  $k = 1, 2, \dots, q$ , the  $(s, k)$ -th entry of  $E_{s,t}^{p,p} A$  is the product of the  $s$ -th row of  $E_{s,t}^{p,p}$  and the  $k$ -th column of  $A$ , and therefore is given by

$$\varepsilon_{s1}a_{1k} + \varepsilon_{s2}a_{2k} + \cdots + \varepsilon_{sp}a_{pk} = a_{tk}.$$

Hence the  $s$ -th row of  $E_{s,t}^{p,p} A$  is  $[a_{t1} \ a_{t2} \ \cdots \ a_{tq}]$ .

Whenever  $g \neq s$ , we have  $\varepsilon_{gh} = 0$  for each  $h$ . Then, no matter which  $k$  is, the  $(g, k)$ -th entry of  $E_{s,t}^{p,p} A$  is a sum of  $p$  copies of 0's, and hence is 0.

#### 4. Examples. (Illustrations of Lemma (1).)

(a) Suppose  $A$  is the  $(3 \times 4)$ -matrix whose  $(i, j)$ -th entry is given by  $a_{ij}$ . Then:

$$\text{i. } E_{1,2}^{3,3}A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

$$\text{ii. } E_{3,1}^{3,3}A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{11} & a_{12} & a_{13} & a_{14} \end{bmatrix}.$$

$$\text{iii. } E_{3,1}^{3,3}A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix}$$

#### 4. Examples. (Illustrations of Lemma (1).)

(a) Suppose  $A$  is the  $(3 \times 4)$ -matrix whose  $(i, j)$ -th entry is given by  $a_{ij}$ . Then:

$$\begin{array}{l}
 \text{i. } E_{1,2}^{3,3} A = \begin{bmatrix} 0 & \textcircled{1} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \\
 \text{ii. } E_{3,1}^{3,3} A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \textcircled{1} & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{11} & a_{12} & a_{13} & a_{14} \end{bmatrix} \\
 \text{iii. } E_{3,3}^{3,3} A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \textcircled{1} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix}
 \end{array}$$

(b) Suppose  $A$  is the  $(4 \times 6)$ -matrix whose  $(i, j)$ -th entry is given by  $a_{ij}$ . Then:

$$\begin{aligned}
 \text{i. } E_{2,4}^{4,4}A &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} . \\
 \text{ii. } E_{3,2}^{4,4}A &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} . \\
 \text{iii. } E_{4,1}^{4,4}A &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \end{bmatrix} .
 \end{aligned}$$



## 5. Lemma (2).

Let  $A$  be an  $(p, q)$ -matrix. Let  $i, k$  be integers between 1 and  $p$ .

- (a) For any real number  $\alpha$ , the resultant of the row operation  $\alpha R_i + R_k$  on  $A$  is  $(I_p + \alpha E_{k,i}^{p,p})A$ .
- (b) For any non-zero real number  $\beta$ , the resultant of the row operation  $\beta R_k$  on  $A$  is  $(I_p + (\beta - 1)E_{k,k}^{p,p})A$ .
- (c) The resultant of the row operation  $R_i \leftrightarrow R_k$  on  $A$  is  $(I_p - E_{i,i}^{p,p} - E_{k,k}^{p,p} + E_{i,k}^{p,p} + E_{k,i}^{p,p})A$ .

**Proof.** Exercise. (Straightforward calculation with the help of Lemma (1).)

## 6. Examples. (Illustrations of Lemma (2).)

Suppose  $A$  is the  $(3 \times 4)$ -matrix whose  $(i, j)$ -th entry is given by  $a_{ij}$ . Then:

(a)

$$\begin{aligned} A \xrightarrow{4R_2+R_1} & \begin{bmatrix} 4a_{21} + a_{11} & 4a_{22} + a_{12} & 4a_{23} + a_{13} & 4a_{24} + a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &= 4 \begin{bmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = 4E_{1,2}^{3,3}A + A = (I_3 + 4E_{1,2}^{3,3})A \end{aligned}$$

## 5. Lemma (2).

Let  $A$  be an  $(p, q)$ -matrix. Let  $i, k$  be integers between 1 and  $p$ .

- (a) For any real number  $\alpha$ , the resultant of the row operation  $\alpha R_i + R_k$  on  $A$  is  $(I_p + \alpha E_{k,i}^{p,p})A$ .
- (b) For any non-zero real number  $\beta$ , the resultant of the row operation  $\beta R_k$  on  $A$  is  $(I_p + (\beta - 1)E_{k,k}^{p,p})A$ .
- (c) The resultant of the row operation  $R_i \leftrightarrow R_k$  on  $A$  is  $(I_p - E_{i,i}^{p,p} - E_{k,k}^{p,p} + E_{i,k}^{p,p} + E_{k,i}^{p,p})A$ .

**Proof.** Exercise. (Straightforward calculation with the help of Lemma (1).)

## 6. Examples. (Illustrations of Lemma (2).)

Suppose  $A$  is the  $(3 \times 4)$ -matrix whose  $(i, j)$ -th entry is given by  $a_{ij}$ . Then:

(a)

$$\begin{aligned}
 A \xrightarrow{4R_2+R_1} & \begin{bmatrix} 4a_{21} + a_{11} & 4a_{22} + a_{12} & 4a_{23} + a_{13} & 4a_{24} + a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\
 &= 4 \begin{bmatrix} a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = 4E_{1,2}^{3,3}A + \underbrace{A}_{I_3 A} = (I_3 + 4E_{1,2}^{3,3})A
 \end{aligned}$$

$\uparrow$   
 $I_3 A$

This is an example of row-operation matrices.

(b)

$$\begin{aligned} A \xrightarrow{R_1 \leftrightarrow R_3} & \begin{bmatrix} a_{31} & a_{32} & a_{33} & a_{34} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{11} & a_{12} & a_{13} & a_{14} \end{bmatrix} \\ &= \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} - \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &+ \begin{bmatrix} a_{31} & a_{32} & a_{33} & a_{34} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{11} & a_{12} & a_{13} & a_{14} \end{bmatrix} \\ &= A - E_{1,1}^{3,3}A - E_{3,3}^{3,3}A + E_{1,1}^{3,3}A + E_{3,3}^{3,3}A = (I_3 - E_{1,1}^{3,3} - E_{3,3}^{3,3} + E_{1,1}^{3,3} + E_{3,3}^{3,3})A. \end{aligned}$$

(c)

$$\begin{aligned} A \xrightarrow{5R_2} & \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ 5a_{21} & 5a_{22} & 5a_{23} & 5a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &= \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} + 4 \begin{bmatrix} 0 & 0 & 0 & 0 \\ a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 0 & 0 \end{bmatrix} = A + 4E_{2,2}^{3,3}A = (I_3 + 4E_{2,2}^{3,3})A \end{aligned}$$

(b)

$$\begin{aligned} A \xrightarrow{R_1 \leftrightarrow R_3} & \begin{bmatrix} a_{31} & a_{32} & a_{33} & a_{34} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{11} & a_{12} & a_{13} & a_{14} \end{bmatrix} \\ &= \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} - \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &+ \begin{bmatrix} a_{31} & a_{32} & a_{33} & a_{34} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ a_{11} & a_{12} & a_{13} & a_{14} \end{bmatrix} \\ &= \underbrace{A - E_{1,1}^{3,3}A - E_{3,3}^{3,3}A + E_{1,1}^{3,3}A + E_{3,3}^{3,3}A}_{(I_3 - E_{1,1}^{3,3} - E_{3,3}^{3,3} + E_{1,1}^{3,3} + E_{3,3}^{3,3})A} = (I_3 - E_{1,1}^{3,3} - E_{3,3}^{3,3} + E_{1,1}^{3,3} + E_{3,3}^{3,3})A. \end{aligned}$$

This is another example of row-operation matrices.

(c)

$$\begin{aligned} A \xrightarrow{5R_2} & \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ 5a_{21} & 5a_{22} & 5a_{23} & 5a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} \\ &= \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} + 4 \begin{bmatrix} 0 & 0 & 0 & 0 \\ a_{21} & a_{22} & a_{23} & a_{24} \\ 0 & 0 & 0 & 0 \end{bmatrix} = \underbrace{A + 4E_{2,2}^{3,3}A}_{(I_3 + 4E_{2,2}^{3,3})A} = (I_3 + 4E_{2,2}^{3,3})A \end{aligned}$$

This is yet another example of row-operation matrices.

## 7. Definition. (Row operation matrices.)

Let  $p$  be a positive integer, and  $M$  be a  $(p, p)$ -square matrix.

The matrix  $M$  is called a row-operation matrix of size  $p$  if any one of the statements below holds:

- (a)  $M = I_p + \alpha E_{k,i}^{p,p}$  for some real number  $\alpha$  and some distinct integers  $i, k$  between 1 and  $p$ .
- (b)  $M = I_p + (\beta - 1)E_{k,k}^{p,p}$  for some non-zero real number  $\beta$  and some integer  $k$  between 1 and  $p$ .
- (c)  $M = I_p - E_{i,i}^{p,p} - E_{k,k}^{p,p} + E_{i,k}^{p,p} + E_{k,i}^{p,p}$  for some distinct integers  $i, k$ .

**Remark.** Now we know that the effect of applying a certain row operation on a matrix, say,  $A$ , is the same as multiplying  $A$  by some row operation matrix from the left.

In fact such a square matrix is uniquely determined by the row operation concerned; it is independent of  $A$ .

Theorem (3) below describes a ‘dictionary’ between the collection of all row operations on matrices with  $p$  rows and the collection of all row-operation matrices of size  $p$ . This ‘dictionary’ tells us the ‘application of row operations’ and the ‘multiplication from the left by row-operation matrices’ are two ways of thinking about the same thing.

## 8. Theorem (3). ('Dictionary' between row operations and matrix multiplication from the left.)

Let  $p, q$  be positive integers.

For any row operation  $\rho$  on  $(p \times q)$ -matrices, there exists some unique  $(p \times p)$ -square matrix  $M[\rho]$  such that for any  $(p \times q)$ -matrix  $A$ , the matrix  $M[\rho]A$  is the resultant of the application of  $\rho$  on  $A$ .

**Proof.** A tedious word game, with reference to the definitions for the notion of row operations and for the notion of row-operation matrices.

**Remark.** The table below summarizes the correspondence between row operations and row-operation matrices:

Row operation changing $C$ to $C'$ .	How $C'$ is obtained from $C$ through row-operation matrix.	'Reverse' row operation changing $C'$ to $C$ .	How $C$ is recovered from $C'$ through row-operation matrix.
$C \xrightarrow{\alpha R_i + R_k} C'$ .	$C' = (I_p + \alpha E_{k,i}^{p,p})C$	$C' \xrightarrow{-\alpha R_i + R_k} C$ .	$C = (I_p - \alpha E_{k,i}^{p,p})C'$
$C \xrightarrow{\beta R_k} C'$ .	$C' = [I_p + (\beta - 1)E_{k,k}^{p,p}]C$	$C' \xrightarrow{(1/\beta)R_k} C$ .	$C = [I_p + (1/\beta - 1)E_{k,k}^{p,p}]C'$
$C \xrightarrow{R_i \leftrightarrow R_k} C'$ .	$C' = (I_p - E_{i,i}^{p,p} - E_{k,k}^{p,p} + E_{i,k}^{p,p} + E_{k,i}^{p,p})C$	$C' \xrightarrow{R_i \leftrightarrow R_k} C$ .	$C = (I_p - E_{i,i}^{p,p} - E_{k,k}^{p,p} + E_{i,k}^{p,p} + E_{k,i}^{p,p})C'$

## 9. Corollary (4).

Let  $C_1, C_2, \dots, C_N$  be  $(p \times q)$ -matrices.

Suppose  $C_1$  is row-equivalent to  $C_N$ , and are joint by some sequence of row operations  $\rho_1, \rho_2, \dots, \rho_{N-1}$ :

$$C_1 \xrightarrow{\rho_1} C_2 \xrightarrow{\rho_2} \cdots \xrightarrow{\rho_{N-2}} C_{N-1} \xrightarrow{\rho_{N-1}} C_N$$

Then there exist row-operation matrices  $H_1, H_2, \dots, H_{N-1}$  of size  $p$  such that  $C_N = H_{N-1}H_{N-2} \cdots H_2H_1C_1$ .

**Proof.** This is an immediate consequence of Theorem (3).

### 9. Corollary (4).

Let  $C_1, C_2, \dots, C_N$  be  $(p \times q)$ -matrices.

Suppose  $C_1$  is row-equivalent to  $C_N$ , and are joint by some sequence of row operations  $\rho_1, \rho_2, \dots, \rho_{N-1}$ :

$$C_1 \xrightarrow{\rho_1} C_2 \xrightarrow{\rho_2} \dots \xrightarrow{\rho_{N-2}} C_{N-1} \xrightarrow{\rho_{N-1}} C_N$$

Then there exist row-operation matrices  $H_1, H_2, \dots, H_{N-1}$  of size  $p$  such that  $C_N = H_{N-1}H_{N-2} \dots H_2H_1C_1$ .

**Proof.** This is an immediate consequence of Theorem (3).

$$\begin{cases} C_2 = H_1 C_1 \\ C_3 = H_2 C_2 \\ \vdots \\ C_N = H_{N-1} C_{N-1} \end{cases}$$

What  $H_j$ 's?  
 For each  $j$ , the row operation  $\rho_j$  has a corresponding row operation matrix; this is  $H_j$ .

$$\begin{aligned} C_N &= H_{N-1} C_{N-1} = H_{N-1} (H_{N-2} C_{N-2}) = H_{N-1} H_{N-2} C_{N-2} \\ &= H_{N-1} H_{N-2} H_{N-3} \dots H_2 H_1 C_1 \end{aligned}$$

How to obtain  $H_{N-1} H_{N-2} \dots H_2 H_1$  explicitly? Through an appropriate sequence of row operations, starting from  $I_p$ .

$$I_p \xrightarrow{\rho_1} H_1 \xrightarrow{\rho_2} H_2 H_1 \xrightarrow{\rho_3} H_3 H_2 H_1 \rightarrow \dots \xrightarrow{\rho_{N-1}} H_{N-1} H_{N-2} \dots H_2 H_1$$



## 10. Examples. (Illustrations on Corollary (4).)

(a) The sequence of row operations below joins  $C$  and  $C''$ :

$$C = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 1 & 0 & 0 & 2 \end{bmatrix} \xrightarrow{1R_1+R_2} C' = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 0 & 0 & 2 \end{bmatrix} \xrightarrow{2R_2+R_1} C'' = \begin{bmatrix} 3 & 4 & 5 & 5 \\ 1 & 2 & 2 & 2 \\ 1 & 0 & 0 & 2 \end{bmatrix}.$$

$$\text{Then } C'' = H_2 H_1 C, \text{ in which } H_1 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, H_2 = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$\text{So } C'' = HC, \text{ in which } H = H_2 H_1 = \begin{bmatrix} 3 & 2 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

(b) The sequence of row operations below joins  $C$  and  $C''$ :

$$C = \begin{bmatrix} 1 & 2 & 2 & -1 \\ 2 & -2 & 1 & 0 \\ 1 & 0 & 0 & 2 \end{bmatrix} \xrightarrow{4R_2} C' = \begin{bmatrix} 1 & 2 & 2 & -1 \\ 8 & -8 & 4 & 0 \\ 1 & 0 & 0 & 2 \end{bmatrix} \xrightarrow{-2R_1} C'' = \begin{bmatrix} -2 & -4 & -4 & 2 \\ 8 & -8 & 4 & 0 \\ 1 & 0 & 0 & 2 \end{bmatrix}.$$

$$\text{Then } C'' = H_2 H_1 C, \text{ in which } H_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}, H_2 = \begin{bmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

$$\text{So } C'' = HC, \text{ in which } H = H_2 H_1 = \begin{bmatrix} -2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

(c) The sequence of row operations below joins  $C$  and  $C''$ :

$$C = \begin{bmatrix} 1 & 2 & 2 & 0 \\ 3 & 0 & 3 & 1 \\ 2 & 1 & 0 & 1 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_2} C' = \begin{bmatrix} 3 & 0 & 3 & 1 \\ 1 & 2 & 2 & 0 \\ 2 & 1 & 0 & 1 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_3} C'' = \begin{bmatrix} 3 & 0 & 3 & 1 \\ 2 & 1 & 0 & 1 \\ 1 & 2 & 2 & 0 \end{bmatrix}.$$

$$\text{Then } C'' = H_2 H_1 C, \text{ in which } H_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, H_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

$$\text{So } C'' = HC, \text{ in which } H = H_2 H_1 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}.$$

(d) The sequence of row operations below joins  $C$  and  $C'''$ :

$$C = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 1 & 0 & 0 & 2 \end{bmatrix} \xrightarrow{1R_1+R_2} C' = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 0 & 0 & 2 \end{bmatrix} \xrightarrow{2R_3} C'' = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 2 & 0 & 0 & 4 \end{bmatrix}$$

$$\xrightarrow{R_1 \leftrightarrow R_3} C''' = \begin{bmatrix} 2 & 0 & 0 & 4 \\ 1 & 2 & 2 & 2 \\ 1 & 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Then } C''' = H_3 H_2 H_1 C, \text{ in which } H_1 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, H_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}, H_3 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

$$\text{So } C''' = HC, \text{ in which } H = H_3 H_2 H_1 = \begin{bmatrix} 0 & 0 & 2 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$