Assignment 2

Deadline: Jan 24, 2018.

Hand in: Section 6.2 no 9, 10, 18; Supplementary exercise no 2, 3a.

Section 6.2: Q.5, 9-12, 17, 18, 19.

Supplementary Exercises

- 1. Let f be a function defined on \mathbb{R} . It is called a periodic function if there is a non-zero number T such that f(x+T) = f(x) for all x. The number T is called a period of f.
 - (a) Show that $nT, n \neq 0 \in \mathbb{Z}$, is also a period if f has a period T.
 - (b) Let f be differentiable. Show that f must be constant if it has a sequence of periods $\{T_n\}, T_n \to 0$. Hint: If f is non-constant, $f'(c) \neq 0$ at some c.
 - (c) (Optional) Let f be differentiable. Show that if f is non-constant, there exists a positive period L satisfying, if T is another period of f, then T = nL for some integer n. This L is called the minimal period of f.
- 2. Let f be a differentiable function defined on $(0, \infty)$. Suppose f satisfies $|f(x)| \leq C\sqrt{x}$ for all $x \in (0, \infty)$ for some constant C > 0. Show that there exists a sequence of numbers $\{x_n\}, x_n \to \infty$, such that $f'(x_n) \to 0$ as $n \to \infty$.
- 3. (a) Let $p : \mathbb{R} \to \mathbb{R}$ be a polynomial $p(x) = a_0 + a_1 x + \dots + a_n x^n$, where $n \in \mathbb{N}$, $a_0, a_1, \dots, a_n \in \mathbb{R}$ and $a_n \neq 0$. Suppose that p has n real roots. Show that p' has n-1 real roots.
 - (b) (Optional) What happens when p does not have n real roots? In this case, there are complex roots. Could you make a guess on the distribution of the roots of p'?
- 4. It has been shown that a differentiable function f on (a, b) satisfying f'(x) = 0 everywhere must be a constant. Show that this result is not true when the assumption is relaxed to the right derivative of f exists and $f'_{+}(x) = 0$ everywhere.