


Remark : The distance formula

⇒ Postulates 2 & 3 of Euclidean geometry also hold in hyperbolic geometry.

Postulate 2 : A line can be produced indefinitely in either direction.



Since $\lim_{r \rightarrow 1} d(0, r)$

$$= \lim_{r \rightarrow 1} \ln \frac{1+r}{1-r} = +\infty$$

$\forall N > 0, \exists 1 > r_1 > r$ st.

$$\ln \frac{1+r_1}{1-r_1} > \ln \frac{1+r}{1-r} + N$$


ie. $d(0, r_1) > d(0, r) + N$

∴ hyperbolic straight line segment
can be produced indefinitely

(ie. longer than any prescribed length)

Postulate 3: A circle can be described with any center and radius.

Pf: Use a transformation, we only need to consider



Note that

$$d(0, re^{i\theta}) = d(0, r) = \ln \frac{1+r}{1-r}$$

\therefore hyperbolic distance from a point $z = re^{i\theta}$ to the center O is a constant

$$d(0, re^{i\theta}) = \ln \frac{1+r}{1-r}$$

depending only on $r \in (0, 1)$.

Hence $R = d(0, re^{i\theta}) = \ln \frac{1+r}{1-r}$

is the (hyperbolic) radius of the hyperbolic circle

And for any $R > 0$, we can solve $R = \ln \frac{1+r}{1-r}$.

to find $r = \frac{e^R - 1}{e^R + 1}$ (check), $r \in (0, 1)$.

Then the Euclidean circle centered at 0 with Euclidean radius $r = \frac{e^R - 1}{e^R + 1}$ is the required hyperbolic circle centered at 0 with hyperbolic radius R . ~~XX~~

Conclusion = The hyperbolic geometry is a non-Euclidean geometry in the strict sense.


(Postulate 4 is automatically satisfied as all transformations in the hyperbolic geometry are conformal. Euclidean angle measurement is invariant and hence provides the required angle in hyperbolic geometry.)

Formula of Lobatchevsky

Thm: Let the point p be given by the hyperbolic distance d from a hyperbolic straight line. Let θ be the angle of parallelism of p with respect to this line. Then

$$\boxed{e^{-d} = \tan \frac{\theta}{2}}$$

Pf:



After a transformation, we may assume $p=0$ and the perpendicular from p to the hyperbolic straight line is the x-axis.

Then the point r as in the figure is given

by
$$d = \ln \frac{1+r}{1-r}$$

and
$$r = \sqrt{1 + \tan^2 \theta} - \tan \theta$$

$$= \frac{1}{\cos \theta} - \tan \theta$$

$$= \frac{1 - \sin \theta}{\cos \theta}$$

$$\Rightarrow e^{-d} = \frac{1-r}{1+r} = \frac{1 - \frac{1 - \sin \theta}{\cos \theta}}{1 + \frac{1 - \sin \theta}{\cos \theta}} = \frac{\cos \theta + \sin \theta - 1}{\cos \theta - \sin \theta + 1}$$

$$= \frac{(\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}) + 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} - 1}{(\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}) - 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} + 1}$$

$$= \frac{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} - 2 \sin^2 \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2} - 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} = \tan \frac{\theta}{2}$$

~~✗~~

The Upper Half Plane Model

Def: The Upper half plane is the subset

$$\mathbb{U} = \{z : \text{Im}z > 0\} \subset \mathbb{C}$$

Let $\overline{\mathbb{H}}$ be the group of transformations (of \mathbb{U}) of the form

$$\left\{ w = Tz = \frac{az+b}{cz+d}, \quad \begin{array}{l} a, b, c, d \in \mathbb{R} \\ \& ad-bc > 0 \end{array} \right\}$$

The pair $(\mathbb{U}, \overline{\mathbb{H}})$ models hyperbolic geometry.


Remark: Both $(\mathbb{D}, \overline{\mathbb{H}})$ and $(\mathbb{U}, \overline{\mathbb{H}})$ are models of the same abstract geometry, namely the hyperbolic geometry. (see (Ex.) in the proof below)

Distance in the upper half plane model


$$l(\gamma) = \int_a^b \frac{|z'(t)|}{y(t)} dt \quad \text{for } \gamma : z(t) = x(t) + iy(t)$$

Pf: Consider the transformation (Möbius, but not in \mathbb{H} , the group of hyperbolic geometry)

$$w = S(z) = i \frac{1+z}{1-z}$$



S



$$S(-1) = 0, \quad S(0) = i, \quad S(1) = \infty$$


Hence • $S: \mathbb{D} \rightarrow \mathbb{U}$

$$\bullet S^{-1}w = \frac{i w + 1}{i w - 1} \quad (\text{check!})$$

$$\bullet Tz = e^{i\theta} \frac{z - z_0}{1 - \bar{z}_0 z}, \quad z_0 \in \mathbb{D}, \theta \in \mathbb{R} \quad (\text{i.e. } T \in \mathbb{H})$$

(Ex!)

$\Leftrightarrow S \circ T \circ S^{-1}$ is a transformation in $\overline{\mathbb{H}}$



$$\begin{array}{c} T \in \mathbb{H} \\ \Downarrow \\ S \circ T \circ S^{-1} \in \overline{\mathbb{H}} \end{array}$$

$\therefore S$ is an isomorphism of the disk and upper half-plane models.

Now let $\gamma = z(t) = x(t) + iy(t)$, $a \leq t \leq b$ be a smooth curve in the upper half plane (i.e. $y(t) > 0$)


$$\text{Then } \hat{\gamma} = \hat{z}(t) = S^{-1}(z(t)) = \frac{iz(t)+1}{iz(t)-1} \quad \text{is a smooth curve in } \mathbb{D}.$$

$$\Rightarrow |\hat{z}'(t)| = \frac{|(iz(t)-1)(iz'(t)) - (iz(t)+1)iz'(t)|}{|iz(t)-1|^2} = \frac{2|z'(t)|}{|iz(t)-1|^2}$$

$$\Rightarrow \underset{\substack{\uparrow \\ \text{upper half plane} \\ \text{model}}}{l(\gamma)} = \underset{\substack{\uparrow \\ \text{disk model}}}{l(\hat{\gamma})} = 2 \int_a^b \frac{|\hat{z}'|}{1-|\hat{z}|^2} dt = 2 \int_a^b \frac{\frac{2|z'|}{|iz-1|^2}}{1 - \left| \frac{iz+1}{iz-1} \right|^2} dt$$


$$\begin{aligned}
&= \int_a^b \frac{4|z'|}{|\bar{i}z-1|^2 - |\bar{i}z+1|^2} dt \\
&= \int_a^b \frac{4|z'|}{|\bar{i}x-y-1|^2 - |\bar{i}x-y+1|^2} dt \\
&= \int_a^b \frac{4|z'|}{[(1+y)^2 + x^2] - [(1-y)^2 + x^2]} dt \\
&= \int_a^b \frac{|z'|}{y} dt \quad \#
\end{aligned}$$

Remark: Hyperbolic straight lines in the upper half plane model are




Ch10 Area (hyperbolic)

Recall: In Euclidean geometry



$$\text{Area}(R) \sim \sum \Delta x \Delta y \longrightarrow \iint_R dx dy$$

Similarly, in hyperbolic geometry (upper half plane model)



$$\left[\begin{array}{l} y + \Delta y \\ y \\ z(t) = x + it, y \leq t \leq y + \Delta y \\ l = \int_y^{y + \Delta y} \frac{|dz|}{x} dt = \ln \frac{y + \Delta y}{y} \sim \frac{\Delta y}{y} \end{array} \right]$$

Hence, we make


Def: The (hyperbolic) area of a region R
in the hyperbolic upper half plane model
is given by

$$A = \iint_R \frac{dx dy}{y^2}$$


Areas of Triangles

(1) Doubly asymptotic triangles


(i.e. triangles with 2 ideal vertexes)



disk model



upper half-plane model



We only need to consider the case that the ideal points at ∞ and -1 , and the "finite" vertex somewhere along the unit circle


(Ex: Hurts = horizontal translations and scaling are transformations of (\mathbb{U}, \mathbb{H}))

Let α = interior angle of the triangle at the "finite" vertex. Then Euclidean geometry \Rightarrow the "finite" vertex has coordinates $(\cos \alpha, \sin \alpha)$

Hence the area of the triangle is

$$A = \iint \frac{dx dy}{y^2}$$

$$= \int_{-1}^{\cos \alpha} \left(\int_{\sqrt{1-x^2}}^{+\infty} \frac{dy}{y^2} \right) dx$$




$$= \int_{-1}^{\cos \alpha} \frac{1}{\sqrt{1-x^2}} dx \quad \left(\begin{array}{l} \text{let } x = \cos \theta, \theta \in [\alpha, \pi] \\ \Rightarrow \sin \theta \geq 0 \\ \Delta x' \leq 0 \end{array} \right)$$


$$= \pi - \alpha$$

ie. $\boxed{A = \pi - \alpha}$

(2) Treblely asymptotic triangle (ideal triangle)
 (ie. all 3 vertexes are ideal points.)



($\alpha = 0$)



By (1), we have


$$\boxed{A = \pi}$$

(for any treblely asymptotic triangle.)

(3) General triangle

We may put the triangle in a way such

that one of the edge is long the y-axis



$$I = \Delta PP^*g^*, \quad II = \Delta gq^*r^*, \quad III = \Delta rr^*p^*$$

(as in the figure) are doubly asymptotic triangles.

Then by

$$\Delta p^*q^*r^* = \Delta pqr \cup I \cup II \cup III$$

(interior disjoint union)

is a trebly asymptotic triangle

we have by (1) & (2)

$$\begin{aligned}\pi &= A + [\pi - (\pi - \alpha)] + [\pi - (\pi - \beta)] + [\pi - (\pi - \gamma)] \\ &= A + (\alpha + \beta + \gamma)\end{aligned}$$

$$\Rightarrow \boxed{A = \pi - (\alpha + \beta + \gamma)}$$

i.e. The area of a triangle equals to π minus the sum of interior angles which is called angular defect.

Thm: The area of a triangle (in hyperbolic geometry) equals its angular defect.

Thm: The sum of the interior angles of a triangle in hyperbolic geometry is less than π radians.