

SCATTERING DIAGRAMS FROM ASYMPTOTIC ANALYSIS ON MAURER-CARTAN EQUATIONS

KWOKWAI CHAN, NAICHUNG CONAN LEUNG, AND ZIMING NIKOLAS MA

ABSTRACT. Let \check{X}_0 be a semi-flat Calabi-Yau manifold equipped with a Lagrangian torus fibration $\check{p} : \check{X}_0 \rightarrow B_0$. We investigate the asymptotic behavior of Maurer-Cartan solutions of the Kodaira-Spencer deformation theory on \check{X}_0 by expanding them into Fourier series along fibres of \check{p} over a contractible open subset $U \subset B_0$, following a program set forth by Fukaya [21] in 2005. We prove that semi-classical limits (i.e. leading order terms in asymptotic expansions) of the Fourier modes of a specific class of Maurer-Cartan solutions naturally give rise to consistent scattering diagrams, which are tropical combinatorial objects that have played a crucial role in works of Kontsevich-Soibelman [37] and Gross-Siebert [28] on the reconstruction problem in mirror symmetry.

1. INTRODUCTION

1.1. Background. The celebrated Strominger-Yau-Zaslow (SYZ) conjecture [45] asserts that mirror symmetry is a *T-duality*, meaning that a mirror pair of Calabi-Yau manifolds should admit fibre-wise dual (special) Lagrangian torus fibrations to the same base. This immediately suggests a construction of the mirror (as a complex manifold): Given a Calabi-Yau manifold X , one first looks for a Lagrangian torus fibration $p : X \rightarrow \check{B}$. The base \check{B} is then an integral affine manifold with singularities. Letting $\check{B}_0 \subset \check{B}$ be the smooth locus and setting

$$\check{X}_0 := T\check{B}_0/\Lambda_{\check{B}_0},$$

where $\Lambda_{\check{B}_0} \subset T\check{B}_0$ denotes the natural lattice locally generated by affine coordinate vector fields, yields a torus bundle $\check{p} : \check{X}_0 \rightarrow \check{B}_0$ which admits a natural complex structure \check{J}_0 , called the *semi-flat complex structure*. This would not produce the correct mirror in general,¹ simply because \check{J}_0 cannot be extended across the singular points \check{B}^{sing} . But the SYZ proposal suggests that the mirror is given by deforming \check{J}_0 using *quantum corrections* coming from holomorphic disks in X with boundary on the Lagrangian torus fibres of p .

The precise mechanism of such a mirror construction was first depicted by Kontsevich-Soibelman [36] using rigid analytic geometry and then by Fukaya [21] using asymptotic analysis. In Fukaya's proposal, he described how instanton corrections would arise near the large volume limit given by scaling of the symplectic structure on X by $\hbar \in \mathbb{R}_{>0}$, which is mirrored to scaling of the complex structure \check{J}_0 on \check{X}_0 . It was conjectured that the desired deformations of \check{J}_0 were given by a specific class of solutions to the Maurer-Cartan equation of the Kodaira-Spencer deformation theory of complex structures on \check{X}_0 , whose expansions into Fourier modes along torus fibres of \check{p} would have semi-classical limits (i.e. leading order terms in asymptotic expansions as $\hbar \rightarrow 0$) concentrated along gradient flow trees of a canonically defined multi-valued Morse function on \check{B}_0 [21, Conjecture 5.3]. On the mirror side, holomorphic disks in X with boundary on fibres of p were conjectured to collapse to gradient flow trees emanating from the singular points $\check{B}^{\text{sing}} \subset \check{B}$ [21, Conjecture 3.2]. From this one sees directly how the mirror complex structure is determined by quantum corrections. Unfortunately, the arguments in [21] were only heuristical and the analysis involved to make them precise seemed intractable at that time.

¹Except in the semi-flat case when $\check{B} = \check{B}_0$ where there are no singular fibres; see [40].

These ideas were later exploited by Kontsevich-Soibelman [37] (for dimension 2) and Gross-Siebert [28] (for general dimensions) to construct families of rigid analytic spaces and formal schemes respectively from integral affine manifolds with singularities, thereby solving the very important *reconstruction problem* in SYZ mirror symmetry. They cleverly got around the analytical difficulties, and instead of solving the Maurer-Cartan equation, used gradient flow trees in \check{B}_0 [37] or tropical trees in the Legendre dual B_0 [28] to encode the modified gluing maps between charts in constructing the mirror family. A key notion in their constructions is that of *scattering diagrams*, which are combinatorial structures encoding possibly very complicated gluing data. It has also been understood (by works of these authors and their collaborators, notably [27]) that these scattering diagrams encode Gromov-Witten data as well.

In this paper, we revisit Fukaya’s original ideas and apply asymptotic analysis motivated by Witten-Morse theory [46]. Our primary goal is to connect consistent scattering diagrams to the asymptotic behavior of a specific class of solutions of the Maurer-Cartan equation. In particular we prove a modified version of (the “scattering part” of) Fukaya’s original conjecture in [21]. As pointed out by Fukaya himself, understanding scattering phenomenon is vital to a general understanding of quantum corrections in mirror symmetry.

We start with a Calabi-Yau manifold X (regarded as a symplectic manifold) equipped with a Lagrangian torus fibration which admits a Lagrangian section \mathfrak{s}

$$(X, \omega, J) \begin{array}{c} \xleftarrow{\mathfrak{s}} \\ \xrightarrow{p} \end{array} \check{B}$$

and whose discriminant locus is given by $\check{B}^{\text{sing}} \subset \check{B}$, over which the integral affine structure develops singularities. Restricting p to the smooth locus $\check{B}_0 = \check{B} \setminus \check{B}^{\text{sing}}$, we obtain a semi-flat symplectic Calabi-Yau manifold $X_0 \hookrightarrow X$, which, by Duistermaat’s action-angle coordinates [16], can be identified as a quotient of the cotangent bundle of the base $X_0 \cong T^*\check{B}_0/\Lambda_{\check{B}_0}^\vee$, where $\Lambda_{\check{B}_0}^\vee \subset T^*\check{B}_0$ is the natural lattice (dual to $\Lambda_{\check{B}_0}$) locally generated by affine coordinate 1-forms. We then have a pair of fibre-wise dual torus bundles over the same base:

$$\begin{array}{ccc} X_0 = T^*\check{B}_0/\Lambda_{\check{B}_0}^\vee & & \check{X}_0 = T\check{B}_0/\Lambda_{\check{B}_0} \\ & \searrow p & \swarrow \check{p} \\ & \check{B}_0 & \end{array}$$

We scale both the complex structure on \check{X}_0 and the symplectic structure on X_0 by introducing a $\mathbb{R}_{>0}$ -valued parameter \hbar (so that $\hbar \rightarrow 0$ give the respective large structure limits) and consider the family of spaces (as well as the associated dgLa’s) parametrized by \hbar .

As suggested by Fukaya [21] (and motivated by the relation between Morse theory and de Rham theory [46, 31, 12]), we consider the Fourier expansion (see Definition 2.9) of the Kodaira-Spencer differential graded Lie algebra (dgLa) $(KS_{\check{X}_0} = \Omega^{0,*}(\check{X}_0, T^{1,0}\check{X}_0), \bar{\partial}, [\cdot, \cdot])$ associated to \check{X}_0 along fibres of \check{p} , and try to solve the Maurer-Cartan (abbrev. MC) equation

$$(1.1) \quad \bar{\partial}\Phi + \frac{1}{2}[\Phi, \Phi] = 0.$$

Remark 1.1. *The idea that Fourier-type transforms should be responsible for the interchange between symplectic-geometric data on one side and complex-geometric data on the mirror side (i.e. T-duality) came from the original SYZ proposal [45]. This has been applied successfully in the toric case: see [34, 36, 13, 14, 10, 11, 22, 23, 24, 1, 2, 17, 18] for compact toric varieties and [41, 33, 25, 26, 5, 6, 3, 9, 8, 29, 39] for toric Calabi-Yau varieties. Nevertheless, no scattering phenomenon was involved in those examples.*

1.2. Main results. Before describing our main results, we first choose a Hessian type metric (see Definition 2.3) on the affine manifold \check{B}_0 which allows us to apply the Legendre transform (see Section 2.3) and work with the *Legendre dual* B_0 . This originates from an idea of Gross-Siebert [28] who suggested that, while tropical trees on B_0 correspond to Morse gradient flow trees on \check{B}_0 under the Legendre transform, the former are easier to work with because of their linear nature.

We will also choose a convex open subset $U \subset B_0$, fix a codimension 2 tropical affine subspace $Q \subset U$ and work locally around Q .² In U , a scattering diagram can be viewed schematically as the process of how new walls are being created from the transversal intersection between non-parallel walls supported on tropical hyperplanes in U . The combinatorics of this process is governed by the algebra of the *tropical vertex group* [27], which will be reviewed in Section 3.

We work with dgLa's over the *formal* power series ring $R = \mathbb{C}[[t]]$ where t is a formal deformation variable. Our goal is to investigate the relation between the scattering process and solutions of the MC equation of the Kodaira-Spencer dgLa $KS_{\check{X}_0}[[t]]$.³

To begin with, let $\mathbf{w} = (P, \Theta)$ be a single wall supported on a tropical hyperplane $P \subset U$ containing Q (although Q does not play any role in this single wall case) and equipped with a wall-crossing factor Θ (as an element in the tropical vertex group). Our first aim is to see how Θ is related to solutions of the MC equation (1.1).

Recall that in Witten-Morse theory [46, 31, 12], the shrinking of a fibre-wise loop $m \in \pi_1(p^{-1}(x), s(x))$ towards a singular fibre indicates the presence of a critical point of the symplectic area function f_m in the singular locus (in B), and the union of gradient flow lines emanating from the singular locus should be interpreted as a stable submanifold associated to that critical point. Furthermore, this codimension one stable submanifold should correspond to a bump differential 1-form with support concentrated along P (see [12]).

Inspired by this, given a wall \mathbf{w} , we are going to write down an ansatz $\Pi \in KS_{\check{X}_0}^1[[t]]$ solving (1.1); see Definition 4.2 for the precise formula. Since $\check{X}_0(U) := \check{X}_0 \times_{B_0} U$ does not admit any non-trivial deformations, the MC solution Π is gauge equivalent to 0, i.e. there exists $\varphi \in KS_{\check{X}_0}^0[[t]]$ such that $e^\varphi * 0 = \Pi$; we further use a gauge fixing condition ($\hat{P}\varphi = 0$) to uniquely determine the gauge φ .

In Proposition 4.28, we demonstrate how the semi-classical limit (as $\hbar \rightarrow 0$) of φ determines the wall-crossing factor Θ (or more precisely, $\text{Log}(\Theta)$); see the introduction of Section 4 for a more detailed description. Moreover, the support of the bump-form-like MC solution Π (see Figure 4) is more and more concentrated along P as $\hbar \rightarrow 0$. In Definition 4.19, we make precise the key notion of having *asymptotic support on P* to describe such asymptotic behavior. We further show that *any* MC solution Π with asymptotic support on P would give rise to the *same* wall crossing factor Θ in Section 4.2.3 (see Remark 4.29).

At this point we are ready to explain the main results of this paper. From now on, unlike the case of a single wall, we will be solving the Maurer-Cartan equation *only up to error terms with exponential order in \hbar^{-1}* , i.e. terms of the form $O(e^{-c/\hbar})$. This is sufficient for our purpose because those error terms tend to zero as one approaches the large volume/complex structure limits when $\hbar \rightarrow 0$, and thus they do not contribute to the semi-classical limits of the MC solutions and the associated scattering diagrams.⁴ To make this precise, we introduce in Section 5.2.1 a dgLa $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$ which is a quotient of a sub-dgLa of $KS_{\check{X}_0}(U)[[t]]$, and we will work with and construct MC solutions of $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$.

²In the language of the Gross-Siebert program [28], we are working locally near a *joint* (i.e. a codimension 2 cell) in a polyhedral decomposition of the singular set $\text{Sing}(\mathcal{D})$ of a scattering diagram \mathcal{D} .

³There are other approaches to the scattering process or wall-crossing formulas such as [7, 19, 44].

⁴This point was also anticipated by Fukaya in [21].

Our first main result relates a specific class of MC solutions (satisfying the two assumptions described below) to *consistent* scattering diagrams (see Definition 3.5 for the precise meaning of consistency). Suppose that we have a countable collection $\{P_a\}_{a \in \mathbb{W}}$ of tropical half-hyperplanes (supports of the walls) sharing the codimension 2 tropical affine subspace Q as their common boundary, as shown in Figure 1. We consider a Maurer-Cartan solution $\Phi \in \widehat{\mathfrak{g}^*/\mathcal{E}^*}(U) \otimes_R \mathfrak{m}$ which admits a




FIGURE 1. A collection of walls sharing a common boundary Q

Fourier decomposition

$$(1.2) \quad \Phi = \sum_{a \in \mathbb{W}} \Phi^{(a)},$$

where the sum is finite modulo \mathfrak{m}^{N+1} for every $N \in \mathbb{Z}_{>0}$ (here \mathfrak{m} is the maximal ideal in $R = \mathbb{C}[[\hbar]]$).

Assumption I (see Assumption 5.48 for the precise statement): Each summand $\Phi^{(a)}$ has asymptotic support on the corresponding half-hyperplane P_a (intuitively meaning that the support of $\Phi^{(a)}$ is more and more concentrated along P_a as $\hbar \rightarrow 0$) and has asymptotic expansion (as $\hbar \rightarrow 0$) of the form

$$\Phi^{(a)} = \Psi^{(a)} + F^{(a)},$$

where $\Psi^{(a)}$ is the leading order term consisting of terms with the leading \hbar order and $F^{(a)}$ is the error term consisting of terms with higher \hbar orders.


From this assumption, we deduce that:

Lemma 1.2 (=Lemma 5.40). *For each $a \in \mathbb{W}$, the summand $\Phi^{(a)}$ is a solution of the Maurer-Cartan equation (1.1) over $U \setminus Q$.*

Now we delete Q from U and work over $A := U \setminus Q$. We also choose an open set \tilde{A}_0 in the universal cover \tilde{A} of A and consider the covering map $\mathfrak{p} : \tilde{A}_0 \rightarrow A$.

Assumption II (see Assumption 5.49 for the precise statement): Applying the homotopy operator $\hat{\mathcal{H}}$ (defined by integration over a homotopy $h : \mathbb{R} \times \tilde{A}_0 \rightarrow \tilde{A}_0$ contracting \tilde{A}_0 to a point in (5.23)) to the pullback of the leading order term $\Psi^{(a)}$ by \mathfrak{p} gives a step function which jumps across the lift of P_a in \tilde{A}_0 and whose restriction to the affine half space $\tilde{\mathbb{H}}(P_a) \setminus P_a$ produces an element $\text{Log}(\Theta_a)$ in the tropical vertex Lie-algebra \mathfrak{h} (defined in Definition 3.1). Figure 2 illustrates the situation in a slice of a tubular neighborhood around Q .

Since $\tilde{X}_0(U) \times_A \tilde{A}_0$ does not admit any non-trivial deformations, each summand $\Phi^{(a)}$ in (1.2) is gauge equivalent to 0, so there exists a unique solution φ_a to $e^{\varphi_a} * 0 = \Phi^{(a)}$ satisfying the gauge fixing condition $\hat{\mathcal{P}}\varphi_a = 0$. We carefully estimate the orders of the parameter \hbar in the asymptotic expansion of the gauge φ_a as in the single wall case above, and obtain the following:

FIGURE 2. A slice in a tubular neighborhood around Q

Lemma 1.3 (=Lemma 5.44). *The asymptotic expansion of the gauge φ_a is of the form (see Notations 4.10 for the precise meaning of $O_{loc}(\hbar^{1/2})$):*

$$\varphi_a = \psi_a + O_{loc}(\hbar^{1/2}),$$

where ψ_a , the semi-classical limit of φ_a as $\hbar \rightarrow 0$, is a step function which jumps across the half-hyperplane P_a and is related to an element Θ_a of the tropical vertex group by the formula

$$\text{Log}(\Theta_a) = \psi_a|_{\hat{H}(P_a) \setminus P_a};$$

here $\hat{H}(P_a) \setminus P_a \subset \tilde{A}_0$ is the open half-space (defined in Notation 5.39) which contains the support of ψ_a .

Thus, each $\Phi^{(a)}$, or more precisely, the gauge φ_a , determines a wall $\mathbf{w}_a = (P_a, \Theta_a)$ supported on a tropical half-hyperplane P_a and equipped with a wall crossing factor Θ_a . Hence the Fourier decomposition (1.2) of the Maurer-Cartan solution Φ defines a scattering diagram $\mathcal{D}(\Phi)$ consisting of the walls $\{\mathbf{w}_a\}_{a \in \mathbb{W}}$. Our first main result is the following:

Theorem 1.4 (=Theorem 5.50). *If Φ is any solution to the Maurer-Cartan equation of $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ satisfying both Assumptions I and II (or more precisely Assumptions 5.48 and 5.49), then the associated scattering diagram $\mathcal{D}(\Phi)$ is consistent, meaning that we have the following identity $\Theta_{\gamma, \mathcal{D}(\Phi)} = \text{Id}$, where the left-hand side is the path ordered product (whose definition will be reviewed in Section 3.2.1) along any embedded loop γ in $U \setminus \text{Sing}(\mathcal{D}(\Phi))$ intersecting $\mathcal{D}(\Phi)$ generically; here $\text{Sing}(\mathcal{D}(\Phi)) = Q$ is the singular set of the scattering diagram $\mathcal{D}(\Phi)$.*

Our second main result studies how a scattering process starting with two non-parallel walls $\mathbf{w}_1 = (P_1, \Theta_1)$, $\mathbf{w}_2 = (P_2, \Theta_2)$ intersecting transversally at $Q = P_1 \cap P_2$ gives rise to a MC solution of $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ satisfying both Assumptions I and II, thereby producing a consistent scattering diagram via Theorem 1.4.⁵

In this case, there are two solutions to the MC equation (1.1) $\Pi_{\mathbf{w}_i} \in KS_{\tilde{X}_0}^1(U)[[t]]$, $i = 1, 2$ associated to the two initial walls $\mathbf{w}_1, \mathbf{w}_2$ respectively (e.g. those provided by our ansatz), but their sum $\Pi := \Pi_{\mathbf{w}_1} + \Pi_{\mathbf{w}_2} \in KS_{\tilde{X}_0}(U)[[t]]$ does *not* solve (1.1), even up to error terms with exponential

⁵Indeed Assumptions I and II (or more precisely Assumptions 5.48 and 5.49) are extracted from properties of the MC solutions we constructed.

order in \hbar^{-1} . Nevertheless, a method of Kuranishi [38] allows us to, after fixing the gauge using an explicit homotopy operator (introduced in Definition 5.14), write down a solution $\Phi = \Pi + \dots$, as a sum over trees (5.14) with input Π , of the equation (1.1) up to error terms with exponential order in \hbar^{-1} , or more precisely, of the MC equation of the dgLa $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$.

The MC solution Φ has a Fourier decomposition as in (1.2) of the form $\Phi = \Pi + \sum_{a \in \mathbb{W}} \Phi^{(a)}$, where the sum is over $a = (a_1, a_2) \in \mathbb{W} := (\mathbb{Z}_{>0}^2)_{\text{prim}}$ which parametrizes the tropical half-hyperplanes P_a 's containing Q and lying in-between P_1 and P_2 , as shown in Figure 3. Our second main result is




FIGURE 3. Scattered walls P_a 's from two initial walls

the following:

Theorem 1.5 (=Theorem 5.46). *The Maurer-Cartan solution Φ satisfies both Assumptions I and II (or more precisely Assumptions 5.48 and 5.49) in Theorem 1.4, and hence the scattering diagram $\mathcal{D}(\Phi)$ associated to Φ is consistent, meaning that we have the following identity⁶*

$$\Theta_{\gamma, \mathcal{D}(\Phi)} = \Theta_1^{-1} \Theta_2 \left(\prod_{a \in \mathbb{W}}^{\gamma} \Theta_a \right) \Theta_1 \Theta_2^{-1} = Id,$$

along any embedded loop γ in $U \setminus \text{Sing}(\mathcal{D}(\Phi))$ which intersects $\mathcal{D}(\Phi)$ generically; here $\text{Sing}(\mathcal{D}(\Phi)) = Q = P_1 \cap P_2$.

The proofs that Φ satisfies both Assumptions 5.48 and 5.49 occupy Sections 5.2.3 and 5.2.4; Assumption 5.48 will be handled in Theorem 5.25 in Section 5.2.3 while Assumption 5.49 will be handled in Lemma 5.31 in Section 5.2.4.

Remark 1.6. *Notice that the scattering diagram $\mathcal{D}(\Phi)$ is the unique (by passing to a minimal scattering diagram if necessary) consistent extension, determined by Kontsevich-Soibelman's Theorem 3.7, of the scattering diagram consisting of two initial walls \mathbf{w}_1 and \mathbf{w}_2 .*

1.3. A reader's guide. The rest of this paper is organized as follows.

In Section 2, we review the Kodaira-Spencer dgLa $KS_{\check{X}_0}$ associated to the semi-flat Calabi-Yau manifold \check{X}_0 , followed by a brief review of the Legendre and Fourier transforms.

In Section 3, we review the tropical vertex group and the theory of scattering diagrams (in particular a theorem due to Kontsevich-Soibelman) following the exposition in [27].

Section 4 is about the single wall scenario. In Section 4.1, we write down an ansatz associated to a given single wall solving the MC equation. In Section 4.2.3, we formulate the key notion of

⁶Another common way to write this identity is as a formula for the commutator of two elements in the tropical vertex group: $\Theta_2^{-1} \Theta_1 \Theta_2 \Theta_1^{-1} = \prod_{a \in \mathbb{W}}^{\gamma} \Theta_a$.

asymptotic support on a tropical polyhedral subset which allows us to define a filtration (4.11) to keep track of the \hbar orders. We also prove two key results, namely, Lemma 4.22 (and its extension Lemma 4.25) and Lemma 4.23, which form the basis for the subsequent asymptotic analysis. Applying them, we prove the main results Lemma 4.27 and Proposition 4.28 for the single wall case. Except Definition 4.19 and the statements of Lemmas 4.22 and 4.23, the reader may skip the rather technical Section 4.2 at first reading.

Section 5 is the heart of this paper where we study the scattering process which starts with two initial walls. In Section 5.1, Kuranishi's method of solving the MC equation of a dgLa is reviewed. In Section 5.2, we introduce the dgLa $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ by which we make precise the meaning of *solving the MC equation of $KS_{\check{X}_0}(U)[[t]]$ up to error terms with exponential order in \hbar^{-1}* . We then begin the asymptotic analysis of the MC solutions of $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$; the key results here are Theorem 5.25 and Lemma 5.35. In Section 5.3, we apply the results obtained in Section 5.2 to prove Lemmas 5.43 and 5.44 (which are parallel to Lemma 4.27 and Proposition 4.28 in Section 4), from which we deduce Theorems 1.4 and 1.5.

ACKNOWLEDGEMENT

We thank Si Li, Marco Manetti and Matt Young for various useful conversations when we were preparing the first draft of this paper. We are also heavily indebted to the anonymous referees for many critical yet very constructive comments and suggestions, which lead to a significant and substantial improvement in the exposition including the introduction of the notion of “asymptotic support” and an illuminating reorganization of many of the arguments in the proofs of our main results. Finally we would like to thank Mark Gross for his interest in our work.

The work of K. Chan was supported by a grant from the Hong Kong Research Grants Council (Project No. CUHK14302015) and direct grants from CUHK. The work of N. C. Leung was supported by grants from the Hong Kong Research Grants Council (Project No. CUHK402012 & CUHK14302215) and direct grants from CUHK. The work of Z. N. Ma was supported by a Professor Shing-Tung Yau Post-Doctoral Fellowship, Center of Mathematical Sciences and Applications at Harvard University, Department of Mathematics at National Taiwan University, Yau Mathematical Sciences Center at Tsinghua University, and Institute of Mathematical Sciences and Department of Mathematics at CUHK.

2. THE KODAIRA-SPENCER DGLA IN THE SEMI-FLAT CASE

In this section, we review the classical Kodaira-Spencer deformation theory of complex structures and the associated dgLa [35, 43] in the semi-flat setting, as well as the Legendre and Fourier transforms [32, 40] which play important roles in semi-flat SYZ mirror symmetry.

2.1. The semi-flat Calabi-Yau manifold \check{X}_0 . We let $\text{Aff}(\mathbb{R}^n) = \mathbb{R}^n \rtimes \text{GL}_n(\mathbb{R})$ be the group of affine linear transformations of \mathbb{R}^n and consider the subgroup $\text{Aff}_{\mathbb{R}}(\mathbb{Z}^n)_0 := \mathbb{R}^n \rtimes \text{SL}_n(\mathbb{Z})$.

Definition 2.1 ([27]). *An n -dimensional smooth manifold B is called tropical affine if it admits an atlas $\{(U_i, \psi_i)\}$ of coordinate charts $\psi_i : U_i \rightarrow \mathbb{R}^n$ such that $\psi_i \circ \psi_j^{-1} \in \text{Aff}_{\mathbb{R}}(\mathbb{Z}^n)_0$ for all i, j .*

Given a (possibly non-compact) tropical affine manifold \check{B}_0 , we set

$$\check{X}_0 := T\check{B}_0/\Lambda_{\check{B}_0},$$

where the lattice subbundle $\Lambda_{\check{B}_0} \subset T\check{B}_0$ is locally generated by the coordinate vector fields $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}$ for a given choice of local affine coordinates $\check{x} = (x^1, \dots, x^n)$ in a contractible open subset $\check{U} \subset \check{B}_0$. Then the natural projection map $\check{p} : \check{X}_0 \rightarrow \check{B}_0$ is a torus fibration. We also let y^j 's be the canonical coordinates on the fibres of \check{p} over \check{U} with respect to the frame $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}$ of $T\check{B}_0$.

Choosing $\beta = \sum_{i,j=1}^n \beta_i^j(\check{x}) dx^i \otimes \frac{\partial}{\partial x^j} \in \Omega^1(T\check{B}_0)$ satisfying $\nabla\beta = 0 \in \Omega^2(T\check{B}_0)$, where ∇ is the natural affine flat connection on \check{B}_0 , we get a one-parameter family of complex structures parametrized by $\hbar \in \mathbb{R}_{>0}$ defined by the family of matrices

$$(2.1) \quad \check{J}_\beta = \begin{pmatrix} -\hbar\beta & \hbar I \\ -\hbar^{-1}(I + \hbar^2\beta^2) & \hbar\beta \end{pmatrix}$$

with respect to the local frame $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}, \frac{\partial}{\partial y^1}, \dots, \frac{\partial}{\partial y^n}$, where we write β as a matrix with respect to the frame $\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}$. Locally, the corresponding holomorphic volume form is given by

$$(2.2) \quad \check{\Omega}_\beta = \bigwedge_{j=1}^n \left((dy^j - \sum_{k=1}^n \beta_k^j dx^k) + i\hbar^{-1} dx^j \right),$$

and a holomorphic frame of $T^{1,0}\check{X}_0$ can be written as

$$(2.3) \quad \check{\partial}_j := \frac{\partial}{\partial \log z^j} = \frac{i}{4\pi} \left(\frac{\partial}{\partial y^j} - i\hbar \left(\sum_{k=1}^n \beta_j^k \frac{\partial}{\partial y^k} + \frac{\partial}{\partial x^j} \right) \right),$$

for $j = 1, \dots, n$. So the local complex coordinates are given by

$$(2.4) \quad z^j = \exp \left(-2\pi i \left(y^j - \sum_k \beta_k^j x^k + i\hbar^{-1} x^j \right) \right).$$

The condition that $\sum_{k=1}^n \beta_k^j(\check{x}) dx^k$ being closed for each $j = 1, \dots, n$ is equivalent to integrability of the almost complex structure \check{J}_β .

2.2. The Kodaira-Spencer dgLa. For a complex manifold \check{X}_0 , the Kodaira-Spencer complex is the space $KS_{\check{X}_0} := \Omega^{0,*}(\check{X}_0, T^{1,0}\check{X}_0)$ of $T^{1,0}\check{X}_0$ -valued $(0,*)$ -forms, which is equipped with the Dolbeault differential $\bar{\partial}$ and a Lie bracket defined in local holomorphic coordinates $z_1, \dots, z_n \in \check{X}_0$ by $[\phi d\bar{z}^I, \psi d\bar{z}^J] = [\phi, \psi] d\bar{z}^I \wedge d\bar{z}^J$, where $\phi, \psi \in \Gamma(T^{1,0}\check{X}_0)$. The triple

$$(KS_{\check{X}_0}, \bar{\partial}, [\cdot, \cdot])$$

defines the *Kodaira-Spencer differential graded Lie algebra (abbrev. dgLa)*, which governs the deformation theory of complex structures on \check{X}_0 . Given an open subset $\check{U} \subset \check{X}_0$, we may also talk about the local Kodaira-Spencer complex $KS_{\check{X}_0}(\check{U})$.

Notations 2.2. We let $R = \mathbb{C}[[t]]$ to be the ring of formal power series and $\mathfrak{m} = (t)$ denote the maximal ideal generated by t , and consider dgLa's over R to avoid convergence issues.

An element $\varphi \in \Omega^{0,1}(\check{X}_0, T^{1,0}\check{X}_0) \otimes \mathfrak{m}$ defines a formal deformation of complex structures if and only if it is a solution to the Maurer-Cartan equation (1.1). The exponential group $KS_{\check{X}_0}^0 \otimes \mathfrak{m}$ acts on the set of Maurer-Cartan solutions $MC_{KS_{\check{X}_0}}(R)$ as automorphisms of the formal family of complex structures over R , and therefore one can define the space of deformations of \check{X}_0 over R by $Def_{KS_{\check{X}_0}}(R) := MC_{KS_{\check{X}_0}}(R) / \exp(KS_{\check{X}_0}^0 \otimes \mathfrak{m})$ via the dgLa $KS_{\check{X}_0}$.

2.3. The Legendre transform. To define the Legendre dual B_0 of \check{B}_0 so that we can work in the tropical world, we need a metric g on \check{B}_0 of Hessian type (see, e.g. [4, Chapter 6]):

Definition 2.3. A Riemannian metric $g = (g_{ij})_{i,j}$ on \check{B}_0 is said to be Hessian type if it is locally given by $g = \sum_{i,j} \frac{\partial^2 \check{\phi}}{\partial x^i \partial x^j} dx^i \otimes dx^j$ in local affine coordinates x^1, \dots, x^n for some convex function $\check{\phi}$.

To construct Kähler structures, we further need a compatibility condition between g and β in (2.1), namely, we assume that

$$(2.5) \quad \sum_{i,j,k} \beta_i^j g_{jk} dx^i \wedge dx^k = 0,$$

when we write $\beta = \sum_{i,j} \beta_i^j(\check{x}) dy_j \wedge dx^i$ in the local coordinates $x^1, \dots, x^n, y_1, \dots, y_n$. Given such a Hessian type metric g , a Kähler form on \check{X}_0 is given by $\check{\omega} = 2i\partial\bar{\partial}\check{\phi} = \sum_{j,k} g_{jk} dy^j \wedge dx^k$.

We can now introduce the Legendre transform following Hitchin [32]; see also [4, Chapter 6]. Given a strictly convex smooth function $\check{\phi} : \check{U}(\subset \check{B}_0) \rightarrow \mathbb{R}$, we trivialize $T^*\check{U} \cong \check{U} \times \mathbb{R}^n$ via affine frames and define the *Legendre transform* $L_{\check{\phi}} : \check{U} \rightarrow \mathbb{R}^n$ by $x = L_{\check{\phi}}(\check{x}) := d\check{\phi}(\check{x}) \in \mathbb{R}^n$, or equivalently, by $x_j = \frac{\partial \check{\phi}}{\partial x^j}$, where $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ denote the dual coordinates. The image $U := L_{\check{\phi}}(\check{U}) \subset \mathbb{R}^n$ is an open subset and $L_{\check{\phi}}$ is a diffeomorphism. The Legendre dual $\phi : U \rightarrow \mathbb{R}$ of $\check{\phi}$ is defined by the equation $\phi(x) := \sum_{j=1}^n x_j x^j - \check{\phi}(\check{x})$, and the dual transform L_{ϕ} is inverse to $L_{\check{\phi}}$.

If $\check{\phi}$ is the semi-flat potential in Definition 2.3 which defines a Hessian type metric, then the dual coordinate charts $U = L_{\check{\phi}}(\check{U})$ actually glue to give another tropical affine manifold B_0 , which we call the *Legendre dual* of \check{B}_0 , whose underlying smooth manifold is same as that of \check{B}_0 (see [4, Chapter 6]). The lattice bundles $\Lambda_{B_0} \cong \Lambda_{\check{B}_0}^\vee$ are interchanged in this process, so that we can write $\check{X}_0 = T^*B_0/\Lambda_{B_0}^\vee$, and using the affine coordinates (x_1, \dots, x_n) on B_0 , we can write

$$\check{\Omega}_\beta = \bigwedge_{k=1}^n \left(dy^k - \sum_{j=1}^n (\beta^{jk} - i\hbar^{-1}g^{jk}) dx_j \right), \quad \check{\omega} = \sum_{k=1}^n dy^k \wedge dx_k$$

2.4. The Fourier transform.

Definition 2.4. *The sheaf of integral affine functions $\text{Aff}_{B_0}^{\mathbb{Z}}$, as a sheaf over B_0 (which is the same as \check{B}_0 as a smooth manifold), is the subsheaf of the sheaf of smooth functions over B_0 whose local sections $\text{Aff}_{B_0}^{\mathbb{Z}}(U)$ over a contractible open set $U \subset B_0$ are defined to be affine linear functions of the form $m(\check{x}) = m_1 x^1 + \dots + m_n x^n + b$ for some $m_i \in \mathbb{Z}$ and $b \in \mathbb{R}$, in local affine coordinates on \check{B}_0 (caution: not B_0). This sheaf fits into the following exact sequence of sheaves over B_0*

$$0 \rightarrow \underline{\mathbb{R}} \rightarrow \text{Aff}_{B_0}^{\mathbb{Z}} \rightarrow \Lambda_{B_0} \rightarrow 0.$$

Since $\check{X}_0 = T\check{B}_0/\Lambda_{\check{B}_0}$, exponentiation of complexification of local affine linear functions on \check{B}_0 give local holomorphic functions on \check{X}_0 as follows.

Definition 2.5. *Given $m \in \text{Aff}_{B_0}^{\mathbb{Z}}(U)$, expressed locally as $m(x) = \sum_j m_j x^j + b$, we let $z^m := e^{\frac{2\pi b}{\hbar}} (z^1)^{m_1} \dots (z^n)^{m_n} \in \mathcal{O}_{\check{X}_0}(\check{p}^{-1}(U))$, where z^j is given in equation (2.4). This defines an embedding $\text{Aff}_{B_0}^{\mathbb{Z}}(U) \hookrightarrow \mathcal{O}_{\check{X}_0}(\check{p}^{-1}(U))$, and we denote the image subsheaf by \mathcal{O}^{aff} , as a sheaf over B_0 .*

We can embed the lattice bundle $\Lambda_{B_0}^\vee \hookrightarrow \check{p}_* T^{1,0} \check{X}_0$ into the push forward of the sheaf of holomorphic vector fields; in local coordinates U , it is given by (cf. equation (2.3))

$$(2.6) \quad n = (n^j) \mapsto \check{\partial}_n := \sum_j n^j \frac{\partial}{\partial \log z^j} = \frac{i}{4\pi} \sum_j n^j \left(\frac{\partial}{\partial y^j} - i\hbar \sum_k \left(\beta_j^k \frac{\partial}{\partial y^k} + g_{jk} \frac{\partial}{\partial x_k} \right) \right)$$

for a local section $n \in \Lambda_{B_0}^\vee(U)$. This embedding is globally defined, and by abuse of notations, we will write $\Lambda_{B_0}^\vee$ to stand for its image subsheaf. For later purpose, we introduce the notation

$$(2.7) \quad \partial_n := \frac{\hbar}{4\pi} \sum_j n^j g_{jk} \frac{\partial}{\partial x_k}.$$

Notations 2.6. *Since we work in a contractible open coordinate chart U , we will fix a rank n lattice $M \cong \mathbb{Z}^n$ and its dual $N = \text{Hom}(M, \mathbb{Z})$, and identify $U \subset M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R} \cong \mathbb{R}^n$ as an open subset containing the origin 0 and write $N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}$. We also trivialize $\Lambda_{B_0}|_U \cong \underline{M}$ and $\Lambda_{B_0}^{\vee}|_U \cong \underline{N}$ and identify $\check{X}_0(U) = \check{p}^{-1}(U) \cong U \times (N_{\mathbb{R}}/N)$. Since $TU \cong \Lambda_{B_0} \otimes_{\mathbb{Z}} \mathbb{R}$, a local section $m \in \underline{M}(U)$ naturally corresponds to an affine integral vector field over U , which will be denoted by m as well. The exact sequence in Definition 2.4 splits and we will call m 's or the associated z^m 's the Fourier modes.*

Definition 2.7. *We consider the sheaf $\mathcal{O}^{\text{aff}} \otimes_{\mathbb{Z}} \Lambda_{B_0}^{\vee}$ over B_0 and define a Lie bracket $[\cdot, \cdot]$ on it by restriction of the usual Lie bracket on $\check{p}_* \mathcal{O}(T^{1,0} \check{X}_0)$.*

Notice that the Lie bracket on $\mathcal{O}^{\text{aff}} \otimes_{\mathbb{Z}} \Lambda_{B_0}^{\vee}$ is well defined because in a small enough affine coordinate chart, we have the following formula from [27]

$$(2.8) \quad \left[z^m \otimes \check{\partial}_n, z^{m'} \otimes \check{\partial}_{n'} \right] = z^{m+m'} \check{\partial}_{(m',n)n' - (m,n)n},$$

which shows that $\mathcal{O}^{\text{aff}} \otimes_{\mathbb{Z}} \Lambda_{B_0}^{\vee}$ is closed under the Lie bracket on $\check{p}_* \mathcal{O}(T^{1,0} \check{X}_0)$.

Notations 2.8. *The pairing (m, n) in (2.8) is the natural pairing between $m \in \Lambda_{B_0}(U)$ and $n \in \Lambda_{B_0}^{\vee}(U)$. Given a local section $m \in \Lambda_{B_0}(U)$, we let $m^{\perp} \subset \Lambda_{B_0}^{\vee}(U)$ be the sub-lattice perpendicular to m with respect to (\cdot, \cdot) .*

Definition 2.9. *On a contractible open subset $U \subset B_0 \cong \check{B}_0$, the Fourier transform*

$$\mathcal{F} : \mathbf{G}^*(U) := \bigoplus_{m \in \Lambda_{B_0}(U)} \Omega^*(U) \cdot z^m \otimes_{\mathbb{Z}} N \hookrightarrow KS_{\check{X}_0}(U)$$

is defined by sending $\alpha \in \Omega^(U)$ to $(\check{p}^*(\alpha))^{0,1}$ (where $(\cdot)^{0,1}$ denotes the $(0,1)$ -part of the 1-form) and $n \in N$ to $\check{\partial}_n$. \mathcal{F} is injective and hence induces a dgLa structure on $\mathbf{G}^*(U)$ from that on $KS_{\check{X}_0}(U)$.⁷*

We also let $\mathbf{G}_N^(U) := \mathbf{G}^*(U) \otimes R/\mathfrak{m}^{N+1}$ and $\widehat{\mathbf{G}}^*(U) := \varprojlim_N \mathbf{G}_N^*(U)$.*

Remark 2.10. *For more details on how Fourier (or SYZ) transforms can be applied to understand (semi-flat) SYZ mirror symmetry, we refer the readers to Fukaya's original paper [21] and a recent survey article [42] by the third named author.*

3. SCATTERING DIAGRAMS

In the section, we review the notion of scattering diagrams introduced in [37, 28]. We will adopt the setting and notations from [27] with slight modifications to fit into our context.

3.1. The sheaf of tropical vertex groups. We start with the same set of data (B_0, g, β) as in Section 2.3, and use $x = (x_1, \dots, x_n)$ as local affine coordinates on B_0 and $\check{x} = (x^1, \dots, x^n)$ as local affine coordinates on \check{B}_0 as before. Given the formal power series ring $R = \mathbb{C}[[t]]$ and its maximal ideal \mathfrak{m} , we consider the sheaf of Lie algebras $\mathfrak{g} := (\mathcal{O}^{\text{aff}} \otimes_{\mathbb{Z}} \Lambda_{B_0}^{\vee}) \widehat{\otimes}_{\mathbb{C}} R$ over B_0 .

Definition 3.1. *The subsheaf $\mathfrak{h} \hookrightarrow \mathfrak{g}$ of Lie algebras is defined as the image of the embedding $\left(\bigoplus_{m \in \Lambda_{B_0}(U)} \mathbb{C} \cdot z^m \otimes_{\mathbb{Z}} (m^{\perp}) \right) \widehat{\otimes}_{\mathbb{C}} R \rightarrow \mathfrak{g}(U)$ over each affine coordinate chart $U \subset B_0$.⁸ The sheaf of tropical vertex groups over B_0 is defined as the sheaf of exponential groups $\exp(\mathfrak{h} \otimes_R \mathfrak{m})$ which act as automorphisms on \mathfrak{h} and \mathfrak{g} .*

⁷Direct computation shows that we have the formula $\bar{\partial}(\sum_m z^m \alpha_m^n \check{\partial}_n) = \sum_m z^m (d\alpha) \check{\partial}_n$.

⁸It is a subsheaf of Lie subalgebras of \mathfrak{g} as can be seen from the formula (2.8).

3.2. Kontsevich-Soibelman's wall crossing formula. This formulation of the wall crossing formula originated from [37] but we will mostly follow [27] as we want to work on B_0 instead of \tilde{B}_0 . From now on, we will work locally in a contractible coordinate chart $U \subset B_0$. We use the same notations as in Section 2.

Definition 3.2. Given $m \in M \setminus \{0\}$ and $n \in m^\perp$, we let $\mathfrak{h}_{m,n} := (\mathbb{C}[z^m] \cdot z^m) \check{\partial}_n \widehat{\otimes}_{\mathbb{C}} \mathfrak{m} \hookrightarrow \mathfrak{g}$ whose general elements are of the form $\sum_{j,k \geq 1} a_{jk} z^{km} \check{\partial}_n t^j$, where $a_{jk} \neq 0$ for only finitely many k 's for each fixed j . This defines an abelian Lie subalgebra of \mathfrak{g} by the formula (2.8).

Definition 3.3. A wall \mathbf{w} in U is a triple (m, P, Θ) , where

- $m \in M \setminus \{0\}$ parallel to P ,
- P is a connected oriented codimension one convex tropical polyhedral subset of U (by a convex tropical polyhedral subset we mean a convex subset which is locally defined by affine linear equations and inequalities defined over \mathbb{Q}),
- $\Theta \in \exp(\mathfrak{h}_{m,n})|_P$ is a germ of sections near P , where $n \in \Lambda_{B_0}^\vee(U) \cong N$ is the unique primitive element satisfying $n \in (TP)^\perp$ and $(\nu_P, n) < 0$, and $\nu_P \in TU \cong U \times M_{\mathbb{R}}$ here is a vector normal to P such that the orientation of $TP \oplus \mathbb{R} \cdot \nu_P$ agrees with that of U .

Definition 3.4. A scattering diagram \mathcal{D} is a set of walls $\{(m_\alpha, P_\alpha, \Theta_\alpha)\}_\alpha$ such that there are only finitely many α 's with $\Theta_\alpha \neq \text{id} \pmod{\mathfrak{m}^N}$ for every $N \in \mathbb{Z}_{>0}$. We define the support of \mathcal{D} to be $\text{supp}(\mathcal{D}) := \bigcup_{\mathbf{w} \in \mathcal{D}} P_{\mathbf{w}}$, and the singular set of \mathcal{D} to be $\text{Sing}(\mathcal{D}) := \bigcup_{\mathbf{w} \in \mathcal{D}} \partial P_{\mathbf{w}} \cup \bigcup_{\mathbf{w}_1 \pitchfork \mathbf{w}_2} P_{\mathbf{w}_1} \cap P_{\mathbf{w}_2}$, where $\mathbf{w}_1 \pitchfork \mathbf{w}_2$ means transversally intersecting walls.⁹

3.2.1. Path ordered products. An embedded path $\gamma : [0, 1] \rightarrow B_0 \setminus \text{Sing}(\mathcal{D})$ is said to be *intersecting* \mathcal{D} *generically* if $\gamma(0), \gamma(1) \notin \text{supp}(\mathcal{D})$, $\text{Im}(\gamma) \cap \text{Sing}(\mathcal{D}) = \emptyset$ and it intersects all the walls in \mathcal{D} transversally. Given such an embedded path γ , we define the *path ordered product* along γ as an element of the form $\Theta_{\gamma, \mathcal{D}} = \prod_{\mathbf{w} \in \mathcal{D}}^{\gamma} \Theta_{\mathbf{w}} \in \exp(\mathfrak{h} \otimes_{\mathbb{R}} \mathfrak{m})_{\gamma(1)}$ in the stalk of $\exp(\mathfrak{h} \otimes_{\mathbb{R}} \mathfrak{m})$ at $\gamma(1)$, following [27]. More precisely, for each $k \in \mathbb{Z}_{>0}$, we define $\Theta_{\gamma, \mathcal{D}}^k \in \exp(\mathfrak{h} \otimes_{\mathbb{R}} (\mathfrak{m}/\mathfrak{m}^{k+1}))_{\gamma(1)}$ and let $\Theta_{\gamma, \mathcal{D}} := \lim_{k \rightarrow +\infty} \Theta_{\gamma, \mathcal{D}}^k$, where $\Theta_{\gamma, \mathcal{D}}^k$ is defined as follows.

Given k , there is a finite subset $\mathcal{D}^k \subset \mathcal{D}$ consisting of walls \mathbf{w} with $\Theta_{\mathbf{w}} \neq \text{Id} \pmod{\mathfrak{m}^{k+1}}$ from Definition 3.4. We then have a sequence of real numbers $0 = t_0 < t_1 < t_2 < \dots < t_s < t_{s+1} = 1$ such that $\{\gamma(t_1), \dots, \gamma(t_s)\} = \gamma \cap \text{supp}(\mathcal{D}^k)$. For each $1 \leq i \leq s$, there are walls $\mathbf{w}_{i,1}, \dots, \mathbf{w}_{i,l_i}$ in \mathcal{D}^k such that $\gamma(t_i) \in P_{i,j} := \text{supp}(\mathbf{w}_{i,j})$ for all $j = 1, \dots, l_i$. Since γ does not hit $\text{Sing}(\mathcal{D})$, we have $\text{codim}(\text{supp}(\mathbf{w}_{i,j_1}) \cap \text{supp}(\mathbf{w}_{i,j_2})) = 1$ for any j_1, j_2 , i.e. the walls $\mathbf{w}_{i,1}, \dots, \mathbf{w}_{i,l_i}$ are overlapping with each other and contained in a common tropical hyperplane. Then we have an element $\Theta_{\gamma(t_i)} := \prod_{j=1}^{l_i} \Theta_{\mathbf{w}_{i,j}}^{\sigma_j}$, where $\sigma_j = 1$ if orientation of $P_{i,j} \oplus \mathbb{R} \cdot \gamma'(t_i)$ agree with that of B_0 and $\sigma_j = -1$ otherwise. (Note that this element is well defined without prescribing the order of the product since the elements $\Theta_{\mathbf{w}_{i,j}}$'s are commuting with each other.) We treat $\Theta_{\gamma(t_i)}$ as an element in $\exp(\mathfrak{h} \otimes_{\mathbb{R}} \mathfrak{m})_{\gamma(1)}$ by parallel transport and take the ordered product along the path γ as $\Theta_{\gamma, \mathcal{D}}^k := \Theta_{\gamma(t_s)} \cdots \Theta_{\gamma(t_i)} \cdots \Theta_{\gamma(t_1)}$.

Definition 3.5. A scattering diagram \mathcal{D} is said to be *consistent* if we have $\Theta_{\gamma, \mathcal{D}} = \text{Id}$, for any embedded loop γ intersecting \mathcal{D} generically. Two scattering diagrams \mathcal{D} and $\tilde{\mathcal{D}}$ are said to be *equivalent* if $\Theta_{\gamma, \mathcal{D}} = \Theta_{\gamma, \tilde{\mathcal{D}}}$ for any embedded path γ intersecting both \mathcal{D} and $\tilde{\mathcal{D}}$ generically.

Remark 3.6. Given a scattering diagram \mathcal{D} , there is a unique representative \mathcal{D}_{\min} from its equivalence class which is minimal. First, we may remove those walls with trivial automorphisms Θ as they do not contribute to the path ordered product. Second, if two walls \mathbf{w}_1 and \mathbf{w}_2 share the same P and m , we can simply take the multiplication $\Theta = \Theta_1 \circ \Theta_2$ and define a single wall \mathbf{w} . After doing so, we obtain a minimal scattering diagram equivalent to \mathcal{D} .

⁹There is a natural (possibly up to further subdivisions) polyhedral decomposition of $\text{Sing}(\mathcal{D})$ whose codimension 2 cells are called *joints* in the Gross-Siebert program [28].

3.2.2. *The wall crossing formula.* Next, we consider the case where \mathcal{D} is a scattering diagram consisting of only two walls $\mathbf{w}_1 = (m_1, P_1, \Theta_1)$ and $\mathbf{w}_2 = (m_2, P_2, \Theta_2)$ where the supports P_i 's are tropical hyperplanes of the form $P_i = Q - \mathbb{R} \cdot m_i$ intersecting transversally in a codimension two tropical subspace $Q := P_1 \cap P_2 \subset U$. In this case, we have the following theorem due to Kontsevich-Soibelman [37]:

Theorem 3.7 (Kontsevich and Soibelman [37]). *Given a scattering diagram \mathcal{D} consisting of two walls $\mathbf{w}_1 = (m_1, P_1, \Theta_1)$ and $\mathbf{w}_2 = (m_2, P_2, \Theta_2)$ supported on tropical hyperplanes P_1, P_2 intersecting transversally in a codimension two tropical subspace $Q := P_1 \cap P_2$, there exists a unique minimal consistent scattering diagram $\mathcal{S}(\mathcal{D}) \supset \mathcal{D}_{\min}$, obtained by adding walls to \mathcal{D}_{\min} supported on tropical half-hyperplanes of the form $Q - \mathbb{R}_{\geq 0} \cdot (a_1 m_1 + a_2 m_2)$ for $a = (a_1, a_2) \in (\mathbb{Z}_{>0}^2)_{\text{prim}}$.*

Remark 3.8. *Interesting relations between these wall crossing factors and relative Gromov-Witten invariants of weighted projective planes were established in [27]. In general it is expected that these wall crossing factors encode counts of holomorphic disks on the mirror A -side, which was conjectured by Fukaya in [21, Section 3] to be closely related to Witten's Morse theory.*

4. SINGLE WALL DIAGRAMS AS DEFORMATIONS

As before, we will work with a contractible open coordinate chart $U \subset B_0$. In this section, we consider a scattering diagram with only one wall $\mathbf{w} = (m, P, \Theta)$, where P is a connected oriented tropical hyperplane in U . Recall that we can write

$$(4.1) \quad \text{Log}(\Theta) = \sum_{j,k \geq 1} a_{jk} z^{km} \check{\partial}_n t^j,$$

where $a_{jk} \neq 0$ for only finitely many k 's for each fixed j .

The hyperplane P divides the base U into two half spaces H_+ and H_- according to the orientation of P , meaning that ν_P should be pointing into H_+ where $\nu_P \in TU$ is the normal to P we choose so that the orientation of $TP \oplus \mathbb{R} \cdot \nu_P$ agrees with that of U . We consider a step-function-like section $\psi \in \Omega^{0,0}(\check{X}_0(U) \setminus \check{p}^{-1}(P), T^{1,0}\check{X}_0)[[t]]$ of the form

$$(4.2) \quad \psi = \begin{cases} \text{Log}(\Theta) & \text{on } H_+, \\ 0 & \text{on } H_-. \end{cases}$$

Our goal is to write down an ansatz $\Pi = \Pi_{\hbar}$ (depending on \hbar) solving the Maurer-Cartan equation (1.1) such that $\Pi = e^\psi * 0 \in \Omega^{0,1}(\check{X}_0(U), T^{1,0})$ represents a smoothing of $e^\psi * 0$ (which is delta-function-like and not well defined by itself), and show that the semi-classical limit of φ is precisely ψ as $\hbar \rightarrow 0$.

4.1. Ansatz corresponding to a single wall. We are going to use the Fourier transform $\mathcal{F} : \widehat{\mathbf{G}}^*(U) \rightarrow KS_{\check{X}_0}(U)[[t]]$ defined in Definition 2.9 to obtain an element $\Pi \in \widehat{\mathbf{G}}^*(U)$, and perform all the computations on $\mathbf{G}_N^*(U)$ or $\widehat{\mathbf{G}}^*(U)$ following Fukaya's ideas [21]. We will omit the Fourier transform \mathcal{F} in our notations, and we will work with tropical geometry on B_0 instead of Witten-Morse theory on \check{B}_0 following Gross-Siebert's idea [28]. We start by choosing some convenient affine coordinates $u_{m,i}$'s (or simply u_i 's, if there is no confusion) on U for each Fourier mode m .

Notations 4.1. *For each Fourier mode $m \in M \setminus \{0\}$, we choose affine coordinates (u_1, \dots, u_n) for U with the properties that u_1 is along $-m$. We will denote the remaining coordinates by $u_{m,\perp} := (u_2, \dots, u_n)$. We further require that the coordinates u_i 's for m and km is the same with $k \in \mathbb{Z}_+$ for convenience.*

Given a wall $\mathbf{w} = (m, P, \Theta)$ as above, we choose u_2 to be the coordinate normal to P and pointing into H_+ . We consider a 1-form depending on $\hbar \in \mathbb{R}_{>0}$ given by

$$(4.3) \quad \delta_m = \delta_{m,\hbar} := \left(\frac{1}{\hbar\pi} \right)^{\frac{1}{2}} e^{-\frac{u_2^2}{\hbar}} du_2,$$

which has the property that $\int_L \delta_m = 1 + O(e^{-c/\hbar})$ for any line $L \cong \mathbb{R}$ intersecting P transversally. This gives a bump form which can be viewed as a smoothing of the delta 1-form over P , as shown in Figure 4. We will sometimes write $\delta_m = e^{-\frac{u_2^2}{\hbar}} \mu_m$, where $\mu_m := \left(\frac{1}{\hbar\pi} \right)^{\frac{1}{2}} du_2$, to avoid repeated appearances of the constant $\left(\frac{1}{\hbar\pi} \right)^{\frac{1}{2}}$.




FIGURE 4. δ_m concentrating along P

Definition 4.2. Given a wall $\mathbf{w} = (m, P, \Theta)$ where $\text{Log}(\Theta)$ is as in (4.1), we let

$$(4.4) \quad \Pi := -\delta_m \cdot \text{Log}(\Theta) = -\delta_m \sum_{j,k \geq 1} a_{jk} z^{km} \check{\partial}_n t^j \in \widehat{\mathbf{G}}^1(U)$$

be the ansatz associated to the wall \mathbf{w} , by viewing \mathbf{G}^* as a module over $\Omega^*(U)$.

Remark 4.3. Our ansatz depends on the choice of the affine coordinates (u_1, \dots, u_n) because δ_m does so, but the property that it has support concentrated along the tropical hyperplane P is an abstract notion which does not depend on the choice of coordinates, as we will see shortly.

Proposition 4.4. The ansatz Π satisfies the Maurer-Cartan (MC) equation $\bar{\partial}\Pi + \frac{1}{2}[\Pi, \Pi] = 0$.

Proof. In fact we will show that both terms $\bar{\partial}\Pi$ and $\frac{1}{2}[\Pi, \Pi]$ vanish. First we have $\bar{\partial}(\delta_m z^{km} \check{\partial}_n) = (d\delta_m) z^{km} \check{\partial}_n = 0$ from the fact that $d(\delta_m) = 0$ which is obvious from (4.3). Next we show that $[\delta_m z^{k_1 m} \check{\partial}_n, \delta_m z^{k_2 m} \check{\partial}_n] = 0$ for any k_1, k_2 . This is simply because $\delta_m = e^{-\frac{u_2^2}{\hbar}} \mu_m$ and μ_m is a covariant constant form (with respect to the affine connection), so we have $[\delta_m z^{k_1 m} \check{\partial}_n, \delta_m z^{k_2 m} \check{\partial}_n] = \mu_m \wedge \mu_m \left[e^{-\frac{u_2^2}{\hbar}} z^{k_1 m} \check{\partial}_n, e^{-\frac{u_2^2}{\hbar}} z^{k_2 m} \check{\partial}_n \right] = 0$. \square

4.2. Relation with the wall crossing factor. Since $\check{X}_0(U) \cong U \times T^n$ has no non-trivial deformations, the element $\check{\Pi}$ must be gauge equivalent to 0. In this subsection, we will explain how the semi-classical limit of the gauge is related to the wall crossing factor $\text{Log}(\Theta)$.

4.2.1. Solving for the gauge φ . So we are going to solve the equation $e^\varphi * 0 = \Pi$ for $\varphi \in \widehat{\mathbf{G}}^*(U)$ with desired asymptotic behavior. Using the definition in [43, Section 1] for gauge action, we are indeed solving

$$(4.5) \quad - \left(\frac{e^{ad_\varphi} - \text{Id}}{ad_\varphi} \right) \bar{\partial}\varphi = \Pi.$$

Solutions φ to (4.5) is not unique. We will make a choice by choosing a homotopy operator \hat{H} . Since $\mathbf{G}^*(U) = \bigoplus_m (\Omega^*(U)z^m) \otimes_{\mathbb{Z}} N$ is a tensor product of N with a direct sum, it suffices to define a homotopy operator \hat{H}_m for each Fourier mode m contracting $\Omega^*(U)$ to its cohomology $H^*(U) \cong \mathbb{C}$.

Definition 4.5. We fix a based point $q \in H_-$. By contractibility, we have the map $\rho_q : [0, 1] \times U \rightarrow U$ satisfying $\rho_q(0, \cdot) = q$ and $\rho_q(1, \cdot) = \text{Id}$, which contracts U to $\{q\}$.

This defines a homotopy operator $\hat{H}_m : \Omega^*(U)z^m \rightarrow \Omega^*(U)[-1]z^m$ by $\hat{H}_m(\alpha z^m) := \int_0^1 \rho_q^*(\alpha) z^m$. We also define the projection $\hat{P}_m : \Omega^*(U)z^m \rightarrow H^*(U)z^m$ by setting $\hat{P}_m(\alpha z^m) = \alpha|_q z^m$ for $\alpha \in \Omega^0(U)$ and 0 otherwise, and $\iota_m : H^*(U)z^m \rightarrow \Omega^*(U)z^m$ by setting $\iota_m : H^*(U)z^m \rightarrow \Omega^*(U)z^m$ to be the embedding of constant functions on U at degree 0 and 0 otherwise.

These operators can be put together to define operators on $\mathbf{G}^*(U)$, $\mathbf{G}_N^*(U)$ or $\hat{\mathbf{G}}^*(U)$ and they are denoted by \hat{H} , \hat{P} and ι respectively.

Remark 4.6. The based point q is chosen so that the semi-classical limit of the gauge φ_0 (as $\hbar \rightarrow 0$) behaves like a step-function across the wall P . There are many possible choices of \hat{H} , corresponding to choices of ρ_q for this purpose. In Definition 5.12, we will write down another particular choice (suitable for later purposes) in the case when the open subset U is spherical (see Section 11).

In the rest of this section, we will fix $q \in H_-$ in the half space H_- and impose the gauge fixing condition $\hat{P}\varphi = 0$ to solve for φ satisfying (4.5); in other words, we look for a solution satisfying $\varphi = \hat{H}\bar{\partial}\varphi + \bar{\partial}\hat{H}\varphi = \hat{H}\bar{\partial}\varphi$ to solve the equation (4.5) order by order; here $\hat{H}\varphi = 0$ by degree reasons. This is possible because of the following lemma which we learn from [43].

Lemma 4.7. Among all solutions of $e^\varphi * 0 = \Pi$, there exists a unique one satisfying $\hat{P}\varphi = 0$.

Proof. Notice that for any $\sigma = \sigma_1 + \sigma_2 + \dots \in t \cdot \hat{\mathbf{G}}^*(U)$ with $\bar{\partial}\sigma = 0$, we have $e^\sigma * 0 = 0$, and hence $e^{\varphi \bullet \sigma} * 0 = \Pi$ is still a solution for the same equation. With $\varphi \bullet \sigma$ given by the Baker-Campbell-Hausdorff formula as $\varphi \bullet \sigma = \varphi + \sigma + \frac{1}{2}\{\varphi, \sigma\} + \dots$, we can then solve the equation $\hat{P}(\varphi \bullet \sigma) = 0$ order by order under the assumption that $\bar{\partial}\sigma = 0$. \square

Under the gauge fixing condition $\hat{P}\varphi = 0$, setting

$$(4.6) \quad \varphi_{s+1} := -\hat{H} \left(\Pi + \sum_{k \geq 0} \frac{ad_{\varphi^s}^k}{(k+1)!} \bar{\partial}\varphi^s \right)_{s+1},$$

where the subscript $s+1$ on the RHS means taking the coefficient of t^{s+1} and $\varphi^s := \varphi_1 + \dots + \varphi_s$, defines $\varphi = \varphi_1 + \varphi_2 + \dots$ inductively.

Remark 4.8. Notice that

$$\bar{\partial} \left(\Pi + \sum_{k \geq 0} \frac{ad_{\varphi^s}^k}{(k+1)!} \bar{\partial}\varphi^s \right)_{s+1} = 0,$$

so the operator \hat{H} , which is defined by integration along paths, is independent of the paths chosen upon applying to these terms.

Remark 4.9. We also observe that the terms φ_s 's vanish on the direct summand $(\Omega^*(U)z^{\hat{m}}) \otimes_{\mathbb{Z}} N$ whenever $\hat{m} \neq km$. Furthermore, we can see that each $\bar{\partial}\varphi_s$ (and all its derivatives) decay exponentially as $O_{s,K}(e^{-c_{s,K}/\hbar})$ on any compact subset $K \subset H_-$ away from P .

We are going to analyze the behavior of φ as $\hbar \rightarrow 0$ to show that it admits an asymptotic expansion with leading order term exactly given by ψ on $\check{X}_0(U) \setminus \check{p}^{-1}(P)$.

4.2.2. *Asymptotic analysis for the gauge φ .* Observe that when we are considering a single wall $\mathbf{w} = (m, P, \Theta)$, the Maurer-Cartan solution Π and hence the gauge φ will be non-trivial only for the summand $(\Omega^*(U)z^{\hat{m}}) \otimes_{\mathbb{Z}} N$ where $\hat{m} = km$ for some $k \in \mathbb{Z}_{>0}$. We use the affine coordinates $u = (u_1, \dots, u_n)$ from Notations 4.1 for each component U for all these summands.

By Remark 4.8, when dealing with closed 1-forms, we can replace the operator $\hat{H}_{\hat{m}}$ by the path integral over any path with the same end points. Let us consider the path ϱ_u defined by

$$\varrho_u = \varrho_{u_1, u_{m,\perp}}(t) = \begin{cases} ((1-2t)u_1^0 + 2tu_1, u_{m,\perp}^0) & \text{if } t \in [0, \frac{1}{2}], \\ (u_1, (2t-1)u_{m,\perp} + (2-2t)u_{m,\perp}^0) & \text{if } t \in [\frac{1}{2}, 1], \end{cases}$$

where $u^0 = q$ (see the left picture of Figure 9). From now on, we will assume that ϱ_u is contained in the contractible open set U by shrinking U if necessary. Then we define the operator \hat{I} by

$$(4.7) \quad \hat{I}(\alpha) =: \int_{\varrho_u} \alpha.$$

By what we just said, we have $\hat{H}_{\hat{m}}(\alpha z^{\hat{m}}) = \hat{I}(\alpha) z^{\hat{m}} = (\int_{\varrho_u} \alpha) z^{\hat{m}}$ for closed 1-forms α .

We are going to apply \hat{I} , instead of \hat{H} , to the closed 1-form

$$\left(\Pi + \sum_{k \geq 0} \frac{ad_{\varphi^s}^k}{(k+1)!} \bar{\partial} \varphi^s \right)_{s+1}$$

to solve for φ_{s+1} because this could somewhat simplify the asymptotic analysis below.

First of all, the first term φ_1 can be explicitly expressed as $\varphi_1 = \sum_k a_{1k} \hat{I}(\delta_m) z^{km} \check{\partial}_n$, where δ_m is the 1-form defined in (4.3) and $\check{\partial}_n$ is the affine vector field defined in (2.6).¹⁰ Since

$$\hat{I}(\delta_m) = \int_{\varrho_u} \delta_m = \left(\frac{1}{\hbar\pi} \right)^{\frac{1}{2}} \int_{\varrho_u} e^{-\frac{u_2^2}{\hbar}} du_2 = \begin{cases} 1 + O_{loc}(\hbar) & \text{if } u \in H_+, \\ O_{loc}(\hbar) & \text{if } u \in H_-, \end{cases}$$

we see that φ_1 has the desired asymptotic expansion, with leading order term given by the coefficient of t^1 in ψ given in (4.2), where the notation $O_{loc}(\hbar)$ means the following:

Notations 4.10. *We say that a function $f(x, \hbar)$ on an open subset $U \times \mathbb{R}_{>0} \subset B_0 \times \mathbb{R}_{>0}$ belongs to $O_{loc}(\hbar^l)$ if it is bounded by $C_K \hbar^l$ for some constant C_K (independent of \hbar) on every compact subset $K \subset U$.*

Next we consider the second term φ_2 . Notice that $[z^{k_1 m} \check{\partial}_n, z^{k_2 m} \check{\partial}_n] = 0$ for all positive k_1, k_2 . Therefore we have

$$(4.8) \quad [\varphi_1, \Pi_1] = - \sum_{k_1, k_2} a_{1k_1} a_{1k_2} \left(\hat{I}(\delta_m) (\nabla_{\partial_n} \delta_m) \check{\partial}_n - \delta_m \nabla_{\partial_n} (\hat{I}(\delta_m)) \check{\partial}_n \right) z^{(k_1+k_2)m},$$

where Π_s refers to the coefficient of t^s in Π and ∂_n was introduced in (2.7).

To compute the order of \hbar in each term in (4.8), we first have $|\hat{I}(\delta_m)| \leq 2$ from the definition of δ_m in (4.3), and using the formula $\delta_m = (\frac{1}{\hbar\pi})^{\frac{1}{2}} (e^{-\frac{u_2^2}{\hbar}} du_2)$, we get

$$|\hat{I}(\hat{I}(\delta_m) \nabla_{\partial_n} \delta_m)| = \frac{\hbar}{4\pi} \left| \int_{\varrho_u} \hat{I}(\delta_m) \nabla_{g_{jk} n^j \frac{\partial}{\partial x_k}} (\delta_m) \right| \leq C \hbar^{1/2} \left| \int_{\varrho_u} (\nabla_{g_{jk} n^j \frac{\partial}{\partial x_k}} e^{-\frac{u_2^2}{\hbar}} du_2) \right| \leq C \hbar^{1/2}.$$

This follows from the fact that $\nabla(u_2)^2$ vanishes along P up to first order, giving an extra $\hbar^{1/2}$ upon integrating against $e^{-\frac{u_2^2}{\hbar}}$. Similarly, we can show that $|\hat{I}(\delta_m \nabla_{\partial_n} (\hat{I}(\delta_m)))| \leq C \hbar^{1/2}$. Therefore we

¹⁰Note that there is a factor $\frac{\hbar}{4\pi}$ in front of the expression of $\check{\partial}_n$ in (2.6), which will become important later when we count the \hbar orders in the asymptotic expansions.

have

$$\varphi_2 = \begin{cases} \sum_{k \geq 1} a_{2k} z^{km} \check{\delta}_n + \sum_{k \geq 1} O_{loc}(\hbar^{1/2}) z^{km} \check{\delta}_n & \text{on } \check{p}^{-1}(H_+), \\ \sum_{k \geq 1} O_{loc}(\hbar^{1/2}) z^{km} \check{\delta}_n & \text{on } \check{p}^{-1}(H_-), \end{cases}$$

where the notation $O_{loc}(\hbar^{1/2}) z^{km} \check{\delta}_n$ means a finite sum of terms of the form $\phi z^{km} \check{\delta}_n$ with $\phi \in O_{loc}(\hbar^{1/2})$.

We would like to argue that the same kind of asymptotic formula holds for a general term φ_s as well. To study the order of \hbar in derivatives of the function $e^{-\frac{u_2^2}{\hbar}}$, we need the following stationary phase approximation (see e.g. [15]).

Lemma 4.11. *Let $U \subset \mathbb{R}^n$ be an open neighborhood of 0 with coordinates x_1, \dots, x_n . Let $\varphi : U \rightarrow \mathbb{R}_{\geq 0}$ be a Morse function with unique minimum $\varphi(0) = 0$ in U . Let $\tilde{x}_1, \dots, \tilde{x}_n$ be a set of Morse coordinates near 0 so that $\varphi(x) = \frac{1}{2}(\tilde{x}_1^2 + \dots + \tilde{x}_n^2)$. For every compact subset $K \subset U$, there exists a constant $C = C_{K,N}$ such that for every $u \in C^\infty(U)$ with $\text{supp}(u) \subset K$, we have*

$$(4.9) \quad \left| \left(\int_K e^{-\varphi(x)/\hbar} u - (2\pi\hbar)^{n/2} \left(\sum_{k=0}^{N-1} \frac{\hbar^k}{2^k k!} \tilde{\Delta}^k \left(\frac{u}{\mathfrak{S}} \right) (0) \right) \right) \right| \leq C \hbar^{n/2+N} \sum_{|\alpha| \leq 2N+n+1} \sup |\partial^\alpha u|,$$

where $\tilde{\Delta} = \sum \frac{\partial^2}{\partial \tilde{x}_j^2}$, $\mathfrak{S} = \pm \det(\frac{d\tilde{x}}{dx})$ and $\mathfrak{S}(0) = (\det \nabla^2 \varphi(0))^{1/2}$. In particular, if u vanishes at 0 up to order L , then we can take $N = \lceil L/2 \rceil$ and get $|\int_K e^{-\varphi(x)/\hbar} u| \leq C \hbar^{n/2+\lceil L/2 \rceil}$.

We will keep track of the order of \hbar in solving the general equation (4.6), and will see that the leading order contribution of φ_{s+1} simply comes from $-H(\Pi_{s+1})$. From the above calculation, we learn that for the 1-form δ_m defined in (4.3), any differentiation $\nabla_{\check{\delta}_n}(\delta_m)$ will contribute an extra vanishing of order $\hbar^{1/2}$, and hence can be considered as an error term. Systematic tracking of these \hbar orders during the iteration (4.6) is necessary. So we extract such properties of δ_m which we need later in the following lemma.

Given a wall $P \subset U$, there is an affine foliation $\{P_q\}_{q \in N}$ of U , where each P_q is a tropical hyperplane parallel to P and N is an affine line transversal to P which parametrizes the leaves, as shown in Figure 5. Given any point $p \in P$ and a neighborhood $V \subset U$ containing p , there is an induced affine foliation $\{(P_{V,q})\}_{q \in N}$ on V .

Lemma 4.12. *Using u_2 as a coordinate for N so that $q = u_2 \in N$ (recall that specific affine coordinates (u_1, \dots, u_n) in U have been chosen in Notations 4.1), and considering the function $g := (u_2)^2$, we have the integral estimate*

$$\int_N u_2^r \left(\sup_{P_{V,u_2}} |\nabla^j(e^{-\frac{g}{\hbar}})| \right) du_2 \leq C_{j,r,V} \hbar^{-\frac{j-r}{2} + \frac{1}{2}}$$

for any $j, r \in \mathbb{Z}_{\geq 0}$.

Proof. First we notice that $\nabla^j(e^{-g/\hbar})$ consists of terms of the form $\hbar^{-M} \left(\prod_{i=1}^M (\nabla^{s_i} g) \right) e^{-g/\hbar}$, where $\sum_i s_i = j$. We see that $\nabla^l \left(\prod_{i=1}^M (\nabla^{s_i} g) \right) |_P \equiv 0$ for $l \leq \sum_{i=1}^M \max(0, 2 - s_i) =: L$. We observe that the terms contributing to the lowest \hbar power are either of the form $\hbar^{-\lfloor \frac{j+1}{2} \rfloor} u_2^r \prod_{i=1}^{\lfloor \frac{j+1}{2} \rfloor} (\nabla^{s_i} g) e^{-g/\hbar}$ having $s_i \leq 2$, or of the form $\hbar^{-j} u_2^r \prod_{i=1}^j (\nabla^{s_i} g) e^{-g/\hbar}$ having $s_i = 1$. In both cases, applying the stationary phase approximation in Lemma 4.11 and counting the vanishing order along P , we obtain $\int_N u_2^r \left(\sup_{P_{V,q}} |\nabla^j(e^{-\frac{g}{\hbar}})| \right) du_2 \leq C_{j,r,V} \hbar^{1/2+(r+j)/2-j} = C_{j,r,V} \hbar^{(r-j)/2+1/2}$. \square

The reader may notice that taking the supremum $\sup_{P_{V,u_2}}$ in Lemma 4.12 is redundant because g is constant along the leaves of the foliation $\{(P_{V,q})\}_{q \in N}$; we write it in this way in order to match Definition 4.19 below.

Remark 4.13. *The order $\hbar^{-\frac{j-r}{2} + \frac{1}{2}}$ which appears in Lemma 4.12 is related to a similar weighted L^2 norm in [30].*

4.2.3. *Differential forms with asymptotic support.* Motivated by the procedure of tracking the \hbar orders as in Lemma 4.12, we would like to formulate the notion of a differential k -form having *asymptotic support on a closed codimension k tropical polyhedral subset $P \subset U$* ; by a tropical polyhedral subset we mean a connected locally convex subset which is locally defined by affine linear equations or inequalities over \mathbb{Q} , as in the codimension 1 case above (Definition 3.3). Before doing so, we first need to define the notion of a differential k -form having exponential decay, or more precisely, having exponential order $O(e^{-c/\hbar})$; the error terms which appear in our later discussion will be of such shape:

Notations 4.14. *We will use the notation $\Omega_{\hbar}^*(B_0)$ (similarly for $\Omega_{\hbar}^*(U)$) to stand for $\Gamma(B_0 \times \mathbb{R}_{>0}, \wedge^* T^* B_0)$, where the extra $\mathbb{R}_{>0}$ direction is parametrized by \hbar .*

Definition 4.15. *We define $\mathcal{W}_k^{-\infty}(U) \subset \Omega_{\hbar}^k(U)$ to be those differential k -forms $\alpha \in \Omega_{\hbar}^k(U)$ such that for each point $q \in U$, there exists a neighborhood V of q where we have $\|\nabla^j \alpha\|_{L^\infty(V)} \leq D_{j,V} e^{-c_V/\hbar}$ for some constants c_V and $D_{j,V}$. The association $U \mapsto \mathcal{W}_k^{-\infty}(U)$ defines a sheaf over B_0 which is denoted by $\mathcal{W}_k^{-\infty}$.*

We will also consider differential forms which only blow up at polynomial orders in \hbar^{-1} :

Definition 4.16. *We define $\mathcal{W}_k^\infty(U) \subset \Omega_{\hbar}^k(U)$ to be those differential k -forms $\alpha \in \Omega_{\hbar}^k(U)$ such that for each point $q \in U$, there exists a neighborhood V of q where we have $\|\nabla^j \alpha\|_{L^\infty(V)} \leq D_{j,V} \hbar^{-N_{j,V}}$ for some constant $D_{j,V}$ and $N_{j,V} \in \mathbb{Z}_{>0}$. The association $U \mapsto \mathcal{W}_k^\infty(U)$ defines a sheaf over B_0 which is denoted by \mathcal{W}_k^∞ .*

Notice that the sheaves $\mathcal{W}_k^{\pm\infty}$ in Definitions 4.15 and 4.16 are closed under application of $\nabla_{\frac{\partial}{\partial x}}$, the deRham differential d and wedge product of differential forms. We also observe the fact that $\mathcal{W}_k^{-\infty}$ is a differential graded ideal of \mathcal{W}_k^∞ ; this will be useful later in Section 5.2. In particular, we can consider the sheaf of differential graded algebras $\mathcal{W}_*^\infty/\mathcal{W}_*^{-\infty}$, equipped with the deRham differential.

The following lemma will be useful in Section 5.3 (readers may skip it until Section 5.3):

Lemma 4.17. (1) *Suppose that $\alpha \in (\mathcal{W}_k^\infty/\mathcal{W}_k^{-\infty})(U) = \mathcal{W}_k^\infty(U)/\mathcal{W}_k^{-\infty}(U)$ (note that the sheaves $\mathcal{W}_*^{\pm\infty}$ are both soft sheaves) satisfies $d\alpha = 0$ in $(\mathcal{W}_{k+1}^\infty/\mathcal{W}_{k+1}^{-\infty})(U)$. Then for any compact family of smooth $(k+1)$ -chain $\{\gamma_u\}_{u \in K}$ in U , we have*

$$\left| \int_{\partial\gamma_u} \hat{\alpha} \right| \leq D_{K,\hat{\alpha}} e^{-c_{K,\hat{\alpha}}/\hbar}$$

for any $u \in K$, where $\hat{\alpha}$ is any choice of lifting of α to $\mathcal{W}_k^\infty(U)$.

(2) *For any $\alpha \in (\mathcal{W}_1^\infty/\mathcal{W}_1^{-\infty})(U)$ and a fixed based point $x^0 \in U$, the path integral $f_\alpha := \int_{x^0}^x \hat{\alpha}$, defined locally by first choosing a contractible compact subset $K \subset U$, then a family of paths $\varrho : [0, 1] \times K \rightarrow U$ joining x^0 to $x \in K$, and also a lifting $\hat{\alpha}$ of α to $\mathcal{W}_1^\infty(U)$, gives a well-defined element in $(\mathcal{W}_0^\infty/\mathcal{W}_0^{-\infty})(U)$, meaning that for different choices of K , ϱ and $\hat{\alpha}$, the path integrals only differ by elements in $\mathcal{W}_0^{-\infty}(U)$.*

Proof. For the first statement, the equation $d\alpha = 0$ in $(\mathcal{W}_k^\infty/\mathcal{W}_k^{-\infty})(U)$ means that we have $d\hat{\alpha} = \beta$ for some $\beta \in \mathcal{W}_{k+1}^{-\infty}(U)$. Stokes' Theorem then implies that $\int_{\partial\gamma_u} \hat{\alpha} = \int_{\gamma_u} \beta = O_{K,\beta}(e^{-c_{K,\beta}/\hbar})$.

For the second statement, we first fix a point $x \in U$ and a contractible compact subset $K \subset U$ such that $x \in \text{int}(K)$, and also a lifting $\hat{\alpha}$ with $d\hat{\alpha} = \beta \in \mathcal{W}_2^{-\infty}(U)$. Suppose that we have two families of paths $\varrho_1, \varrho_2 : [0, 1] \times K \rightarrow U$ parametrized by K . Using contractibility of U , we have a homotopy $h : [0, 1]^2 \times K \rightarrow U$ between ϱ_1 and ϱ_2 satisfying $h(0, \cdot) = \varrho_0$, $h(1, \cdot) = \varrho_1$, $h(\cdot, 0) = x^0$ and $h(\cdot, 1) = \text{id}_K$. Therefore we have $\varrho_1 - \varrho_0 = \partial h$, and hence the difference of the two path integrals is given by

$$\int_{\varrho_1(\cdot, x)} \hat{\alpha} - \int_{\varrho_0(\cdot, x)} \hat{\alpha} = \int_{h(\cdot, x)} d\hat{\alpha} = \int_{h(\cdot, x)} \beta = \int_{[0, 1]^2} h^*(\beta).$$

Taking the covariant derivatives by ∇^j of this difference, we have $\nabla^j \left(\int_{\varrho_1(\cdot, x)} \hat{\alpha} - \int_{\varrho_0(\cdot, x)} \hat{\alpha} \right) = \int_{[0, 1]^2} \nabla^j(h^*(\beta))$. From the fact that $\beta \in \mathcal{W}_2^{-\infty}(U)$, we have $|\nabla^j(h^*(\beta))|(s, t, u) \leq D_{j, K, h} e^{-c_{j, K, h}/\hbar}$ for any point $(s, t, u) \in [0, 1]^2 \times K$, and therefore $\|\nabla^j(f_{1, \hat{\alpha}} - f_{2, \hat{\alpha}})\|_{L^\infty(K)} \leq D_{j, K, h} e^{-c_{j, K, h}/\hbar}$. Hence f_α , as an element of $(\mathcal{W}_0^\infty/\mathcal{W}_0^{-\infty})(U)$, is independent of the choice of the family of paths ϱ .

Now if K_1, K_2 are two contractible compact subsets with $x \in \text{int}(K_1 \cap K_2)$, we can change the family of paths parametrized by each K_i to an auxiliary one parametrized by $K_1 \cap K_2$. By above, the path integral will only differ by elements in $\mathcal{W}_0^{-\infty}(U)$. Finally, for two different liftings $\hat{\alpha}_1, \hat{\alpha}_2$ of α , we have $\hat{\alpha}_1 - \hat{\alpha}_2 \in \mathcal{W}_1^{-\infty}(U)$ and so $\int_{x_0}^x \hat{\alpha}_1 - \hat{\alpha}_2 \in \mathcal{W}_0^{-\infty}(U)$. This completes the proof of the second statement. \square

Notations 4.18. Let $P \subset U$ be a closed codimension k tropical polyhedral subset.

- (1) There is a natural foliation $\{P_q\}_{q \in N}$ in U obtained by parallel transporting the tangent space of P (at some interior point in P) to every point in U by the affine connection ∇ on B_0 .

We let $\nu_P \in \Gamma(U, \wedge^k(N_P^*))$ be a top covariant constant form (i.e. $\nabla(\nu_P) = 0$) in the conormal bundle N_P^* of P (which is unique up to scaling by constants); we regard ν_P as a volume form on space of leaves N if it admits a smooth structure. We also let $\nu_P^\vee \in \wedge^k N_P$ be a volume element dual to ν_P , and choose a lifting of ν_P^\vee as an element in $\wedge^k TU$ (which will again be denoted by ν_P^\vee by abusing notations).




FIGURE 5. The foliation near P

- (2) For any point $p \in P$, we choose a sufficiently small convex neighborhood $V \subset U$ containing p so that there exists a slice $N_V \subset V$ transversal to the foliation $\{P_q \cap V\}$ given by intersection of $\{P_q\}_{q \in N}$ with V , i.e. a dimension k affine subspace which is transversal to all the leaves in $\{P_q \cap V\}$; we denote this foliation on V by $\{(P_{V, q})\}_{q \in N_V}$, using N_V as the parameter space. See Figure 5 for an illustration.

In V , we take local affine coordinates $x = (x_1, \dots, x_n)$ such that $x' := (x_1, \dots, x_k)$ parametrizes N_V with $x' = 0$ corresponding to the unique leaf containing P . Using these coordinates, we can write $\nu_P = dx_1 \wedge \dots \wedge dx_k$ and $\nu_P^\vee = \frac{\partial}{\partial x_1} \wedge \dots \wedge \frac{\partial}{\partial x_k}$.

Definition 4.19. A differential k -form $\alpha \in \mathcal{W}_k^\infty(U)$ is said to have asymptotic support on a closed codimension k tropical polyhedral subset $P \subset U$ if the following conditions are satisfied:

- (1) For any $p \in U \setminus P$, there is a neighborhood $V \subset U \setminus P$ of p such that $\alpha|_V \in \mathcal{W}_k^{-\infty}(V)$ on V .
- (2) There exists a neighborhood W_P of P in U such that we can write

$$\alpha = h(x, \hbar)\nu_P + \eta,$$

where ν_P is the volume form Notations 4.18(1), $h(x, \hbar) \in C^\infty(W_P \times \mathbb{R}_{>0})$ and η is an error term satisfying $\eta \in \mathcal{W}_k^{-\infty}(W_P)$ on W_P (see Figure 4).

- (3) For any $p \in P$, there exists a sufficiently small convex neighborhood V containing p such that using the coordinate system chosen in Notations 4.18(2) and considering the foliation $\{(P_{V,x'})\}_{x' \in N_V}$ in V , we have, for all $j \in \mathbb{Z}_{\geq 0}$ and multi-index $\beta = (\beta_1, \dots, \beta_k) \in \mathbb{Z}_{\geq 0}^k$, the estimate

$$(4.10) \quad \int_{x' \in N_V} (x')^\beta \left(\sup_{P_{V,x'}} |\nabla^j(\iota_{\nu_P} \alpha)| \right) \nu_P \leq D_{j,V,\beta} \hbar^{-\frac{j+s-|\beta|-k}{2}},$$

for some constant $D_{j,V,\beta}$ and some $s \in \mathbb{Z}$, where $|\beta| = \sum_l \beta_l$ is the vanishing order of the monomial $(x')^\beta = x_1^{\beta_1} \cdots x_k^{\beta_k}$ along $P_{x'=0}$.

Remark 4.20. Note that condition (3) in Definition 4.19 is independent of the choice of the convex neighborhood V , the transversal slice N_V and the choice of the local affine coordinates $x = (x_1, \dots, x_n)$ (although the constant $D_{j,V,\beta}$ may depends these choices). Therefore this condition can be checked by choosing a sufficiently nice neighborhood V at every point $p \in P$.

Remark 4.21. The idea of putting the weight $(x')^\beta$ and the differentiation ∇^j in condition (3) in Definition 4.19 comes from a similar weighted L^2 norm used in [30]. In this paper, instead of L^2 norms, we use a mixture of L^∞ and L^1 norms for the purpose of Lemma 4.22.

The estimate in condition (3) of Definition 4.19 defines the following filtration


$$(4.11) \quad \mathcal{W}_k^{-\infty} \cdots \subset \mathcal{W}_P^{-s} \subset \cdots \subset \mathcal{W}_P^{-1} \subset \mathcal{W}_P^0 \subset \mathcal{W}_P^1 \subset \mathcal{W}_P^2 \subset \cdots \subset \mathcal{W}_P^s \subset \cdots \subset \mathcal{W}_k^\infty \subset \Omega_\hbar^k(U),$$

where, for any given $s \in \mathbb{Z}$, $\mathcal{W}_P^s = \mathcal{W}_P^s(U)$ denotes the set of k -forms $\alpha \in \mathcal{W}_k^\infty(U)$ with asymptotic support on P such that the estimate (4.10) holds with the given integer s . Note that the degree k of the differential forms has to be equal to the codimension of P . Also note that the sets $\mathcal{W}_k^{\pm\infty}(U)$ are independent of the choice of P . This filtration keeps track of the polynomial order of \hbar for k -forms with asymptotic support on P , and it provides a convenient tool for us to prove and express our results in the subsequent asymptotic analysis. In these terms, Lemma 4.12 simply means $\delta_m \in \mathcal{W}_P^1(U)$, where P is the tropical hyperplane supporting a wall.

The filtration satisfies $\nabla_{\frac{\partial}{\partial x_l}} \mathcal{W}_P^s(U) \subset \mathcal{W}_P^{s+1}(U)$ for any $l = 1, \dots, n$, and $(x')^\beta \mathcal{W}_P^s(U) \subset \mathcal{W}_P^{s-|\beta|}(U)$ for any affine monomial $(x')^\beta$ with vanishing order $|\beta|$ along P , so we have the nice property that

$$(4.12) \quad (x')^\beta \nabla_{\frac{\partial}{\partial x_{l_1}}} \cdots \nabla_{\frac{\partial}{\partial x_{l_j}}} \mathcal{W}_P^s(U) \subset \mathcal{W}_P^{s+j-|\beta|}(U).$$

Lemma 4.22. For two closed tropical polyhedral subsets $P_1, P_2 \subset U$ of codimension k_1, k_2 respectively, we have $\mathcal{W}_{P_1}^s(U) \wedge \mathcal{W}_{P_2}^r(U) \subset \mathcal{W}_P^{r+s}(U)$ for any codimension $k_1 + k_2$ polyhedral subset P containing $P_1 \cap P_2$ normal to $\nu_{P_1} \wedge \nu_{P_2}$ if they intersect transversally,¹¹ and $\mathcal{W}_{P_1}^s(U) \wedge \mathcal{W}_{P_2}^r(U) \subset \mathcal{W}_{k_1+k_2}^{-\infty}(U)$ if their intersection is not transversal.

FIGURE 6. Foliation in the neighborhood V

Before giving the proof, let us clarify that, when we say two closed tropical polyhedral subsets $P_1, P_2 \subset U$ of codimension k_1, k_2 are *intersecting transversally*, we mean the affine subspaces containing P_1, P_2 and of codimension k_1, k_2 respectively are intersecting transversally; this definition also applies to the case when $\partial P_i \neq \emptyset$, as shown in Figure 6.

Proof of Lemma 4.22. We first consider the case when P_1 and P_2 are not intersecting transversally. Part (2) of Definition 4.19 says that we have neighborhoods W_{P_i} of P_i such that we can write $\alpha_i = h_i \nu_{P_i} + \eta_i$ for $i = 1, 2$. Since $\nu_{P_1} \wedge \nu_{P_2} = 0$ in $W_{P_1} \cap W_{P_2}$ by the non-transversal assumption, we have $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_k^{-\infty}(W_{P_1} \cap W_{P_2})$ near $P_1 \cap P_2$, and hence $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_k^{-\infty}(U)$ by condition (1) in Definition 4.19 and the fact that $\mathcal{W}_k^{-\infty}(U)$ is a differential ideal of $\mathcal{W}_k^{\infty}(U)$.

Next we assume that $P_1 \pitchfork P_2 = Q$. Let $\alpha_1 \in \mathcal{W}_{P_1}^s(U)$ and $\alpha_2 \in \mathcal{W}_{P_2}^r(U)$. Using again the fact that $\mathcal{W}_k^{-\infty}(U)$ is a differential ideal of $\mathcal{W}_k^{\infty}(U)$, same reasoning as above shows that condition (1) in Definition 4.19 holds for $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_Q^{r+s}(U)$. Condition (2) in Definition 4.19 is also satisfied because in this case we have $\nu_Q = \nu_{P_1} \wedge \nu_{P_2}$ in $W_Q = W_{P_1} \cap W_{P_2}$. So it remains to prove condition (3) in Definition 4.19.

Fixing a point $p \in Q$, we take an affine convex coordinate chart given by $V(\subset T_p U \cong M_{\mathbb{R}}) \rightarrow U$ centered at $0 \in T_p U$. Then $V_Q := V \cap T_p Q$ is a neighborhood of 0 in $T_p Q$. We take \mathbb{Q} -affine bases $m_2^1, \dots, m_2^{k_2}$ of $T_p P_1 / T_p Q$ and $m_1^1, \dots, m_1^{k_1}$ of $T_p P_2 / T_p Q$ respectively, and the corresponding dual bases in $(T_p U / T_p P_2)^*$ and $(T_p U / T_p P_1)^*$. We use $x^i \cdot m_i = \sum_{j=1}^{k_i} x_j^i m_j^i$, $i = 1, 2$ to stand for the natural pairing between $x^1 = (x_1^1, \dots, x_{k_1}^1) \in (T_p U / T_p P_1)^*$ and $m_1^1, \dots, m_1^{k_1}$, and that between $x^2 = (x_1^2, \dots, x_{k_2}^2) \in (T_p U / T_p P_2)^*$ and $m_2^1, \dots, m_2^{k_2}$, respectively. By shrinking V if necessary, we can write it as $V = \bigcup_{\substack{x^1 \in (-\delta, \delta)^{k_1} \\ x^2 \in (-\delta, \delta)^{k_2}}} (x^1 \cdot m_1 + x^2 \cdot m_2 + V_Q)$ for some small $\delta > 0$,

as shown in Figure 6. Then we can parametrize the foliations induced by Q , P_1 and P_2 respectively as $Q_{V, (x^1, x^2)} = x^1 \cdot m_1 + x^2 \cdot m_2 + V_Q$, $(P_1)_{V, x^1} = x^1 \cdot m_1 + \bigcup_{x^2 \in (-\delta, \delta)^{k_2}} (x^2 \cdot m_2 + V_Q)$ and $(P_2)_{V, x^2} = x^2 \cdot m_2 + \bigcup_{x^1 \in (-\delta, \delta)^{k_1}} (x^1 \cdot m_1 + V_Q)$. We also extend (x^1, x^2) to local affine coordinates $(x_1^1, \dots, x_{k_1}^1, x_1^2, \dots, x_{k_2}^2, x_{k_1+k_2+1}, \dots, x_n)$ on V .

Now for $\alpha_1 \in \mathcal{W}_{P_1}^r(U)$ and $\alpha_2 \in \mathcal{W}_{P_2}^s(U)$, we first observe that we can write $\alpha_i = h_i(x, \hbar) dx^i + \eta_i = h_i(x, \hbar) dx_1^i \wedge \dots \wedge dx_{k_i}^i + \eta_i$ for $i = 1, 2$, and we have $\nabla^j(h_1 h_2) = \sum_{j_1+j_2=j} (\nabla^{j_1} h_1)(\nabla^{j_2} h_2)$. Also, any affine monomial $(x')^\beta$ (in the coordinates $(x_1^1, \dots, x_{k_1}^1, x_1^2, \dots, x_{k_2}^2)$) with vanishing order $|\beta|$ along Q can be rewritten in the form $(x^1)^{\beta_1} (x^2)^{\beta_2}$, where $(x^i)^{\beta_i}$ has vanishing order $|\beta_i|$ along Q .

¹¹In particular we can take $P = P_1 \cap P_2$ if $\text{codim}_{\mathbb{R}}(P_1 \cap P_2) = k_1 + k_2$.

Since the error terms η_i 's are not contributing when we count the polynomial order in \hbar^{-1} , it remains to estimate a term of the form $(x^1)^{\beta_1}(x^2)^{\beta_2}(\nabla^{j_1} h_1)(\nabla^{j_2} h_2)$. We have

$$\begin{aligned}
& \int_{(x^1, x^2) \in (-\delta, \delta)^{k_1+k_2}} (x^1)^{\beta_1}(x^2)^{\beta_2} \sup_{Q_{V, (x^1, x^2)}} |(\nabla^{j_1} h_1)(\nabla^{j_2} h_2)| dx^1 dx^2 \\
&= \int_{x^2 \in (-\delta, \delta)^{k_2}} (x^2)^{\beta_2} \left(\int_{x^1 \in (-\delta, \delta)^{k_1}} (x^1)^{\beta_1} \sup_{Q_{V, (x^1, x^2)}} |(\nabla^{j_1} h_1)(\nabla^{j_2} h_2)| dx^1 \right) dx^2 \\
&\leq \int_{x^2} (x^2)^{\beta_2} \sup_{(P_2)_{V, x^2}} |(\nabla^{j_2} h_2)| \left(\int_{x^1} (x^1)^{\beta_1} \sup_{Q_{V, (x^1, x^2)}} |(\nabla^{j_1} h_1)| dx^1 \right) dx^2 \\
&\leq D_{j_1, V, (x^1)^{\beta_1}} \hbar^{-\frac{j_1+s-|\beta_1|-k_1}{2}} \int_{x^2} (x^2)^{\beta_2} \sup_{(P_2)_{V, x^2}} |(\nabla^{j_2} h_2)| dx^2 \\
&\leq D_{j_1, V, (x^1)^{\beta_1}} D_{j_2, V, (x^2)^{\beta_2}} \hbar^{-\frac{j_1+s-|\beta_1|-k_1}{2}} \hbar^{-\frac{j_2+r-|\beta_2|-k_2}{2}} \leq D_{j, V, x^\beta} \hbar^{-\frac{j+s+r-|\beta|-k}{2}},
\end{aligned}$$

which gives the desired estimate in condition (3) of Definition 4.19. \square

For a given closed tropical polyhedral subset $P \subset U$, we choose a reference tropical hyperplane $R \subset U$ which divides the base U as $U \setminus R = U_+ \cup U_-$ such that $P \subset U_+$, together with an affine vector field v (meaning $\nabla v = 0$) not tangent to R pointing into U_+ . We let

$$(4.13) \quad I(P) := (P + \mathbb{R}_{\geq 0} v) \cap U$$

be the image swept out by P under the flow of v .

By shrinking U if necessary, we can assume that for any point $p \in U$, the unique flow line of v in U passing through p intersects R uniquely at a point $x \in R$. Then the time- t flow along v defines a diffeomorphism $\tau : W \rightarrow U$, $(t, x) \mapsto \tau(t, x)$, where $W \subset \mathbb{R} \times R$ is the maximal domain of definition of τ (namely, for any $x \in R$, there is a maximal time interval $I_x \subset \mathbb{R}$ so that the flow line through x has its image lying inside U). For any point $x \in R$, we denote by $\tau_x(t) := \tau(t, x)$ the flow line of v passing through x . Figure 7 illustrates the situation.




FIGURE 7. The flow along v and $I(P)$

We now define an integral operator I as

$$(4.14) \quad I(\alpha)(t, x) := \int_0^t \iota_{\frac{\partial}{\partial s}}(\tau^*(\alpha))(s, x) ds.$$

Note that I depends on the choice of the tropical hyperplane R .

Lemma 4.23. *For $\alpha \in \mathcal{W}_P^s(U)$, we have $I(\alpha) \in \mathcal{W}_{k-1}^{-\infty}(U)$ if v is tangent to P , and $I(\alpha) \in \mathcal{W}_{I(P)}^{s-1}(U)$ if v is not tangent to P , where $I(P)$ is defined in (4.13).*

Proof. In order to simplify notations in this proof, we will omit τ^* in the definition (4.14) of I by treating $\tau : W \rightarrow U$ as an affine coordinate chart.

Suppose that v is tangent to P . By condition (2) of Definition 4.19, we have a neighborhood $W_P \subset U$ such that $\alpha = h\nu_P + \eta$. For each point $x \in R$, the path $\tau_x(t)$ is tangent to the foliation $\{P_q\}_{q \in N}$ in W_P whenever $\tau_x(t) \in W_P$ by the tangency assumption. This means $\iota_{\frac{\partial}{\partial t}}(\nu_P) = 0$ in $\tau_x^{-1}(W_P)$ and hence we have

$$I(\alpha)(t, x) = \int_{[0,t]} \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds = \int_{[0,t] \cap \tau_x^{-1}(U \setminus W_P)} \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds + \int_{[0,t] \cap \tau_x^{-1}(W_P)} \iota_{\frac{\partial}{\partial s}} \eta(s, x) ds.$$

So we have $I(\alpha) \in \mathcal{W}_{k-1}^{-\infty}(U)$ by conditions (1) and (2) of Definition 4.19.

Now suppose that v is not tangent to P . Let $I(W_P) := \bigcup_{t \geq 0} (W_P + t \cdot v) \cap U$, which gives an open neighborhood of $I(P)$. Concerning condition (1) in Definition 4.19, we take $\tau(t_0, x_0) \in U \setminus I(P)$, and then a neighborhood V of $\tau(t_0, x_0)$ in $U \setminus I(P)$ and a neighborhood $W'_P \subset W_P$ of P , such that, for any point $\tau(t, x) \in V$, the flow line joining $\tau(t, x)$ to R does not hit $\overline{W'_P}$. This implies that $I(\alpha)|_V \in \mathcal{W}_{k-1}^{-\infty}(V)$ since we have $\alpha|_{U \setminus \overline{W'_P}} \in \mathcal{W}_k^{-\infty}(U \setminus \overline{W'_P})$ and

$$I(\alpha)(t, x) = \int_0^t \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds = \int_{[0,t] \cap \tau_x^{-1}(U \setminus \overline{W'_P})} \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds.$$

So condition (1) in Definition 4.19 holds for $I(\alpha)$.

Concerning condition (2) in Definition 4.19, we first note that $v = \frac{\partial}{\partial t}$ is tangent to $I(P)$, so by parallel transporting the form $\iota_{\frac{\partial}{\partial t}} \nu_P$ to the neighborhood $I(W_P)$, we obtain a volume element in the normal bundle of $I(P)$, which we denote by $\nu_{I(P)}$. For a point $q \in I(W_P)$, we take a small neighborhood V near q , and for $\tau(t, x) \in V$, we write


$$\begin{aligned} I(\alpha)(t, x) &= \int_0^t \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds = \int_{[0,t] \cap \tau_x^{-1}(U \setminus W_P)} \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds + \int_{[0,t] \cap \tau_x^{-1}(W_P)} \iota_{\frac{\partial}{\partial s}} (h\nu_P + \eta)(s, x) ds \\ &= \left(\int_{[0,t] \cap \tau_x^{-1}(W_P)} h(s, x) ds \right) \nu_{I(P)} + \int_{[0,t] \cap \tau_x^{-1}(W_P)} \iota_{\frac{\partial}{\partial s}} \eta(s, x) ds + \int_{[0,t] \cap \tau_x^{-1}(U \setminus W_P)} \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds, \end{aligned}$$

where the last two terms are in $\mathcal{W}_{k-1}^{-\infty}(V)$, and condition (2) in Definition 4.19 holds for $I(\alpha)$.

Concerning condition (3) in Definition 4.19, we fix a point $p = \tau(b, x) \in I(P)$ and let $p' = \tau(a, x) \in P$ be the unique point such that p, p' lie on the same flow line τ_x . We take local affine coordinates $x = (x_1, \dots, x_{k-1}, x_k, \dots, x_{n-1}) \in (-\delta, \delta)^{n-1}$ of R centered at p' (meaning that $p' = (a, 0)$) such that $x' = (x_1, \dots, x_{k-1})$ are normal to the tropical polyhedral subset $p_R(\tau^{-1}(P)) \subset R$, where $p_R : W(\subset \mathbb{R} \times R) \rightarrow R$ is the natural projection.

By taking δ small enough, we have $\tau : (a - \delta, b + \delta) \times (-\delta, \delta)^{n-1} \rightarrow U$ mapping diffeomorphically onto its image, such that it contains the part the flow line $\tau_{0|[a,b]}$ joining p' to p . We can also take $V = \tau((b - \delta, b + \delta) \times (-\delta, \delta)^{n-1})$ with $\tau(b, 0) = p$, and arrange that $V' = \tau((a - \delta, a + \delta) \times (-\delta, \delta)^{n-1}) \subset W_P$ with $\tau(a, 0) = p'$. Notice that there is a possibility that $p = p' \in P$ and therefore $a = b$ in the above description which means $V = V'$. Figure 8 illustrates the situation.

Recall that there is a foliation $\{P_q\}_{q \in N}$ codimension k affine subspaces parallel to P . Then the induced foliation $\{P_{t,x'}\}_{(t,x') \in N_{V'}}$ of the neighborhood V' can be parametrized by $N_{V'} := (a - \delta, a + \delta) \times (-\delta, \delta)^{k-1}$. Therefore the foliation of V induced by $I(P)$ is parametrized as $\{I(P)_{x'}\}_{x' \in N_V}$, where $I(P)_{x'} = \bigcup_{t \in (b - \delta, b + \delta)} (P_{0,x'} + tv)$ and $N_V = (-\delta, \delta)^{k-1}$.

FIGURE 8. Neighborhood along the flow line $\tau_0(t)$

For $\alpha \in \mathcal{W}_P^s$, we consider $I(\alpha) = \int_0^t \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds$ in the neighborhood V , and what we need to estimate is the term $\int_{N_V} (x')^\beta \sup_{I(P)_{x'}} |\nabla^j \iota_{\nu_{I(P)}} \alpha| \nu_{I(P)}$. The integral $I(\alpha)$ can be split into two parts as $\int_0^t = \int_0^{a-\delta} + \int_{a-\delta}^t$ and we only have to control the second part $I_{a-\delta}(\alpha) := \int_{a-\delta}^t \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds$ because $\alpha \in \mathcal{W}_P^s(U)$ satisfies condition (1) in Definition 4.19, so $\int_0^{a-\delta} \iota_{\frac{\partial}{\partial s}} \alpha(s, x) ds$, as a function of (t, x) which is constant in t , lies in $\mathcal{W}_{k-1}^{-\infty}(U)$ (as the integral misses the support P of α). Writing $\nabla^j = \nabla_\perp^{j_1} \nabla_{\frac{\partial}{\partial t}}^{j_2}$, where $\nabla_\perp(t) = 0$, we have two cases depending on whether $j_2 = 0$ or $j_2 > 0$:

Case 1: $j_2 = 0$. Then we have

$$|\nabla_\perp^j \iota_{\nu_{I(P)}} (I_{a-\delta}(\alpha))| \leq \int_{a-\delta}^{a+\delta} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds + \int_{a+\delta}^{b+\delta} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds.$$

The latter term can be dropped because the domain $\int_{a+\delta}^{b+\delta}$ misses the support of P , so it lies in $\mathcal{W}_{k-1}^{-\infty}$. For the first term, we treat $\int_{a-\delta}^{a+\delta} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds$ as a function of (t, x) on V which is constant along the t -direction. Therefore we estimate

$$\begin{aligned} & \int_{x'} (x')^\beta \sup_{I(P)_{x'}} \left(\int_{a-\delta}^{a+\delta} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds \right) \nu_{I(P)} = \int_{x'} (x')^\beta \sup_{P_{0,x'}+bv} \left(\int_{a-\delta}^{a+\delta} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds \right) \nu_{I(P)} \\ & \leq \int_{x'} \sup_{P_{0,x'}+bv} \left(\int_{a-\delta}^{a+\delta} (x')^\beta \sup_{P_{s,x'}} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds \right) \nu_{I(P)} = \int_{x'} \left(\int_{a-\delta}^{a+\delta} (x')^\beta \sup_{P_{s,x'}} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds \right) \nu_{I(P)} \\ & \leq C_{j,V',\beta} \hbar^{-\frac{j+s-|\beta|-k}{2}}, \end{aligned}$$

where the first inequality follows from the inequality

$$\int_{a-\delta}^{a+\delta} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds \leq \int_{a-\delta}^{a+\delta} \sup_{P_{t,x'}} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds,$$

and the second equality is due to the fact that $\int_{a-\delta}^{a+\delta} \sup_{P_{t,x'}} |\nabla_\perp^j (\iota_{\nu_P} \alpha)| ds$, treated as function on V , is constant along the leaf $P_{0,x'} + bv$. Writing $j + s - |\beta| - k = j + (s-1) - |\beta| - (k-1)$, we obtain the desired estimate so that $\alpha \in \mathcal{W}_{I(P)}^{s-1}(U)$.

Case 2: $j_2 > 0$. Then we have $\nabla_{\frac{\partial}{\partial t}}^{j_2} \iota_{\nu_{I(P)}^\vee} (I_{a-\delta}(\alpha)) = \nabla_{\frac{\partial}{\partial t}}^{j_2-1} (\iota_{\nu_P^\vee} \alpha)$. We can rewrite it as

$$\nabla_{\perp}^{j_1} \nabla_{\frac{\partial}{\partial t}}^{j_2} \iota_{\nu_{I(P)}^\vee} (I_{a-\delta}(\alpha))(t, x) = \int_{a-\delta}^t \nabla^j (\iota_{\nu_P^\vee} \alpha)(s, x) ds + \left(\nabla_{\frac{\partial}{\partial t}}^{j_2-1} \nabla_{\perp}^{j_1} (\iota_{\nu_P^\vee} \alpha) \right) (a - \delta, x),$$

where the latter term lies in $\mathcal{W}_{k-1}^{-\infty}$ because it misses the support P of α , and the first term is bounded by

$$\left| \int_{a-\delta}^t \nabla^j (\iota_{\nu_P^\vee} \alpha)(s, x) ds \right| \leq \int_{a-\delta}^{a+\delta} |\nabla^j (\iota_{\nu_P^\vee} \alpha)|(s, x) ds + \int_{a+\delta}^{b+\delta} |\nabla^j (\iota_{\nu_P^\vee} \alpha)|(s, x) ds.$$

The same argument as Case 1 can then be applied to get the desired estimate. \square

Remark 4.24. *Lemmas 4.22 and 4.23 say that we can relate the differential-geometric operations \wedge and I to intersection and suspension of asymptotic supports. These properties are essential for relating Maurer-Cartan solutions, which are differential-geometric in nature, to combinatorics of scattering diagrams.*

In order to apply the notion of asymptotic support to keep track of the \hbar order in asymptotic expansions of the gauge element $\varphi^s = \varphi_1 + \varphi_2 + \dots + \varphi_s$, we will restrict our attention to the dg Lie subalgebra $(\bigoplus_m \mathcal{W}_*^\infty(U)z^m) \otimes_{\mathbb{Z}} N \subset \mathbf{G}^*(U)$, whose elements are finite sums of the form $\sum_{m,n} \alpha_m^n z^m \check{\partial}_n$, where $\alpha_m^n \in \mathcal{W}_*^\infty(U)$. Restriction of \hat{H} defined in Definition 4.5 to $(\bigoplus_m \mathcal{W}_*^\infty(U)z^m) \otimes_{\mathbb{Z}} N$ gives the homotopy operator $\hat{H} : (\bigoplus_m \mathcal{W}_*^\infty(U)z^m) \otimes_{\mathbb{Z}} N \rightarrow (\bigoplus_m \mathcal{W}_{*-1}^\infty(U)z^m) \otimes_{\mathbb{Z}} N$ defined as $\hat{H} \left(\sum_{m,n} \alpha_m^n z^m \check{\partial}_n \right) = \sum_{m,n} \int_0^1 \rho_q^*(\alpha_m^n) z^m \check{\partial}_n$, using ρ_q in Definition 4.5. We also write $\hat{I} \left(\sum_{m,n} \alpha_m^n z^m \check{\partial}_n \right) = \sum_{m,n} \hat{I}(\alpha_m^n) z^m \check{\partial}_n$ when the α_m^n 's are 1-forms. Extending Lemma 4.22 to this dg Lie subalgebra, we have the following:

Lemma 4.25. *Given $m_1, m_2 \in M$, $n_1, n_2 \in N$, and $\alpha_1 \in \mathcal{W}_{P_1}^s(U)$ and $\alpha_2 \in \mathcal{W}_{P_2}^r(U)$. If P_1 of codimension k_1 intersects P_2 of codimension k_2 transversally, then we have*

$$[\alpha_1 z^{m_1} \check{\partial}_{n_1}, \alpha_2 z^{m_2} \check{\partial}_{n_2}] \in \alpha_1 \wedge \alpha_2 z^{m_1+m_2} \check{\partial}_{(m_2, n_1)n_2 - (m_1, n_2)n_1} + \mathcal{W}_P^{r+s-1}(U) z^{m_1+m_2} \otimes_{\mathbb{Z}} N$$

and $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_P^{r+s}(U)$ for any codimension $k_1 + k_2$ polyhedral subset P containing $P_1 \cap P_2$ normal to $\nu_{P_1} \wedge \nu_{P_2}$. If the intersection is not transversal, then we have $[\alpha_1 z^{m_1} \check{\partial}_{n_1}, \alpha_2 z^{m_2} \check{\partial}_{n_2}] \in \mathcal{W}_k^{-\infty}(U) z^{m_1+m_2} \otimes_{\mathbb{Z}} N$.

Proof. From the definition of the Lie bracket we have

$$\begin{aligned} [\alpha_1 z^{m_1} \check{\partial}_{n_1}, \alpha_2 z^{m_2} \check{\partial}_{n_2}] &= \alpha_1 \wedge \alpha_2 z^{m_1+m_2} \check{\partial}_{(m_2, n_1)n_2 - (m_1, n_2)n_1} \\ &\quad + \alpha_1 \wedge \nabla_{\partial_1}(\alpha_2) z^{m_1+m_2} \check{\partial}_{n_2} \pm \alpha_2 \wedge \nabla_{\partial_2}(\alpha_1) z^{m_1+m_2} \check{\partial}_{n_1} \end{aligned}$$

When P_1 and P_2 are intersecting transversally and let P as above, Lemma 4.22 says that $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_P^{r+s}(U)$, so it remains to show that the last two terms are lying in $\mathcal{W}_P^{r+s-1}(U) z^{m_1+m_2} \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee$. Notice that we have $\nabla_{\partial_{n_1}}(\alpha_1) \in \mathcal{W}_{P_1}^{s-1}(U)$ and $\nabla_{\partial_{n_1}}(\alpha_2) \in \mathcal{W}_{P_2}^{r-1}(U)$ and hence result follows by applying Lemma 4.22 again. When P_1 and P_2 are not intersecting transversally, it follows from the non-transversal case of Lemma 4.22 that all the terms lie in $\mathcal{W}_k^{-\infty}(U) z^{m_1+m_2} \otimes_{\mathbb{Z}} N$. \square

Remark 4.26. *The terms $z^{m_1+m_2} \check{\partial}_{(m_2, n_1)n_2 - (m_1, n_2)n_1}$ which appear in Lemma 4.25 come from $[z^{m_1} \check{\partial}_{n_1}, z^{m_2} \check{\partial}_{n_2}]$ using the formula 2.8. In particular, if we have both $(m_1, n_1) = 0$ and $(m_2, n_2) = 0$ (which means that both $z^{m_1} \check{\partial}_{n_1}$ and $z^{m_2} \check{\partial}_{n_2}$ are elements in the tropical vertex group), then the leading order term of $[\alpha_1 z^{m_1} \check{\partial}_{n_1}, \alpha_2 z^{m_2} \check{\partial}_{n_2}]$ is given by $\alpha_1 \wedge \alpha_2 z^{m_1+m_2} \check{\partial}_{(m_2, n_1)n_2 - (m_1, n_2)n_1}$, and $z^{m_1+m_2} \check{\partial}_{(m_2, n_1)n_2 - (m_1, n_2)n_1}$ is an element in the tropical vertex group as well. This property will be important to us in Section 5.3.*

At this point we are ready to go back to the asymptotic analysis of the gauge φ . Recall that there are two integral operators: \hat{I} defined in (4.7) and I defined in (4.14). If we restrict ourselves

to differential 1-forms, we can treat both \hat{I} and I as path integrals, where the choices of paths differ only by a path lying inside R , as shown in Figure 9. (Indeed, the requirement that, for any point $p \in U$, the unique flow line of v in U passing through p intersects R uniquely at a point $x \in R$ when we define I is equivalent to the condition that U contains the path ϱ_u when we define \hat{I} .)




FIGURE 9. The difference between I and \hat{I}

The key observation is that Lemma 4.23, which applies to the operator I , can be applied to \hat{I} as well because R is chosen so that $R \cap P = \emptyset$, and hence integration of terms with asymptotic support on R over any path in R will produce elements in $\mathcal{W}_*^{-\infty}(U)$.

We show by induction that the term $\hat{I} \left(\sum_{k \geq 0} \frac{ad_{\varphi^s}^k}{(k+1)!} \bar{\partial} \varphi^s \right)_{s+1}$ does not contribute to the leading \hbar order term in φ_{s+1} defined in (4.6). For that we take P to be codimension 1 hyperplane in U .

Lemma 4.27. *For the gauge $\varphi = \varphi_1 + \varphi_2 + \dots$ defined iteratively by (4.6), we have*

$$\varphi_s \in \bigoplus_{k \geq 1} \mathcal{W}_{I(P)}^0(U) z^{km} \check{\partial}_n t^s; \quad ad_{\varphi^s}^l(\bar{\partial} \varphi^s) \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq s(l+1)}} \mathcal{W}_P^0(U) z^{km} \check{\partial}_n t^j$$

for all $s \geq 1$ and $l \geq 1$, where $\varphi^s = \varphi_1 + \varphi_2 + \dots + \varphi_s$.

Proof. We prove by induction on s . The $s = 1$ case concerns the term $\varphi_1 = -\hat{I}(\Pi_1)$, and we have $ad_{\varphi_1}^l(\bar{\partial} \varphi_1) = -ad_{\varphi_1}^l(\Pi_1)$. Now $\Pi_1 \in \bigoplus_{k \geq 1} \mathcal{W}_P^1(U) z^{km} \check{\partial}_n t^1$ from Definition 4.2, so we have $\varphi_1 = -\hat{I}(\Pi_1) \in \bigoplus_{k \geq 1} \mathcal{W}_{I(P)}^0(U) z^{km} \check{\partial}_n t^1$ by Lemma 4.23. Applying Lemma 4.25 l times, together with the fact that $I(P)$ and P intersect transversally, we have $-ad_{\varphi_1}^l(\Pi_1) \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq l+1}} \mathcal{W}_P^0(U) z^{km} \check{\partial}_n t^j$.

The key here is that all the terms are linear combinations of $z^{km} \check{\partial}_n$'s, between which the Lie bracket vanish since m is tangent to P and n is normal to P , and hence the leading contribution in Lemma 4.25 vanishes.

Now we assume that the statement is true for all $s' \leq s$. The induction hypothesis together with the fact that $\Pi_{s+1} \in \bigoplus_{k \geq 1} \mathcal{W}_P^1(U) z^{km} \check{\partial}_n t^{s+1}$ imply that

$$\bar{\partial} \varphi_{s+1} = - \left(\Pi + \sum_{l \geq 0} \frac{ad_{\varphi^s}^l}{(l+1)!} \bar{\partial} \varphi^s \right)_{s+1} \in \bigoplus_{k \geq 1} \mathcal{W}_P^1(U) z^{km} \check{\partial}_n t^{s+1}.$$

Applying Lemma 4.23 to $\varphi_{s+1} = -\hat{I} \left(\Pi + \sum_{l \geq 0} \frac{ad_{\varphi^s}^l}{(l+1)!} \bar{\partial} \varphi^s \right)_{s+1}$ then gives $\varphi_{s+1} \in \bigoplus_{k \geq 1} \mathcal{W}_{I(P)}^0(U) z^{km} \check{\partial}_n t^{s+1}$.

The second statement follows by applying Lemma 4.25 multiple times with the same reasoning as above. This completes the proof. \square

By Lemma 4.27, we have

$$(4.15) \quad \varphi_s \in -\hat{I}(\Pi_s) + \bigoplus_{l \geq 1} \hat{I}(\mathcal{W}_P^0)(U) z^{lm} \check{\partial}_n t^s$$

for all s . Lemma 4.23 tells us that $\hat{I}(\mathcal{W}_P^0(U)) \subset \mathcal{W}_{I(P)}^{-1}(U)$, so $-\hat{I}(\Pi_s) \in \mathcal{W}_{I(P)}^0(U)$ is the only term which contributes to the leading order in \hbar . Since $I(P)$ is of codimension 0, $\mathcal{W}_{I(P)}^{-1}(U) \subset O_{loc}(\hbar^{1/2})$ (where $O_{loc}(\hbar^{1/2})$ is defined in Notations 4.10). We conclude that:

Proposition 4.28. *For the gauge $\varphi = \varphi_1 + \varphi_2 + \dots$ defined iteratively by (4.6), we have*

$$\varphi_s \in \begin{cases} \sum_{k \geq 1} a_{sk} z^{km} \check{\partial}_n t^s + \bigoplus_{k \geq 1} \mathcal{W}_{I(P)}^{-1}(U) z^{km} \check{\partial}_n t^s & \text{on } \check{p}^{-1}(H_+), \\ \bigoplus_{k \geq 1} \mathcal{W}_0^{-\infty}(U) z^{km} \check{\partial}_n t^s & \text{on } \check{p}^{-1}(H_-); \end{cases}$$

which implies that $\varphi = \psi + \sum_{j,k \geq 1} O_{loc}(\hbar^{1/2}) z^{km} \check{\partial}_n t^j$ over $\check{p}^{-1}(U \setminus P)$, or equivalently, $\check{\varphi} := \mathcal{F}^{-1}(\varphi) = \check{\psi} + \sum_{j,k \geq 1} O_{loc}(\hbar^{1/2}) z^{km} \check{\partial}_n t^j$ over $\check{X}_0 \setminus \check{p}^{-1}(P)$.

Remark 4.29. *Recall that the ansatz in Definition 4.2 is defined by multiplying δ_m to the wall crossing factor $\text{Log}(\Theta)$. But indeed the only properties that we need are $\delta_m \in \mathcal{W}_P^1(U)$, and $\hat{I}(\delta_m)$ has its leading \hbar order term given by 1 on H_+ .*

So Proposition 4.28 still holds for any solution to the Maurer-Cartan equation in Proposition 4.4 (or more generally, to the Maurer-Cartan equation of the quotient $\widehat{\text{dgLa}} \mathfrak{g}^*/\mathcal{E}^*(U)$ to be introduced in Section 5.2.1) of the form $\Pi \in -\sum_{j,k \geq 1} (a_{jk} \delta_{jk} + \mathcal{W}_P^0(U)) z^{km} \partial_n t^j$ with $\bar{\partial}\Pi = 0$ such that each $\delta_{jk}^{(i)} \in \mathcal{W}_{P_i}^1(U)$ and can be written as $\delta_{jk}^{(i)} = (\pi \hbar)^{-1/2} e^{-\frac{x^2}{\hbar}} dx$ in some neighborhood W_{P_i} of P_i , where x is some affine linear function on W_{P_i} such that P_i is defined by $x = 0$ locally and $\iota_{W_{P_i}} dx > 0$.

5. MAURER-CARTAN SOLUTIONS AND SCATTERING DIAGRAMS

In this section, we interpret the local scattering process, which produces a consistent extension $\mathcal{S}(\mathcal{D})$ of a scattering diagram \mathcal{D} consisting of two non-parallel walls, as arising from semiclassical limits (as $\hbar \rightarrow 0$) of a solution of the Maurer-Cartan (MC) equation.

5.1. Solving Maurer-Cartan equations in general. Let us begin by reviewing the process of solving MC equations in a general $\text{dgLa} (\mathbf{G}^*, \partial, [\cdot, \cdot])$. We will apply Kuranishi's method [38] to solve the MC equation using a homotopy which retracts \mathbf{G}^* to its cohomology and acts as the gauge fixing (see e.g. [35]).

Suppose that we are given an input $\Pi = \Pi_1 + \Pi_2 + \dots \in \mathbf{G}^1 \otimes \mathfrak{m}$ satisfying $\bar{\partial}\Pi = 0$, where $\Pi_k \in \mathbf{G}^1 \otimes \mathfrak{m}^k$ is homogeneous of degree k in t . We attempt to find $\Xi = \Xi_2 + \Xi_3 + \dots \in \mathbf{G}^1 \otimes \mathfrak{m}$, where $\Xi_k \in \mathbf{G}^1 \otimes \mathfrak{m}^k$ is homogeneous of degree k in t , such that $\Phi := \Phi_1 + \Phi_2 + \dots \in \mathbf{G}^1 \otimes \mathfrak{m}$, where each term $\Phi_k := \Pi_k + \Xi_k \in \mathbf{G}^1 \otimes \mathfrak{m}^k$ is homogeneous of degree k in t , gives a solution of the following MC equation, i.e.

$$(5.1) \quad \bar{\partial}\Phi + \frac{1}{2}[\Phi, \Phi] = 0.$$

We assume that there are chain maps ι, \mathcal{P} and homotopy H

$$(5.2) \quad H^*(G^*) \begin{array}{c} \xrightarrow{\iota} \\ \xleftarrow{\mathcal{P}} \end{array} G^* \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} H,$$

such that $\mathcal{P} \circ \iota = \text{Id}$, and $\text{Id} - \iota \circ \mathcal{P} = \bar{\partial}H + H\bar{\partial}$. Then, instead of the MC equation, we look for solutions Φ of the equation

$$(5.3) \quad \Phi = \Pi - \frac{1}{2}H[\Phi, \Phi].$$

This originates from a method of Kuranishi [38] used to solve the MC equation of the Kodaira-Spencer dgLa. His method can be generalized to a general dgLa as follows (see e.g. [43])

Proposition 5.1. *Suppose that Φ satisfies the equation (5.3). Then Φ satisfies the MC equation (5.1) if and only if $\mathcal{P}[\Phi, \Phi] = 0$.*

In general, the k -th equation of the above equation (5.3) is given by

$$(5.4) \quad \Xi_k + \sum_{j+l=k} \frac{1}{2}H[\Phi_j, \Phi_l] = 0,$$

and Ξ_k (recall that $\Xi = \Phi - \Pi$) is uniquely determined by Ξ_j , $j < k$. In this way, the solution Ξ to (5.3) is uniquely determined.

There is a beautiful way to express the unique solution Ξ as a sum of terms involving the input Π over directed trees (reminiscent of a Feynman sum). To this end, we will introduce the notions of a *directed tree* and a *directed tree with ribbon structure*, following [20].

Definition 5.2. *A (directed) k -tree T consists of the following data:*

- a finite set of vertices $\bar{T}^{[0]}$ together with a decomposition $\bar{T}^{[0]} = T_{in}^{[0]} \sqcup T^{[0]} \sqcup \{v_o\}$, where $T_{in}^{[0]}$, called the set of incoming vertices, is a set of size k and v_o is called the outgoing vertex (we also write $T_{\infty}^{[0]} := T_{in}^{[0]} \sqcup \{v_o\}$),
- a finite set of edges $\bar{T}^{[1]}$, and
- two boundary maps $\partial_{in}, \partial_o : \bar{T}^{[1]} \rightarrow \bar{T}^{[0]}$ (here ∂_{in} stands for incoming and ∂_o stands for outgoing)

satisfying all of the following conditions:

- (1) Every vertex $v \in T^{[0]}$ is trivalent, and satisfies $\#\partial_o^{-1}(v) = 2$ and $\#\partial_{in}^{-1}(v) = 1$.
- (2) Every vertex $v \in T_{in}^{[0]}$ has valency one, and satisfies $\#\partial_o^{-1}(v) = 0$ and $\#\partial_{in}^{-1}(v) = 1$; we let $T^{[1]} := \bar{T}^{[1]} \setminus \partial_{in}^{-1}(T_{in}^{[0]})$.
- (3) For the outgoing vertex v_o , we have $\#\partial_o^{-1}(v_o) = 1$ and $\#\partial_{in}^{-1}(v_o) = 0$; we let $e_o := \partial_o^{-1}(v_o)$ be the outgoing edge and denote by $v_r \in T_{in}^{[0]} \sqcup T^{[0]}$ the unique vertex (which we call the root vertex) with $e_o = \partial_{in}^{-1}(v_r)$.
- (4) The topological realization $|\bar{T}| := (\coprod_{e \in \bar{T}^{[1]}} [0, 1]) / \sim$ of the tree T is connected and simply connected; here \sim is the equivalence relation defined by identifying boundary points of edges if their images in $T^{[0]}$ are the same.

Two k -trees T_1 and T_2 are isomorphic if there are bijections $\bar{T}_1^{[0]} \cong \bar{T}_2^{[0]}$ and $\bar{T}_1^{[1]} \cong \bar{T}_2^{[1]}$ preserving the decomposition $\bar{T}_i^{[0]} = T_{i,in}^{[0]} \sqcup T_i^{[0]} \sqcup \{v_{i,o}\}$ and boundary maps $\partial_{i,in}$ and $\partial_{i,o}$. The set of isomorphism classes of k -trees will be denoted by \mathbb{T}_k . For a k -tree T , we will abuse notations and use T (instead of $[T]$) to denote its isomorphism class.

Definition 5.3. *A ribbon structure on a k -tree is a cyclic ordering of $\partial_{in}^{-1}(v) \sqcup \partial_o^{-1}(v)$ for each $v \in T^{[0]}$. Equivalently, it can be regarded as an embedding $|T| \hookrightarrow D$ of $|T|$ into the unit disc $D \subset \mathbb{R}^2$ mapping $T_{\infty}^{[0]}$ to ∂D , from which the cyclic ordering is induced by the clockwise orientation on D . We will use \mathbf{T} to denote a ribbon k -tree, and $\underline{\mathbf{T}}$ to denote the k -tree underlying \mathbf{T} .*

Two ribbon k trees \mathbf{T}_1 and \mathbf{T}_2 are isomorphic if they are isomorphic as k -trees and the isomorphism preserves the cyclic ordering. The set of isomorphism classes of ribbon k -trees will be denoted by \mathbb{RT}_k . We will again abuse notations by using \mathbf{T} to denote an isomorphism class of ribbon k -trees.

Definition 5.4. Given a ribbon k -tree $\mathbf{T} \in \mathbb{RT}_k$, we label the incoming vertices by v_1, \dots, v_k according to its cyclic ordering (or the clockwise orientation on D if we use the embedding $|T| \hookrightarrow D$). We define the operator $\mathfrak{l}_{k,\mathbf{T}} : L[1]^{\otimes k} \rightarrow L[1]$ by

- (1) aligning the inputs $\zeta_1, \dots, \zeta_k \in L$ at the vertices v_1, \dots, v_k respectively,
- (2) applying m_2 at each vertex in $\mathbf{T}^{[0]}$, where $m_2 : L[1] \otimes L[1] \rightarrow L[1]$ is the the graded symmetric operator on $L[1]$ ($= L$ shifted by degree 1) defined by $m_2(\alpha, \beta) := (-1)^{\bar{\alpha}(\bar{\beta}+1)}[\alpha, \beta]$ (here $\bar{\alpha}$ and $\bar{\beta}$ denote degrees of the elements $\alpha, \beta \in L$ respectively), and
- (3) applying the homotopy operator $-H$ to each edge in $\mathbf{T}^{[1]}$.

We then define $\mathfrak{l}_k : L[1]^{\otimes k} \rightarrow L[1]$ by $\mathfrak{l}_k := \sum_{\mathbf{T} \in \mathbb{RT}_k} \frac{1}{2^{k-1}} \mathfrak{l}_{k,\mathbf{T}}$.

The operation $\mathfrak{l}_{k,\mathbf{T}}$ can be symmetrized to give the following operation $\mathfrak{J}_{k,T}$ associated to a k -tree $T \in \mathbb{T}_k$:

Definition 5.5. Given a k -tree $T \in \mathbb{T}_k$, let $\mathbf{T} \in \mathbb{RT}_k$ be a ribbon tree whose underlying tree is $\underline{\mathbf{T}} = T$. We consider the set $\Sigma_k := \{\sigma \mid \sigma : T_{in}^{[0]} \rightarrow \{1, \dots, k\}\}$. Then we define the operator $\mathfrak{J}_{k,T} : \text{Sym}^k(L[1]) \rightarrow L[1]$ by

$$\mathfrak{J}_{k,T}(\zeta_1, \dots, \zeta_k) := \sum_{\sigma \in \Sigma_k} (-1)^{\chi(\sigma, \vec{\zeta})} \mathfrak{l}_{k,\mathbf{T}}(\zeta_{\sigma(1)}, \dots, \zeta_{\sigma(k)});$$

here the sign $(-1)^{\chi(\sigma, \vec{\zeta})}$ is determined by the rule that, when the permutation $(\zeta_1, \dots, \zeta_k) \mapsto (\zeta_{\sigma(v_1)}, \dots, \zeta_{\sigma(v_k)})$ is decomposed as a product of transpositions, each transposition interchanging ζ_i and ζ_j contributes $(-1)^{(\zeta_i+1)(\zeta_j+1)}$ (where $\bar{\zeta}_i$ denotes the degree of $\zeta_i \in L$). Note that $\mathfrak{J}_{k,T}$ is independent of the choice of the ribbon tree \mathbf{T} . We then define $\mathfrak{J}_k : \text{Sym}^k(L[1]) \rightarrow L[1]$ by $\mathfrak{J}_k = \sum_{T \in \mathbb{T}_k} \frac{\mathfrak{J}_{k,T}}{|Aut(T)|}$, where $|Aut(T)|$ is the order of the automorphism group of a k -tree T .

Setting

$$(5.5) \quad \Xi := \sum_{k \geq 2} \frac{1}{k!} \mathfrak{J}_k(\Pi, \dots, \Pi) = \sum_{k \geq 2} \mathfrak{l}_k(\Pi, \dots, \Pi),$$

then

$$(5.6) \quad \Phi := \Pi + \Xi = \sum_{k \geq 1} \frac{1}{k!} \mathfrak{J}_k(\Pi, \dots, \Pi) = \sum_{k \geq 1} \mathfrak{l}_k(\Pi, \dots, \Pi),$$

is the unique solution to the equation (5.3) obtained from recursively solving (5.4)

The equality between the two sums in (5.5) (and hence those in (5.6)) follows from the facts that the inputs are all the same and of degree 1, and simple combinatorial arguments in counting of trees. Also note that the sums in (5.6) are finite sums (mod \mathbf{m}^{N+1}) for every $N \in \mathbb{Z}_{>0}$ because $\Pi = \Pi_1 + \Pi_2 + \dots$ and $\Pi_k \in \mathbf{G}^1 \otimes_R \mathbf{m}^k$ so that, modulo \mathbf{m}^{N+1} , there are only finitely many trees and finitely many Π_k 's involved.

Remark 5.6. Both the operators $\mathfrak{J}_{k,T}$ and $\mathfrak{l}_{k,\mathbf{T}}$ will be used, but for different purposes: $\mathfrak{l}_{k,\mathbf{T}}$ does not involve automorphisms of trees, so it will be used in Section 5.2.3 to simplify some of the notations; while $\mathfrak{J}_{k,T}$ is conceptually more relevant to operations on $dgLa$'s, as we will see later.

Remark 5.7. Sum-over-trees formulas similar to 5.6 appear quite often in the literature, in particular in applications of the homological perturbation lemma [36] and study of L_∞ (or A_∞) structures [20].

5.2. Scattering of two non-parallel walls. Suppose we are given two non-parallel walls $\mathbf{w}_1 = (m_1, P_1, \Theta_1)$, and $\mathbf{w}_2 = (m_2, P_2, \Theta_2)$ where P_1, P_2 are oriented tropical hyperplanes intersecting in a codimension 2 tropical subspace $Q := P_1 \cap P_2$ in an affine convex coordinate chart $U \subset B_0$. The ansatz in Definition 4.2 gives two Maurer-Cartan (abbrev. MC) solutions $\Pi_{\mathbf{w}_i} \in \widehat{\mathbf{G}}^1(U)$, $i = 1, 2$, but their sum $\Pi := \Pi_{\mathbf{w}_1} + \Pi_{\mathbf{w}_2} \in \widehat{\mathbf{G}}^1(U)$ does *not* solve the MC equation (5.1).

As we mentioned in the Introduction, the method of Kuranishi [38] with a specific choice of the homotopy operator allow us to construct from Π a MC solution Φ of $\widehat{\mathbf{G}}^1(U)$ up to errors terms with exponential order in \hbar^{-1} , i.e. terms of the form $O(e^{-c/\hbar})$.¹² More precisely, we will construct MC solutions of the dgLa $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$, which is a quotient of a sub-dgLa of $\widehat{\mathbf{G}}^*(U)$, and show that they naturally give rise to consistent scattering diagrams.

We will first introduce the dgLa $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$ in Section 5.2.1 and construct a specific homotopy operator H in Section 5.2.2, before starting the asymptotic analysis of the MC solutions we constructed in Sections 5.2.3 and 5.2.4. The key results are Theorem 5.25 and Lemma 5.35.

5.2.1. *Solving the MC equation modulo error terms with exponential order in \hbar^{-1} .*

Definition 5.8. We define a dg-Lie subalgebra in $\mathbf{G}_N^*(U)$ by

$$\mathbf{g}_N^*(U) := \left(\bigoplus_m \mathcal{W}_*^\infty(U) z^m \right) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) \otimes_{\mathbb{C}} (R/\mathbf{m}^{N+1}),$$

where $\mathcal{W}_*^\infty(U) \subset \Omega_{\hbar}^*(U)$ is the space of differential forms with polynomial \hbar^{-1} order defined in 4.16. A general element of $\mathbf{g}_N^*(U)$ is a finite sum of the form $\sum_j \sum_{m,n} \alpha_{jm}^n z^m \check{\partial}_n t^j$, where $\alpha_{jm}^n \in \mathcal{W}_*^\infty(U)$. We have the inverse limit $\widehat{\mathbf{g}}^*(U) := \varprojlim \mathbf{g}_N^*(U)$.

There is a dg-Lie ideal $\mathcal{E}_N^*(U)$ of $\mathbf{g}_N^*(U)$ containing exponentially decay errors terms in \hbar^{-1} :

$$(5.7) \quad \mathcal{E}_N^*(U) := \left(\bigoplus_m \mathcal{W}_*^{-\infty}(U) z^m \right) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) \otimes_{\mathbb{C}} (R/\mathbf{m}^{N+1}),$$

where $\mathcal{W}_*^{-\infty}(U) \subset \Omega_{\hbar}^*(U)$ is the space of differential forms with exponential \hbar^{-1} order as in 4.15.

Then we take the quotient

$$(5.8) \quad \mathbf{g}_N^*(U)/\mathcal{E}_N^*(U) = \left(\bigoplus_m (\mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U)) z^m \right) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) \otimes_{\mathbb{C}} (R/\mathbf{m}^{N+1})$$

and define the dgLa $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$ as the inverse limit $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U) := \varprojlim (\mathbf{g}_N^*(U)/\mathcal{E}_N^*(U))$.

Remark 5.9. The advantage of working with the quotient $\mathcal{W}_*^\infty/\mathcal{W}_*^{-\infty}$ is that, given any element $\alpha \in \mathcal{W}_P^s(U)$ and any cut off function χ (independent of \hbar) such that $\chi \equiv 1$ in a neighborhood of P , we have $\alpha = \chi\alpha$ in the quotient $\mathcal{W}_P^s(U)/\mathcal{W}_*^{-\infty}(U)$, so an element in $\mathcal{W}_P^s(U)/\mathcal{W}_*^{-\infty}(U)$ can be treated as a delta function supported along P .

Lemma 5.10. For the dgLa $\mathbf{g}_N^*(U)/\mathcal{E}_N^*(U)$ in a contractible open subset U , we have $H^{>0}(\mathbf{g}_N^*(U)/\mathcal{E}_N^*(U)) = 0$ and

$$H^0(\mathbf{g}_N^*(U)/\mathcal{E}_N^*(U)) = \left(\bigoplus_m H^0(\mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U)) z^m \right) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) \otimes_{\mathbb{C}} (R/\mathbf{m}^{N+1})$$

where

$$H^0(\mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U)) = \frac{\{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq C\hbar^{-N} \text{ for some } C \text{ and } N\}}{\{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq Ce^{-c/\hbar} \text{ for some } c \text{ and } C\}}.$$

¹² Π is not a MC solution even up to such errors terms.

The proof of the above Lemma 5.10 relies on construction of homotopy operator. We will give the proof using the homotopy operator H constructed in Section 5.2.2 when U is a spherical neighborhood as described in Notation 5.11; the proof of a general contractible U works exactly in the same way by using the homotopy operator constructed from the map $\rho_{x^0} : [0, 1] \times U \rightarrow U$ which contracts U to a point x^0 .

5.2.2. *Construction of the homotopy operator.* Recall from Section 5.1 that a homotopy operator H (sometimes called a propagator) is needed for gauge fixing if we want to apply Kuranishi's method to solve the MC equation. To define this (and other operators), we may need to shrink U to a spherical neighborhood as follows.

Notations 5.11. *Suppose that we have two non-parallel walls $\mathbf{w}_i = (m_i, P_i, \Theta_i)$ ($i = 1, 2$) intersecting transversally in a codimension 2 tropical subspace $Q := P_1 \cap P_2$ in an affine convex open subset $V \subset B_0$. We fix a point $q_0 \in Q$. By reversing the orientations on P_i 's (and replacing Θ_i by Θ_i^{-1} accordingly) if necessary, we can choose the oriented normals of P_1 and P_2 to be $-m_1 = -\nu_{P_2}$ and $-m_2 = \nu_{P_1}$. We orient the rank 2 normal bundle NQ by the ordered basis $\{-m_1, -m_2\}$. By identifying an open neighborhood of the zero section in the normal bundle NQ with a tubular neighborhood of Q in B_0 , we see that the two walls are dividing $V \cap NQ$ into 4 quadrants; this can be visualized in the 2-dimensional slice $V \cap NQ_{q_0}$ in $V \cap NQ$, as shown in Figure 10*




FIGURE 10. The slice $V \cap NQ_{q_0}$ in $V \cap NQ$.


Now we fix local affine coordinates in V near q_0 , and choose an affine flat metric g_V with the property that m_1 , m_2 and TQ are perpendicular to each other. Then we choose a point x^0 in the third quadrant in NQ_{q_0} (see Figure 10) with $x^0 \notin (P_1 \cup P_2)$ and a ball $U \subset V$ (defined using the metric g_V) centered at x^0 which contains q_0 . We fix this neighborhood U centered at x^0 and call it a spherical neighborhood; see Figure 11. From this point on, we will work with a spherical neighborhood $U \subset B_0$ for the rest of this paper.

Since $TU = \Lambda_{B_0} \otimes_{\mathbb{Z}} \mathbb{R}$ and $\Gamma(U, \Lambda_{B_0}) \cong M$, we can identify an element $0 \neq m \in M$ with an affine vector field $m \in \Gamma(U, \Lambda_{B_0}) \subset \Gamma(U, TU)$ (with respect to the affine structure on B_0). We denote by U_m^\perp the tropical hyperplane perpendicular to m with respect to the metric g_V . Then U_m^\perp divides U into two half-spheres U_m^+ and U_m^- , which are named so that $-m$ is pointing into U_m^+ . The property that m_1 , m_2 and TQ are perpendicular to each other then implies that

$$(5.9) \quad Q \cap U \subset U_{a^1 m_1 + a^2 m_2}^+$$

for all $(a^1, a^2) \in (\mathbb{Z}_{\geq 0})^2 \setminus \{0\}$; see Figure 11.

To define a homotopy operator on $\mathbf{g}_N^*(U)$, we will first define one on the direct sum $\bigoplus_m \mathcal{W}_*^\infty(U) z^m$ and extend it by taking tensor product. For each $m \in M$, recall that we have $\bar{\partial}(\alpha z^m) = d(\alpha) z^m$,

FIGURE 11. The spherical neighborhood U .

where d is the deRham differential on U . So the cohomology $H^*(\mathcal{W}_*^\infty(U)z^m, \bar{\partial}) = \{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq C\hbar^{-N} \text{ for some } C \text{ and } N\}$ is represented by functions depending only on \hbar and with polynomial growth in \hbar^{-1} .

We are going to construct a homotopy operator H_m on $\mathcal{W}_*^\infty(U)z^m$ which retracts to its cohomology $H^*(\mathcal{W}_*^\infty(U)z^m) = H^*(\mathcal{W}_*^\infty(U))z^m$. The hyperplane U_m^\perp we chose above is playing the role of the reference hyperplane R when we define the operator I in (4.14) in Section 4.2.3. In the current situation, we need a family of reference hyperplanes $U_{a^1 m_1 + a^2 m_2}^\perp$, and the condition (5.9) is to ensure that we can define H_m in the same way as I and apply Lemma 4.23 for each $m = a^1 m_1 + a^2 m_2$.

Now for $0 \neq m \in M$, as in Lemma 4.23, we use flow lines along the affine vector field $-m$ to define a diffeomorphism $\tau_m : W_m \rightarrow U$, where $W_m \subset \mathbb{R} \times U_m^\perp$ is the maximal domain of definition of τ . Under the diffeomorphism τ_m , we obtain affine coordinates $(t =: u_{m,1}, u_{m,2}, \dots, u_{m,n})$ on U such that $x^0 = (0, \dots, 0)$. Note that these coordinates satisfy the condition in Notations 4.1 and we will set $u_{m,\perp} := (u_{m,2}, \dots, u_{m,n})$. For $m = 0 \in M$, we will choose an arbitrary set of local affine coordinates $(u_{0,1}, u_{0,2}, \dots, u_{0,n})$ in defining the homotopy operator H .

In the coordinates $(u_{m,1}, \dots, u_{m,n})$, we decompose a differential form $\alpha \in \mathcal{W}_*^\infty(U)$ uniquely as

$$(5.10) \quad \alpha = \alpha_0 + du_{m,1} \wedge \alpha_1,$$

where $\iota_{\frac{\partial}{\partial u_{m,1}}} \alpha_0 = \iota_{\frac{\partial}{\partial u_{m,1}}} \alpha_1 = 0$. We define a contraction $\rho_{m,\perp} : \mathbb{R} \times U_m^\perp \rightarrow U_m^\perp$ by $\rho_{m,\perp}(r, u_{m,\perp}) := ru_{m,\perp}$.

Definition 5.12. We define the homotopy operator $H_m : \mathcal{W}_*^\infty(U)z^m \rightarrow \mathcal{W}_{*-1}^\infty(U)z^m$ by $H_m(\alpha z^m) := (I_{m,er}(\alpha) + I_m(\alpha))z^m$, where we set

$$I_{m,er}(\alpha)(u_{m,1}, u_{m,\perp}) := \int_0^1 \rho_{m,\perp}^*(\alpha_0(0, \cdot)) = \int_0^1 (\iota_{\frac{\partial}{\partial r}} \rho_{m,\perp}^*(\alpha_0|_{U_m^\perp})) dr,$$

$$I_m(\alpha)(u_{m,1}, u_{m,\perp}) := \int_0^{u_{m,1}} \alpha_1(s, u_{m,\perp}) ds$$

using the decomposition of differential forms $\alpha \in \mathcal{W}_*^\infty(U)$ specified in (5.10).

We define the projection $\mathcal{P}_m : \mathcal{W}_*^\infty(U)z^m \rightarrow H^*(\mathcal{W}_*^\infty(U))z^m$ by

$$\mathcal{P}_m(\alpha z^m) := \begin{cases} (\alpha|_{x^0})z^m & \text{for } \alpha \text{ of degree } 0, \\ 0 & \text{otherwise,} \end{cases}$$

where $\alpha|_{x^0}$ is evaluation of α at the point x^0 and is to be treated as a constant function along U , and the operator $\iota_m : H^*(\mathcal{W}_*^\infty(U))z^m \rightarrow \mathcal{W}_*^\infty(U)z^m$ by $\iota_m(\alpha z^m) := \iota(\alpha)z^m$, where $\iota : H^*(\mathcal{W}_*^\infty(U)) \hookrightarrow \mathcal{W}_*^\infty(U)$ is the embedding of constant functions over U at degree 0 and 0 otherwise.

We will abuse notations by treating H_m , \mathcal{P}_m and ι_m as acting on the spaces $\mathcal{W}_*^\infty(U)$ and $H^*(\mathcal{W}_*^\infty(U))$.

Proposition 5.13. *The operator H_m is a homotopy retract of $\mathcal{W}_*^\infty(U)z^m$ onto its cohomology $H^*(\mathcal{W}_*^\infty(U))z^m$, i.e. we have $Id - \iota_m \mathcal{P}_m = \bar{\partial}H_m + H_m\bar{\partial}$.*

The integral operator H_m preserves $\mathcal{W}_*^{-\infty}(U)$ because the path integrals preserve terms with exponential decay in \hbar as one can see from Definition 4.15. Also, we have the natural identifications

$$\begin{aligned} H^*(\mathcal{W}_*^\infty(U), d) &= \{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq C\hbar^{-N} \text{ for some } C \text{ and } N\}, \\ H^*(\mathcal{W}_*^{-\infty}(U), d) &= \{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq Ce^{-c/\hbar} \text{ for some } c \text{ and } C\}, \end{aligned}$$

under which we see that the operators \mathcal{P}_m and ι_m can be descended to the quotient:

$$\begin{aligned} \mathcal{P}_m &: \mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U) \rightarrow H^*(\mathcal{W}_*^\infty(U), d)/H^*(\mathcal{W}_*^{-\infty}(U), d), \\ \iota_m &: H^*(\mathcal{W}_*^\infty(U), d)/H^*(\mathcal{W}_*^{-\infty}(U), d) \rightarrow \mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U), \end{aligned}$$

again in view of Definition 4.15. Therefore, Proposition 5.13 holds in the quotient $\mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U)$ as well.

Definition 5.14. *We define the operators $H := \bigoplus H_m$, $\mathcal{P} := \bigoplus \mathcal{P}_m$ and $\iota := \bigoplus \iota_m$ acting on the direct sum $\bigoplus_m \mathcal{W}_*^\infty(U)z^m$ and its cohomology. These operators extend naturally to the tensor product $\mathfrak{g}_N^*(U) = (\bigoplus_m \mathcal{W}_*^\infty(U)z^m) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) \otimes_{\mathbb{C}} (R/\mathfrak{m}^{N+1})$ and descend to the quotient $\mathfrak{g}_N^*(U)/\mathcal{E}_N^*(U)$ (since $\mathcal{E}_N^*(U)$ is a dg ideal of $\mathfrak{g}_N^*(U)$). We then take the inverse limit to define the operators H , \mathcal{P} and ι acting on $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$.*

Remark 5.15. *We remark that the homotopy operators defined above depend on the choices of U , the affine coordinates, etc, and so does the MC solution Φ that we are going to construct. However, the scattering diagram $\mathcal{D}(\Phi)$ associated to Φ is independent of these choices.*

Proof of Lemma 5.10. We prove this lemma for each direct summand $(\mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U))z^m$, which will be identified with $(\mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U))$ so that the Witten differential $\bar{\partial}$ becomes the usual deRham differential d . We have the following operators

$$\begin{aligned} H_m &: \mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U) \rightarrow \mathcal{W}_{*-1}^\infty(U)/\mathcal{W}_{*-1}^{-\infty}(U) \\ \mathcal{P}_m &: \mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U) \rightarrow \frac{\{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq C\hbar^{-N} \text{ for some } C \text{ and } N\}}{\{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq Ce^{-c/\hbar} \text{ for some } c \text{ and } C\}} \\ \iota_m &: \frac{\{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq C\hbar^{-N} \text{ for some } C \text{ and } N\}}{\{f : \mathbb{R}_{>0} \rightarrow \mathbb{R} \mid |f(\hbar)| \leq Ce^{-c/\hbar} \text{ for some } c \text{ and } C\}} \rightarrow \mathcal{W}_*^\infty(U)/\mathcal{W}_*^{-\infty}(U) \end{aligned}$$

defined in Definition 5.14. By descending the formula in Proposition 5.13 to the quotient by $\mathcal{W}_*^{-\infty}(U)$, we have $I - \iota_m \circ \mathcal{P}_m = \bar{\partial}H_m + H_m\bar{\partial}$. The result follows by a natural extension of this homotopy equation to $\mathfrak{g}_N^*(U)$. \square

5.2.3. Asymptotic analysis of Maurer-Cartan solutions. Going back to the two given non-parallel walls $\mathbf{w}_1 = (m_1, P_1, \Theta_1)$, $\mathbf{w}_2 = (m_2, P_2, \Theta_2)$. Recall that each wall crossing factor Θ_i is of the form $\text{Log}(\Theta_i) = \sum_{j,k \geq 1} a_{jk}^{(i)} w^{km_i} \check{\partial}_{n_i} t^j$, where $n_i \in \Lambda_{B_0}^\vee(U) \cong N$ is the unique primitive element satisfying $n_i \in (TP_i)^\perp$ and $(\nu_{P_i}, n_i) < 0$, and $\nu_{P_i} \in TU$ a normal to P_i so that the orientation on $TP_i \oplus \mathbb{R} \cdot \nu_{P_i}$ agrees with that on U (see Definition 3.3). As in Remark 4.29 in Section 4, we assume that the two inputs $\Pi^{(1)}, \Pi^{(2)}$ associated to the walls $\mathbf{w}_1, \mathbf{w}_2$ are of the following form:

Assumption 5.16. We assume that there are two MC solutions $\Pi^{(1)}, \Pi^{(2)}$ of $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$,¹³ which can be represented by elements in $\widehat{\mathfrak{g}^*}(U)$ of the form

$$(5.11) \quad \Pi^{(i)} \in - \sum_{j,k \geq 1} a_{jk}^{(i)} (\delta_{jk}^{(i)} + \mathcal{W}_{P_i}^0(U)) z^{km_i} \check{\partial}_{n_i} t^j$$

with $a_{jk}^{(i)} \neq 0$ only for finitely many k 's for each fixed j , and each $\delta_{jk}^{(i)} \in \mathcal{W}_{P_i}^1(U)$ can be written as $\delta_{jk}^{(i)} = (\pi\hbar)^{-1/2} e^{-\frac{\eta^2}{\hbar}} d\eta$ in some neighborhood W_{P_i} of P_i for some affine linear function η on W_{P_i} such that P_i is defined by $\eta = 0$ locally and $\iota_{\nu_{P_i}} d\eta > 0$.

For convenience, we will abuse notations and use $\Pi^{(i)} \in \widehat{\mathfrak{g}^*}(U)$ to denote its class in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ as well. We will also denote by

$$(5.12) \quad \check{\Pi}^{(i)} := - \sum_{j,k \geq 1} a_{jk}^{(i)} \delta_{jk}^{(i)} z^{km_i} \check{\partial}_{n_i} t^j$$

the leading order term of the input $\Pi^{(i)}$ for $i = 1, 2$. Then we have

$$(5.13) \quad \check{\Pi}^{(i)} \in \left(\bigoplus_{k \geq 1} \mathcal{W}_{P_i}^1(U) z^{km_i} \check{\partial}_{n_i} \right) [[t]]; \quad \Pi^{(i)} - \check{\Pi}^{(i)} \in \left(\bigoplus_{k \geq 1} \mathcal{W}_{P_i}^0(U) z^{km_i} \check{\partial}_{n_i} \right) [[t]].$$

We now solve the MC equation of the dgLa $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ by solving the equation (5.3) with the input data $\Pi := \Pi^{(1)} + \Pi^{(2)}$. Using the homotopy operator H and applying the sum over trees formula (5.6), we obtain an element

$$(5.14) \quad \Phi := \sum_{k \geq 1} \frac{1}{k!} \mathcal{J}_k(\Pi, \dots, \Pi) = \sum_{k \geq 1} \mathfrak{l}_k(\Pi, \dots, \Pi),$$

in $\widehat{\mathfrak{g}^*}(U)$, whose class in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ will also be denoted by Φ .

Lemma 5.17. The solution Φ constructed from the input $\Pi = \Pi^{(1)} + \Pi^{(2)}$ using the equation (5.3) and the homotopy operator H defined in Definition 5.14 is a MC solution in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$, i.e. we have $\mathcal{P}[\Phi, \Phi] = 0$ in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$.

We postpone the proof of Lemma 5.17 to Section 5.2.3. From Lemmas 5.17 and 5.10, we obtain a unique element $\varphi \in \widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ satisfying $e^\varphi * 0 = \Phi$ and $\mathcal{P}(\varphi) = 0$ in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ using Lemma 4.7. To start the asymptotic analysis of Φ and φ , we first decompose the Lie bracket $[\cdot, \cdot]$ on $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ into three types of operators:

Definition 5.18. For $\alpha = fz^m \check{\partial}_n$ and $\beta = gz^{m'} \check{\partial}_{n'}$ where $f, g \in \mathcal{W}_*^\infty(U)$, we decompose the Lie bracket $[\cdot, \cdot]$ into three operators $\mathfrak{h}, \mathfrak{h}'$ and \mathfrak{b} defined by

$$\begin{aligned} \mathfrak{h}(\alpha, \beta) &:= (-1)^{\bar{f}(\bar{g}+1)} fg [z^m \check{\partial}_n, z^{m'} \check{\partial}_{n'}], \\ \mathfrak{h}'(\alpha, \beta) &:= (-1)^{\bar{f}(\bar{g}+1)} f(\nabla_{\partial_n} g) z^{m+m'} \check{\partial}_{n'}, \\ \mathfrak{b}(\alpha, \beta) &:= (-1)^{(\bar{f}+1)\bar{g}} g(\nabla_{\partial_{n'}} f) z^{m+m'} \check{\partial}_n; \end{aligned}$$

here \bar{f} and \bar{g} denote the degrees of f and g respectively. These operators extend by linearity to $\mathfrak{g}_N^*(U)$ by treating a general element as a polynomial on the basis $\{z^m t^j\}$, and descend to the quotient $\mathfrak{g}_N^*(U)/\mathcal{E}_N^*(U)$, and can be further extended to $\widehat{\mathfrak{g}^*}(U)$ and $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ by taking inverse limits.

¹³Obviously MC solutions of the dg-Lie subalgebra $\widehat{\mathfrak{g}^*}(U) \subset KS_{\check{X}_0}(U)[[t]]$ descend to the quotient to give MC solutions of $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$.

Next we will decompose the operation \mathfrak{l}_k defined in Definition 5.4 according to the above decomposition of the Lie bracket $[\cdot, \cdot]$, the powers of the formal variable t and the Fourier modes $\{z^m\}$. For this purpose, we need to introduce the notion of a *labeled k -tree*:

Definition 5.19. *A labeled ribbon k -tree is a ribbon k -tree \mathcal{T} together with*

- a labeling of each trivalent vertex $v \in \mathcal{T}^{[0]}$ by \natural, \sharp or \flat , and
- a labeling of each incoming edge $e \in \partial_{in}^{-1}(\mathcal{T}_{in}^{[0]})$ by a pair (m_e, j_e) , where $m_e = km_i$ (for some $k > 0$ and $i = 1, 2$) specifies the Fourier mode and $j_e \in \mathbb{Z}_{>0}$ specifies the order of the formal variable t (corresponding to the input term $z^{km_i}t^{j_e}$ in $\hat{\mathfrak{g}}^*(U)$).

Similarly, we define a labeled k -tree as a k -tree \mathbb{T} together with a labeling of the trivalent vertices $\mathbb{T}^{[0]}$ by \natural or $\sharp + \flat$ (as only symmetric operations are allowed if there is no ribbon structure) and the same labeling of the incoming edges $\partial_{in}^{-1}(\mathbb{T}_{in}^{[0]})$ as above. We use $\underline{\mathbb{T}}$ to denote the underlying labeled k -tree of a labeled ribbon k -tree \mathcal{T} .

Two labeled ribbon k -trees \mathcal{T}_1 and \mathcal{T}_2 (resp. two labeled k -trees \mathbb{T}_1 and \mathbb{T}_2) are said to be isomorphic if they are isomorphic as ribbon k -trees (resp. k -trees) and the isomorphism preserves the labeling. The set of isomorphism classes of labeled ribbon k -trees (resp. labeled k -trees) will be denoted by LRT^k (resp. LT^k). As before, we will abuse notations by using \mathcal{T} (resp. \mathbb{T}) to stand for an isomorphism class of labeled ribbon k -trees (resp. labeled k -trees).

Notations 5.20. *For a labeled ribbon k -tree \mathcal{T} (resp. labeled k -tree \mathbb{T}), there is an induced labeling of all the edges in $\mathcal{T}^{[1]}$ (resp. $\mathbb{T}^{[1]}$) by the rule that at any trivalent vertex $v \in \mathcal{T}^{[0]}$ (resp. $v \in \mathbb{T}^{[0]}$) with two incoming edges e_1, e_2 and one outgoing edge e_3 , we set $(m_{e_3}, j_{e_3}) := (m_{e_1}, j_{e_1}) + (m_{e_2}, j_{e_2})$. We also write $(m_{\mathcal{T}}, j_{\mathcal{T}})$ (resp. $(m_{\mathbb{T}}, j_{\mathbb{T}})$) for the labeling of the unique edge e_o attached to the outgoing vertex v_o .*

Definition 5.21. *Given a labeled ribbon k -tree \mathcal{T} , we label the incoming vertices by v_1, \dots, v_k according to its cyclic ordering. We define the operator (similar to Definition 5.4) $\mathfrak{l}_{k, \mathcal{T}} : (\hat{\mathfrak{g}}^*(U)[1])^{\otimes k} \rightarrow \hat{\mathfrak{g}}^*(U)[1]$, for inputs ζ_1, \dots, ζ_k by*

- (1) extracting the coefficient of the term $z^{m_{e_i}}t^{j_{e_i}}$ in ζ_i and aligning it as the input at v_i ,
- (2) applying the operators \natural, \sharp or \flat to each trivalent vertex $v \in \mathcal{T}^{[0]}$ according to the labeling,
- (3) and applying the homotopy operator $-H$ to each edge in $\mathcal{T}^{[1]}$.

The operator $\mathfrak{l}_{k, \mathcal{T}}$ descends to $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ and will be denoted by the same notation.

Notations 5.22. *We decompose the set LRT^k of isomorphism classes of labeled ribbon k -trees into two parts: $\text{LRT}^k = \text{LRT}_0^k \sqcup \text{LRT}_1^k$, where LRT_0^k consists of trees whose trivalent vertices are all labeled by \natural and $\text{LRT}_1^k := \text{LRT}^k \setminus \text{LRT}_0^k$. We then consider the operators*

$$\mathfrak{l}_{k,0} := \sum_{\mathcal{T} \in \text{LRT}_0^k} \frac{1}{2^{k-1}} \mathfrak{l}_{k, \mathcal{T}}, \quad \mathfrak{l}_{k,1} := \sum_{\mathcal{T} \in \text{LRT}_1^k} \frac{1}{2^{k-1}} \mathfrak{l}_{k, \mathcal{T}}.$$

It is easy to see that for each labeled ribbon k -tree \mathcal{T} , the labeling $m_{\mathcal{T}}$ associated to the unique outgoing edge is of the form $m_{\mathcal{T}} = l(a_1m_1 + a_2m_2)$ for some $l \in \mathbb{Z}_{>0}$ and $(a_1, a_2) \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$, where $(\mathbb{Z}_{\geq 0})_{\text{prim}}^2$ denotes the set of all primitive elements in $(\mathbb{Z}_{\geq 0})^2 \setminus \{0\}$, so the solution Φ can be decomposed as a sum of Fourier modes parametrized by $(\mathbb{Z}_{\geq 0})_{\text{prim}}^2$.

Notations 5.23. *We let $m_a := a_1m_1 + a_2m_2$ for $a = (a_1, a_2) \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$. Note that we have $m_{(1,0)} = m_1$ and $m_{(0,1)} = m_2$. For each $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$, we let P_a be the tropical half-hyperplane $P_a = Q - \mathbb{R}_{\geq 0} \cdot m_a$. We equip each P_a with a normal ν_{P_a} such that $\{-m_a, \nu_{P_a}\}$ agrees with orientation given by $\{-m_1, -m_2\}$ on NQ , and this gives an orientation on P_a such that the orientation of $TP_a \oplus \mathbb{R} \cdot \nu_{P_a}$ agrees with that of B_0 (so that ν_{P_a} satisfies the condition in Definition 3.3 as well).*

Definition 5.24. Given the input $\Pi = \Pi^{(1)} + \Pi^{(2)} \in \hat{\mathbf{g}}^*(U)$, we put $\Psi := \sum_{k \geq 1} \mathfrak{l}_{k,0}(\check{\Pi}, \dots, \check{\Pi}) = \sum_{a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2} \Psi^{(a)}$, where

$$\Psi^{(a)} \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq N}} \sum_{n \in \Lambda_{B_0}^\vee(U) : n \perp m_a} \mathcal{W}_*^\infty(U) \cdot z^{km_a} \check{\partial}_n t^j \pmod{\mathfrak{m}^{N+1}}$$

for each $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$, and $F := \sum_{k \geq 1} (\mathfrak{l}_k(\Pi, \dots, \Pi) - \mathfrak{l}_{k,0}(\check{\Pi}, \dots, \check{\Pi})) = \sum_{a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2} F^{(a)}$, where

$$F^{(a)} \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq N}} \sum_{n \in \Lambda_{B_0}^\vee(U)} \mathcal{W}_*^\infty(U) \cdot z^{km_a} \check{\partial}_n t^j \pmod{\mathfrak{m}^{N+1}}$$

for each $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$.

Then we have $\Phi = \Psi + F$, where Ψ are the leading \hbar order terms and F are the error terms, as $\hbar \rightarrow 0$. We also put $\Phi^{(a)} := \Psi^{(a)} + F^{(a)}$. The key result on the asymptotic analysis of Φ is the following:

Theorem 5.25. For each $a \in (\mathbb{Z}_{>0})_{\text{prim}}^2$, we have

$$\Psi^{(a)} \in \left(\bigoplus_{k \geq 1} \mathcal{W}_{P_a}^1(U) z^{km_a} \check{\partial}_{n_a} \right) [[t]]; \quad F^{(a)} \in \left(\bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} \mathcal{W}_{P_a}^0(U) z^{km_a} \check{\partial}_n \right) [[t]],$$

where $n_a \in \Lambda_{B_0}^\vee(U)$ is the unique primitive normal to P_a such that $(\nu_{P_a}, n_a) < 0$.

Proof. According to the definitions of $\Psi^{(a)}$ and $F^{(a)}$ in Definition 5.24, this theorem is equivalent to the following statements:

$$\left\{ \begin{array}{l} \mathfrak{l}_{k,\mathcal{T}}(\check{\Pi}, \dots, \check{\Pi}) \in \mathcal{W}_{P_a}^1(U) z^{m_{\mathcal{T}}} \check{\partial}_{n_a} t^{j_{\mathcal{T}}} \quad \text{if } \mathcal{T} \in \text{LRT}_0^k, \\ \mathfrak{l}_{k,\mathcal{T}}(\Pi, \dots, \Pi) - \mathfrak{l}_{k,\mathcal{T}}(\check{\Pi}, \dots, \check{\Pi}) \in \sum_n \mathcal{W}_{P_a}^0(U) z^{\mathcal{T}} \check{\partial}_n t^{j_{\mathcal{T}}} \quad \text{if } \mathcal{T} \in \text{LRT}_0^k, \\ \mathfrak{l}_{k,\mathcal{T}}(\Pi, \dots, \Pi) \in \sum_n \mathcal{W}_{P_a}^0(U) z^{\mathcal{T}} \check{\partial}_n t^{j_{\mathcal{T}}} \quad \text{if } \mathcal{T} \in \text{LRT}_1^k. \end{array} \right.$$

The condition $n_a \perp P_a$ in the first statement follows from a simple induction argument using the formula (2.8); see the proof of Lemma 5.35 for more details. All other statements follow from Lemma 5.27 below. \square

To state Lemma 5.27, we fix a labeled ribbon k -tree \mathcal{T} whose incoming edges are e_1, \dots, e_k with labeling $(m_{e_1}, j_{e_1}), \dots, (m_{e_k}, j_{e_k})$ respectively.

We then consider the operation $\mathfrak{l}_{k,\mathcal{T}}((\alpha_1 \check{\partial}_{n_1}) z^{m_{e_1}} t^{j_{e_1}}, \dots, (\alpha_k \check{\partial}_{n_k}) z^{m_{e_k}} t^{j_{e_k}})$ for given $\alpha_1, \dots, \alpha_k \in \mathcal{W}_*^\infty(U)$ and $n_1, \dots, n_k \in \Gamma(U, \Lambda_{B_0}^\vee)$, defined in exactly the same way as in Definition 5.21, and treat it as an operation on $\alpha_1 \check{\partial}_{n_1}, \dots, \alpha_k \check{\partial}_{n_k}$ via the formula

$$(5.15) \quad \mathfrak{l}_{k,\mathcal{T}}(\alpha_1 \check{\partial}_{n_1}, \dots, \alpha_k \check{\partial}_{n_k}) z^{m_{\mathcal{T}}} t^{j_{\mathcal{T}}} := \mathfrak{l}_{k,\mathcal{T}}((\alpha_1 \check{\partial}_{n_1}) z^{m_{e_1}} t^{j_{e_1}}, \dots, (\alpha_k \check{\partial}_{n_k}) z^{m_{e_k}} t^{j_{e_k}}),$$

which will further be abbreviated as $\mathfrak{l}_{k,\mathcal{T}}(\vec{\alpha}, \vec{n}) := \mathfrak{l}_{k,\mathcal{T}}(\alpha_1 \check{\partial}_{n_1}, \dots, \alpha_k \check{\partial}_{n_k})$, where we put $\vec{\alpha} := (\alpha_1, \dots, \alpha_k)$ and $\vec{n} := (n_1, \dots, n_k)$.

Notations 5.26. Given a labeled ribbon k -tree \mathcal{T} and suppose that for each incoming edge e_i , we have assigned a closed codimension 1 tropical polyhedral subset P_{e_i} , which is either one of the two initial hyperplanes P_1, P_2 or one of the half-hyperplanes P_a 's introduced in Notation 5.23. We then inductively assign a (possibly empty) tropical hyperplane or half-hyperplane P_e to each edge $e \in \mathbb{T}^1$ as follows:

If e_1 and e_2 are two incoming edges meeting at a vertex v with an outgoing edge e_3 for which P_{e_1} and P_{e_2} are defined beforehand, we set $P_{e_3} := (Q - \mathbb{R}_{\geq 0} m_{e_3}) \cap U$ if both P_{e_1} and P_{e_2} are

non-empty and they intersect transversally at $Q := P_{e_1} \cap P_{e_2}$ and $P_{e_3} := \emptyset$ otherwise (recall that transversal intersection between two closed tropical polyhedral subsets, including the case when they have nonempty boundaries, was defined right before the proof of Lemma 4.22).

We denote the hyperplane or half-hyperplane associated to the unique outgoing edge e_o by $P_{\mathcal{T}}$. Note that if $P_{\mathcal{T}} \neq \emptyset$, then $P_{\mathcal{T}} = P_a$ for some $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$.

Lemma 5.27. *Given a labeled ribbon k -tree \mathcal{T} , each of whose incoming edges e_i is assigned with a closed codimension 1 tropical polyhedral subset P_{e_i} , which is either one of the two initial hyperplanes P_1, P_2 or one of the half-hyperplanes P_a 's introduced in Notation 5.23. Also given $\alpha_1, \dots, \alpha_k \in \mathcal{W}_*^\infty(U)$ and $n_1, \dots, n_k \in \Gamma(U, \Lambda_{B_0}^\vee)$ and suppose that α_i has asymptotic support (Definition 4.19) on P_{e_i} with either $\alpha_i \in \mathcal{W}_{P_{e_i}}^1(U)$ or $\alpha_i \in \mathcal{W}_{P_{e_i}}^0(U)$ for each $i = 1, \dots, k$, then we have*

$$\left\{ \begin{array}{ll} \mathfrak{l}_{k, \mathcal{T}}(\vec{\alpha}, \vec{n}) \in \mathcal{W}_{P_{\mathcal{T}}}^1(U) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) & \text{if } \mathcal{T} \in \text{LRT}_0^k, P_{\mathcal{T}} \neq \emptyset \text{ and } \alpha_i \in \mathcal{W}_{P_{e_i}}^1(U) \text{ for all } i, \\ \mathfrak{l}_{k, \mathcal{T}}(\vec{\alpha}, \vec{n}) \in \mathcal{W}_{P_{\mathcal{T}}}^0(U) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) & \text{if } \mathcal{T} \in \text{LRT}_0^k, P_{\mathcal{T}} \neq \emptyset \text{ and } \exists i \text{ such that } \alpha_i \in \mathcal{W}_{P_{e_i}}^0(U), \\ \mathfrak{l}_{k, \mathcal{T}}(\vec{\alpha}, \vec{n}) \in \mathcal{W}_{P_{\mathcal{T}}}^0(U) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) & \text{if } \mathcal{T} \in \text{LRT}_1^k \text{ and } P_{\mathcal{T}} \neq \emptyset, \\ \mathfrak{l}_{k, \mathcal{T}}(\vec{\alpha}, \vec{n}) \in \mathcal{W}_1^{-\infty}(U) \otimes_{\mathbb{Z}} \Lambda_{B_0}^\vee(U) & \text{if } P_{\mathcal{T}} = \emptyset. \end{array} \right.$$

For the purpose of the induction argument used to prove Lemma 5.27, we will temporarily relax the condition that $m_{e_i} = km_i$ for $i = 1, 2$ on the labeling of the incoming edges e_i 's in Definition 5.19 and replace it by the condition that $m_{e_i} = km_a$ for some $k > 0$ and some $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$.

Proof. We prove by induction on the number of vertices of a labeled ribbon k -tree \mathcal{T} . The initial step is trivial because $\text{LRT}^1 = \text{LRT}_0^1$ and $\mathfrak{l}_{1,0}$ is the identity.

We illustrate the induction step by considering the simplest non-trivial case, namely, when we have a labeled ribbon 2-tree \mathcal{T} with only one trivalent vertex v , two incoming edges e_1, e_2 and one outgoing edge e_o meeting v . Suppose that the incoming edges e_1, e_2 are assigned labeling $(m_{e_1}, j_{e_1}), (m_{e_2}, j_{e_2})$ and inputs $\alpha_1 \check{\partial}_{n_1} z^{e_1} t^{j_{e_1}}, \alpha_2 \check{\partial}_{n_2} z^{e_2} t^{j_{e_2}}$ respectively.

If $\mathcal{T} \in \text{LRT}_0^k$ (i.e. with labeling \natural at every $v \in \mathcal{T}^{[0]}$) and $P_{\mathcal{T}} \neq \emptyset$, then we have

$$\mathfrak{l}_{k, \mathcal{T}}(\vec{\alpha}, \vec{n}) = \mathfrak{l}_{k, \mathcal{T}}(\alpha_1 \check{\partial}_{n_1}, \alpha_2 \check{\partial}_{n_2}) = -H_{m_{\mathcal{T}}}(\alpha_1 \wedge \alpha_2) \check{\partial}_{n_{\mathcal{T}}},$$

where $n_{\mathcal{T}} = (m_{e_2}, n_1)n_2 - (m_{e_1}, n_2)n_1$ is given by the formula (2.8) (here we are viewing $H_{m_{\mathcal{T}}}$ as an operator on $\mathcal{W}_*^\infty(U)$ as in Definition 5.12).

The first case is when $\alpha_1, \alpha_2 \in \mathcal{W}_{P_{e_i}}^1(U)$. Since $P_{\mathcal{T}} \neq \emptyset$, the walls P_{e_1} and P_{e_2} are intersecting transversally at Q , so we have $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_{P_{e_1} \cap P_{e_2}}^2(U) = \mathcal{W}_Q^2(U)$ by Lemma 4.22. Recall the decomposition $H_{m_{\mathcal{T}}} = I_{m_{\mathcal{T}}} + I_{m_{\mathcal{T}}, er}$ in Definition 5.12. For the second integral $I_{m_{\mathcal{T}}, er}$, its domain of integration lies inside the hyperplane $U_{m_{\mathcal{T}}}^\perp$ which does not intersect Q by our choice of the spherical neighborhood U in Notations 5.11 (see (5.9) and Figure 11), so it produces terms in $\mathcal{W}_*^{-\infty}(U)$. For the first integral $I_{m_{\mathcal{T}}}$, applying Lemma 4.23 gives $-I_{m_{\mathcal{T}}}(\alpha_1 \wedge \alpha_2) \in \mathcal{W}_{P_{\mathcal{T}}}^1(U)$, where $P_{\mathcal{T}} = (Q - \mathbb{R}_{\geq 0} m_{\mathcal{T}}) \cap U$ as described in notation 5.26. This proves the first case.

For the second case, either $\alpha_1 \in \mathcal{W}_{P_{e_1}}^0(U)$ or $\alpha_2 \in \mathcal{W}_{P_{e_1}}^0(U)$, so we have $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_Q^1(U)$ by Lemma 4.22. The rest of the argument is the same as in the first case.

In the third case, we have $\mathcal{T} \in \text{LRT}_1^k$ and $P_{\mathcal{T}} \neq \emptyset$. This means that either \sharp or \flat is applied at v ; we will only give the proof for the case when \sharp is applied because the other case is similar. In such a case, we have $\mathfrak{l}_{k, \mathcal{T}}(\alpha_1 \check{\partial}_{n_1}, \alpha_2 \check{\partial}_{n_2}) = -H_{m_{\mathcal{T}}}(\alpha_1 \wedge (\nabla_{\partial_{n_1}} \alpha_2)) \check{\partial}_{n_2}$. Now $\nabla_{\partial_{n_1}}(\alpha_2) \in \mathcal{W}_{P_{e_2}}^0(U)$ by (4.12) and (2.7), so we get $\alpha_1 \wedge (\nabla_{\partial_{n_1}} \alpha_2) \in \mathcal{W}_Q^1(U)$ and the rest of the proof is same as in the first case.

Finally, for the fourth case we have $P_{\mathcal{T}} = \emptyset$, meaning that P_{e_1} and P_{e_2} not intersecting transversally. Then we have $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_2^{-\infty}(U)$ by Lemma 4.22 and the integral operator $H_{m_{\mathcal{T}}}$ preserves $\mathcal{W}_*^{-\infty}(U)$ by its definition in Definition 4.15. This completes the proof of labeled ribbon 2-tree.

Next, suppose that we have a general labeled ribbon k -tree \mathcal{T} , and $v_r \in \mathcal{T}^{[0]}$ is the unique trivalent vertex adjacent to the unique outgoing edge e_o . Assuming that \acute{e}_1 and \acute{e}_2 are the incoming edges connecting to v_r so that the edges $\acute{e}_1, \acute{e}_2, e_o$ are arranged in clockwise orientation. We split \mathcal{T} at v_r to obtain two trees $\mathcal{T}_1, \mathcal{T}_2$ with outgoing edges \acute{e}_1, \acute{e}_2 and k_1, k_2 incoming edges respectively such that $k = k_1 + k_2$. We split the inputs $(\vec{\alpha}, \vec{n})$ into two accordingly as $\vec{\alpha}_1 = (\alpha_1, \dots, \alpha_{k_1})$, $\vec{n}_1 = (n_1, \dots, n_{k_1})$ and $\vec{\alpha}_2 = (\alpha_{k_1+1}, \dots, \alpha_k)$, $\vec{n}_2 = (n_{k_1+1}, \dots, n_k)$. We then consider the operation $\mathfrak{l}_{k_i, \mathcal{T}_i}(\vec{\alpha}_i, \vec{n}_i)$ associated to each \mathcal{T}_i .

If one of $P_{\mathcal{T}_i}$'s is empty, say, if $P_{\mathcal{T}_1} = \emptyset$, then $\mathfrak{l}_{k_1, \mathcal{T}_1}(\vec{\alpha}_1, \vec{n}_1) \in \mathcal{W}_1^{-\infty}(U) \otimes_{\mathbb{Z}} \Lambda_{B_0}^{\vee}(U)$ by the induction hypothesis. Hence we also have $\mathfrak{l}_{k, \mathcal{T}}(\vec{\alpha}, \vec{n}) \in \mathcal{W}_1^{-\infty}(U) \otimes_{\mathbb{Z}} \Lambda_{B_0}^{\vee}(U)$ since $\mathcal{W}_*^{-\infty}(U)$ is a dg-Lie ideal of $\mathcal{W}_*^{\infty}(U)$ and H_m preserves $\mathcal{W}_*^{-\infty}(U)$.

So it remains to consider the case when $P_{\mathcal{T}_i} \neq \emptyset$ for $i = 1, 2$. Note that $\mathcal{T} \in \text{LRT}_0^k$ if and only if $\mathcal{T}_i \in \text{LRT}_0^{k_i}$ for both $i = 1, 2$ and the labeling of the root vertex v_r is also \natural , and $P_{\mathcal{T}} \neq \emptyset$ if and only if $P_{\acute{e}_1}$ intersects $P_{\acute{e}_2}$ transversally at Q . The induction step is completed by replacing $\alpha_i \check{\partial}_{n_i}$ with $\mathfrak{l}_{k_i, \mathcal{T}_i}(\vec{\alpha}_i, \vec{n}_i)$ for $i = 1, 2$ and using the same argument as in the case for labeled ribbon 2-tree. \square

Proof of Lemma 5.17. Recall that the spherical neighborhood U was chosen so that $x^0 \notin P_a$ for any $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$, where x^0 is the center of U (see Notations 5.11, (5.9) and Figure 11). By Theorem 5.25, we have a neighborhood $V \subset U$ of x^0 such that \bar{V} is compact and $\Phi|_V \in \mathcal{E}_N^*(V) \pmod{\mathfrak{m}^{N+1}}$ for every $N \in \mathbb{Z}_{>0}$. This implies that $[\Phi, \Phi]|_V \in \mathcal{E}_N^*(V) \pmod{\mathfrak{m}^{N+1}}$ for every $N \in \mathbb{Z}_{>0}$ since $\mathcal{E}_N^*(V)$ is closed under the bracket $[\cdot, \cdot]$. As the operator \mathcal{P} preserves $\mathcal{E}_N^*(V)$, we have $\mathcal{P}[\Phi, \Phi]|_V \in \mathcal{E}_N^*(V) \pmod{\mathfrak{m}^{N+1}}$ for every $N \in \mathbb{Z}_{>0}$. But then this means that $\mathcal{P}[\Phi, \Phi] = 0$ in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ since \mathcal{P} is the evaluation at $x^0 \in V \subset U$. \square

5.2.4. Leading \hbar order terms of a Maurer-Cartan solution. Recall from Section 5.2.3 that the leading order term Ψ (constructed in Definition 5.24) is a sum over labeled ribbon trees in LRT_0^k (i.e. those with only \natural labeling on trivalent vertices; see Notation 5.22) with inputs $\check{\Pi}^{(i)}$ (defined in (5.12)). This operation is closely related to the tropical vertex group as well as tropical counting. We are going to discuss the precise correspondence in this subsection.

For this purpose, it would conceptually be more appropriate to use labeled trees rather than labeled ribbon trees, because tropical trees are not equipped with ribbon structures. As in the case of labeled ribbon k -trees, we split the set of isomorphism classes of labeled k -trees into two components $\text{LT}^k = \text{LT}_0^k \sqcup \text{LT}_1^k$, where LT_0^k consists of those whose trivalent vertices are all labeled by \natural and $\text{LT}_1^k := \text{LT}^k \setminus \text{LT}_0^k$. Then given a labeled k -tree $\mathbb{T} \in \text{LT}_0^k$ and taking an arbitrary labeled ribbon k -tree \mathcal{T} with $\underline{\mathcal{T}} = \mathbb{T}$, we can define the operation $\mathfrak{J}_{k, \mathbb{T}}$ by $\mathfrak{J}_{k, \mathbb{T}}(\zeta_1, \dots, \zeta_k) := \sum_{\sigma \in \Sigma_k} (-1)^{\chi(\sigma, \vec{\zeta})} \mathfrak{l}_{k, \mathcal{T}}(\zeta_{\sigma(1)}, \dots, \zeta_{\sigma(k)})$, as in Definition 5.5.

Nonetheless, we prefer to work with labeled ribbon k -trees \mathcal{T} instead to simplify the formulas. There is a combinatorial relation $\frac{1}{k!|\text{Aut}(\mathbb{T})|} \mathfrak{J}_{k, \mathbb{T}}(\check{\Pi}, \dots, \check{\Pi}) = \sum_{\underline{\mathcal{T}} = \mathbb{T}} \frac{1}{2^{k-1}} \mathfrak{l}_{k, \mathcal{T}}(\check{\Pi}, \dots, \check{\Pi})$, where the sum is over all labeled ribbon k -trees \mathcal{T} with underlying labeled k -tree $\underline{\mathcal{T}} = \mathbb{T}$ and $|\text{Aut}(\mathbb{T})|$ is the order of the automorphism group of \mathbb{T} . Furthermore, since $\check{\Pi} \in \widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ has cohomological degree 1, we have

$$(5.16) \quad \frac{1}{k!|\text{Aut}(\mathbb{T})|} \mathfrak{J}_{k, \mathbb{T}}(\check{\Pi}, \dots, \check{\Pi}) = \frac{\#\{\underline{\mathcal{T}} = \mathbb{T}\}}{2^{k-1}} \mathfrak{l}_{k, \mathcal{T}_0}(\check{\Pi}, \dots, \check{\Pi}),$$

for an arbitrary labeled ribbon k -tree \mathcal{T}_0 with underlying labeled k -tree $\underline{\mathcal{T}_0} = \mathbb{T}$. We will obtain results for $\mathfrak{J}_{k, \mathbb{T}}$ by working with $\mathfrak{l}_{k, \mathcal{T}}$ through equation (5.16) in this section.

Given $\mathcal{T} \in \text{LRT}_0^k$ with $P_{\mathcal{T}} \neq \emptyset$ (recall from Notations 5.26 that $P_{\mathcal{T}}$ is the wall attached to the unique outgoing edge e_o), we have an alternative way to describe the operation $\mathfrak{l}_{k, \mathcal{T}}$. Recall that

$\mathcal{T}^{[1]}$ is the set of edges excluding the incoming edges (but including the outgoing edge) by Definition 5.2. We let τ_s^e be the flow of the affine vector field $-m_e$ for time s ($s \in \mathbb{R}_{\leq 0}$ so it is flowing backward in time), where (m_e, j_e) is the labeling of the edge $e \in \mathcal{T}^{[0]}$.

Definition 5.28. Given a sequence of edges $\mathfrak{e} = (e_0, e_1, \dots, e_l)$, as a path which starts from e_0 and ends at e_l following the direction of the tree \mathcal{T} , we define a map $\tau^\mathfrak{e} : W_\mathfrak{e} \rightarrow U$, by $\tau^\mathfrak{e}(\vec{s}, x) = \tau_{s_0}^{e_0} \circ \tau_{s_1}^{e_1} \circ \dots \circ \tau_{s_l}^{e_l}(x)$, where s_j is the time coordinate for the flow of $-m_{e_j}$, $\mathcal{T}_\mathfrak{e}^{[1]}$ is the subset $\{e_0, e_1, \dots, e_l\} \subset \mathcal{T}^{[1]}$ and $W_\mathfrak{e} \subset \mathbb{R}_{\leq 0}^{|\mathcal{T}_\mathfrak{e}^{[1]}|} \times U$ is the maximal domain such that the image of the flow $\tau^\mathfrak{e}$ lies in U . It can also be extended naturally to a map $\hat{\tau}^\mathfrak{e} : \hat{W}_\mathfrak{e} \rightarrow \mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]} \setminus \mathcal{T}_\mathfrak{e}^{[1]}|} \times U$, where $\hat{W}_\mathfrak{e} := \mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]} \setminus \mathcal{T}_\mathfrak{e}^{[1]}|} \times W_\mathfrak{e}$, by taking direct product with $\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]} \setminus \mathcal{T}_\mathfrak{e}^{[1]}|}$. Notice that this definition does not depend on the ribbon structure on \mathcal{T} , so it can be regarded as a definition for a labeled k -tree $\mathbb{T} := \underline{\mathcal{T}}$.

Definition 5.29. We attach a differential form ν_e on $\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|}$ to each $e \in \bar{\mathcal{T}}^{[1]}$ recursively by letting $\nu_e := 1$ for each incoming edge e , and $\nu_{e_3} = (-1)^{\bar{\nu}_{e_2}} \nu_{e_1} \wedge \nu_{e_2} \wedge ds_{e_3}$ (here $\bar{\nu}_{e_2}$ is the cohomological degree of ν_{e_2}) if v is an internal vertex with incoming edges $e_1, e_2 \in \bar{\mathcal{T}}_0$ and outgoing edge e_3 such that e_1, e_2, e_3 is clockwise oriented. We let $\nu_\mathcal{T}$ be the differential form attached to the unique outgoing edge $e_o \in \mathcal{T}^{[1]}$, which defines a volume form or orientation on $\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|}$.

As usual, we let v_1, \dots, v_k be the clockwise ordered incoming vertices of \mathcal{T} and e_1, \dots, e_k the incoming edges respectively. We associate to each e_i a unique sequence \mathfrak{e}_i of edges in $\mathcal{T}^{[1]}$ (excluding the incoming edge e_i itself) joining e_i to the outgoing edge e_o (including the outgoing edge e_o) along the direction of \mathcal{T} .

Remark 5.30. The subset $W_\mathcal{T} = (\bigcap_{i=1}^k \hat{W}_{\mathfrak{e}_i}) \subset \mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|} \times U$, can be viewed as a moduli space of tropical trees in U , denoted by $\mathfrak{M}_\mathcal{T}(U)$, with prescribed slope data $\{m_e\}_{e \in \mathcal{T}^{[1]}}$ (as in Notations 5.20) as follows (this will not be necessary for the rest of the paper): A point $(\vec{s}, x) \in W_\mathcal{T} \subset \mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|} \times U$ will prescribe the location of the vertices $\mathcal{T}^{[0]} \sqcup \{v_o\}$ of \mathcal{T} . First, $x \in U$ is the image of the outgoing vertex v_o . For any trivalent vertex $v \in \mathcal{T}^{[0]}$, there is a unique sequence of edges $\mathfrak{e} = (e_0, e_1, \dots, e_l)$ connecting v to v_o and $\tau^\mathfrak{e}(\vec{s}, x)$ is the image of v . The images of these vertices are allowed to overlap with each other as \vec{s} is taken from $\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|}$. Figure 12 illustrates the generic situation.




FIGURE 12. $W_\mathcal{T}$ parametrizing tropical trees in U

Recall that Definition 5.21, which defines the operator $\mathfrak{I}_{k, \mathcal{T}}$, uses the labeling (m_{e_i}, j_{e_i}) for each incoming edge $e \in \partial_{in}^{-1}(\mathcal{T}_{in}^{[0]})$ to extract the coefficient of $z^{m_{e_i} t^{j_{e_i}}}$ in $\check{\Pi}$ and then treat it as the input

at v_i . For the input $\alpha_i \check{\partial}_{n_i}$, we have $n_i \in TP_{e_i}^\perp$ and $(\nu_{P_i}, n_i) > 0$, and $\alpha_i = -\delta_{jk}^{(1)}$ or $-\delta_{jk}^{(2)}$ (see (5.12)) so that $\alpha_i \in \mathcal{W}_{P_{e_i}}^1(U)$. We decompose the output $\mathfrak{l}_{k,\mathcal{T}}(\vec{\alpha}, \vec{n})$ (defined in equation (5.15)) into a differential form part $\alpha_{\mathcal{T}} \in \mathcal{W}_{P_{\mathcal{T}}}^1(U)$ and a vector field part $\check{\partial}_{n_{\mathcal{T}}}$:

$$(5.17) \quad \mathfrak{l}_{k,\mathcal{T}}(\vec{\alpha}, \vec{n}) = \alpha_{\mathcal{T}} \check{\partial}_{n_{\mathcal{T}}},$$

where $\check{\partial}_{n_{\mathcal{T}}} := \mathfrak{l}_{k,\mathcal{T}}(\check{\partial}_{n_1}, \dots, \check{\partial}_{n_k})$; note that $n_{\mathcal{T}} \in \mathbb{Z} \cdot n_a$ if we write $P_{\mathcal{T}} = P_a$ for some $a \in (\mathbb{Z}_{\geq 0})_{\text{prim}}^2$, where $n_a \in \Lambda_{B_0}^\vee(U)$ is the unique primitive normal to $P_{\mathcal{T}} = P_a$ such that $(\nu_{P_a}, n_a) < 0$. The following lemma shows how $\alpha_{\mathcal{T}}(x)$ can be expressed as an integral over the space $\mathcal{I}_x := \left(\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|} \times \{x\} \right) \cap \left(\bigcap_{i=1}^k \hat{W}_{\epsilon_i} \right)$, for any $x \in U$ up to error terms of exponential order in \hbar^{-1} .

Lemma 5.31. *We have the identity*

$$\alpha_{\mathcal{T}}(x) = (-1)^{k-1} \int_{\mathcal{I}_x} (\tau^{\epsilon_1})^*(\alpha_1) \wedge \dots \wedge (\tau^{\epsilon_k})^*(\alpha_k)$$

in $\mathcal{W}_1^\infty(U)/\mathcal{W}_1^{-\infty}(U)$, where we use the volume form $\nu_{\mathcal{T}}$ on $\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|}$ for the integration on the right hand side.

Proof. We prove the lemma by induction on the number of vertices of a labeled ribbon k -tree \mathcal{T} (as in the proof of Lemma 5.27). In the initial step, \mathcal{I}_x is just the point $\{x\}$ and the right hand side is nothing but evaluation at x , so the result follows from the fact that $\mathfrak{l}_{k,\mathcal{T}}$ is the identity.

As in the proof of Lemma 5.27, we illustrate the induction step by considering the simplest non-trivial case when we are given a labeled ribbon 2-tree \mathcal{T} with only one trivalent vertex v , two incoming edges e_1, e_2 and one outgoing edge e_o meeting v . Suppose that the incoming edges e_1, e_2 are assigned labeling $(m_{e_1}, j_{e_1}), (m_{e_2}, j_{e_2})$ and inputs $\alpha_1 \check{\partial}_{n_1} z^{e_1} t^{j_{e_1}}, \alpha_2 \check{\partial}_{n_2} z^{e_2} t^{j_{e_2}}$ respectively. The operator $\mathfrak{l}_{k,\mathcal{T}}$ associated to \mathcal{T} is explicitly expressed as

$$(5.18) \quad \alpha_{\mathcal{T}} \check{\partial}_{n_{\mathcal{T}}} = \mathfrak{l}_{k,\mathcal{T}}(\alpha_1 \check{\partial}_{n_1}, \alpha_2 \check{\partial}_{n_2}) = -H_{m_{\mathcal{T}}}(\alpha_1 \wedge \alpha_2) \check{\partial}_{n_{\mathcal{T}}},$$

with $n_{\mathcal{T}} = (m_{e_2}, n_1)n_2 - (m_{e_1}, n_2)n_1$ given by the formula (2.8).

There are two cases depending on whether P_{e_1} and P_{e_2} are intersecting transversally or not, as in proof of Lemma 5.27. In both cases we can treat $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_Q^2(U)$ (because if intersection is not transversal, then we have $\alpha_1 \wedge \alpha_2 \in \mathcal{W}_2^{-\infty}(U) \subset \mathcal{W}_Q^2(U)$). From Definition 5.12, we have the decomposition $H_{m_{\mathcal{T}}} = I_{m_{\mathcal{T}}} + I_{m_{\mathcal{T}},er}$. By our choice of the spherical neighborhood U in Notations 5.11, the domain of integration for the second integral $I_{m_{\mathcal{T}},er}$ is supported away from Q , so it gives a term in $\mathcal{W}_1^{-\infty}(U)$. Thus we have $\alpha_{\mathcal{T}} \in -I_{m_{\mathcal{T}}}(\alpha_1 \wedge \alpha_2) + \mathcal{W}_1^{-\infty}(U)$.

In the current case, $\mathcal{T}^{[1]}$ consists of the unique outgoing edge e_o , so $\epsilon_1 = \epsilon_2 = (e_o)$ and hence the map $\tau^{\epsilon_1} = \tau^{\epsilon_2} : W_{\epsilon_i} \rightarrow U$ is simply given by the flow associated to $-m_{\mathcal{T}} = -m_{e_o}$. Now we have the half space $U_{m_{\mathcal{T}}}^+$ containing Q , as shown in Figure 11. Using the coordinates (t, u^\perp) on U where u^\perp are local affine coordinates on $U_{m_{\mathcal{T}}}^\perp$, we obtain a maximal interval $(a_{u^\perp}, b_{u^\perp})$ for each point u^\perp on $U_{m_{\mathcal{T}}}^\perp$ such that the interval $(a_{u^\perp}, b_{u^\perp}) \times \{u^\perp\}$ lies in U . Then the two integrals that we want to compare are $I_m(\alpha_1 \wedge \alpha_2)(t, u^\perp) = \int_0^t \iota_{\frac{\partial}{\partial s}}(\alpha_1 \wedge \alpha_2)(s, u^\perp) ds$ and $\int_{\mathcal{I}_{(t, u^\perp)}} (\tau^{\epsilon_1})^*(\alpha_1 \wedge \alpha_2) = \int_{a_{u^\perp}}^t \iota_{\frac{\partial}{\partial s}}(\alpha_1 \wedge \alpha_2)(s, u^\perp) ds$, the difference between which is given by $\int_{a_{u^\perp}}^0 \iota_{\frac{\partial}{\partial s}}(\alpha_1 \wedge \alpha_2)(s, u_{m_{\mathcal{T}}}^\perp) ds$, which produces a term in $\mathcal{W}_1^{-\infty}(U)$ because it misses the asymptotic support Q of $\alpha_1 \wedge \alpha_2$. This proves the statement for the current case.

Next we consider the induction step. We will adapt the same notations as the induction step in the proof of Lemma 5.27 (see Notations 5.26). So we take a general labeled ribbon k -tree $\mathcal{T} \in \text{LRT}_0^k$, and then split it at the unique vertex $v_r \in \mathcal{T}^{[0]}$ adjacent to the unique outgoing edge e_o to obtain two trees $\mathcal{T}_1, \mathcal{T}_2$ with incoming edges e_1, \dots, e_{k_1} and e_{k_1+1}, \dots, e_k respectively. We denote by $\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{k_1}$ (resp.

$\tilde{\mathbf{e}}_{k_1+1}, \dots, \tilde{\mathbf{e}}_k$) the sequences of edges in \mathcal{T}_1 (resp. \mathcal{T}_2) associated to the incoming edges e_1, \dots, e_{k_1} (resp. e_{k_1+1}, \dots, e_k) obtained respectively from the sequences $\mathbf{e}_1, \dots, \mathbf{e}_{k_1}$ (resp. $\mathbf{e}_{k_1+1}, \dots, \mathbf{e}_k$) of edges in \mathcal{T} by removing the unique outgoing edge e_o .

By the induction hypothesis, we have $(-1)^{k_1-1} \alpha_{\mathcal{T}_1}(x) = \int_{\mathcal{I}_{1,x}} (\tau^{\tilde{\mathbf{e}}_1})^*(\alpha_1) \wedge \dots \wedge (\tau^{\tilde{\mathbf{e}}_{k_1}})^*(\alpha_{k_1})$ modulo $\mathcal{W}_1^{-\infty}(U)$, where $\mathcal{I}_{1,x} = \left(\mathbb{R}_{\leq 0}^{|\mathcal{T}_1^{[1]}|} \times \{x\} \right) \cap \left(\bigcap_{i=1}^{k_1} \hat{W}_{\mathbf{e}_i}^{(1)} \right)$, and $(-1)^{k_2-1} \alpha_{\mathcal{T}_2}(x) = \int_{\mathcal{I}_{2,x}} (\tau^{\tilde{\mathbf{e}}_{k_1+1}})^*(\alpha_{k_1+1}) \wedge \dots \wedge (\tau^{\tilde{\mathbf{e}}_k})^*(\alpha_k)$ where $\mathcal{I}_{2,x} = \left(\mathbb{R}_{\leq 0}^{|\mathcal{T}_2^{[1]}|} \times \{x\} \right) \cap \left(\bigcap_{i=k_1+1}^k \hat{W}_{\mathbf{e}_i}^{(2)} \right)$; here $\hat{W}_{\mathbf{e}_i}^{(1)} \subset \mathbb{R}_{\leq 0}^{|\mathcal{T}_1^{[1]}|} \times U$ is the domain associated to $\tilde{\mathbf{e}}_i$ for the tree \mathcal{T}_1 as in Definition 5.28.

Fixing a point $x \in U$, we consider the flow $\tau^{e_o} : W_{e_o}(\subset \mathbb{R} \times U) \rightarrow U$ by $-m_{e_o} = -m_{\mathcal{T}}$ and let $\mathcal{I}_x^{e_o} := (\mathbb{R}_{\leq 0} \times \{x\}) \cap W_{e_o}$. From its definition, we have, for $x \in U$,

$$\mathcal{I}_x = \bigcup_{s \in \mathcal{I}_x^{e_o}} \mathcal{I}_{1, \tau_s^{e_o}(x)} \times \mathcal{I}_{2, \tau_s^{e_o}(x)} \times \{s\} \subset \mathbb{R}_{\leq 0}^{|\mathcal{T}_1^{[1]}|} \times \mathbb{R}_{\leq 0}^{|\mathcal{T}_2^{[1]}|} \times \mathbb{R}_{\leq 0}.$$

Using the same reasoning as in the 2-tree case, together with the fact that $\alpha_{\mathcal{T}_1} \wedge \alpha_{\mathcal{T}_2}$ is again having asymptotic support on Q , we have

$$\begin{aligned} (-1)^{k-1} \alpha_{\mathcal{T}}(x) &= (-1)^{k-2} \int_{\mathcal{I}_x^{e_o}} (\tau^{e_o})^*(\alpha_{\mathcal{T}_1} \wedge \alpha_{\mathcal{T}_2}) \\ &= \int_{\mathcal{I}_x^{e_o}} (\tau^{e_o})^* \left(\int_{\mathcal{I}_{1, \tau_s^{e_o}(x)}} (\tau^{\tilde{\mathbf{e}}_1})^*(\alpha_1) \dots (\tau^{\tilde{\mathbf{e}}_{k_1}})^*(\alpha_{k_1}) \wedge \int_{\mathcal{I}_{2, \tau_s^{e_o}(x)}} (\tau^{\tilde{\mathbf{e}}_{k_1+1}})^*(\alpha_{k_1+1}) \dots (\tau^{\tilde{\mathbf{e}}_k})^*(\alpha_k) \right) \\ &= \int_{\bigcup_{s \in \mathcal{I}_x^{e_o}} \mathcal{I}_{1, \tau_s^{e_o}(x)} \times \mathcal{I}_{2, \tau_s^{e_o}(x)} \times \{s\}} (\tau^{e_o})^* \left((\tau^{\tilde{\mathbf{e}}_1})^*(\alpha_1) \dots (\tau^{\tilde{\mathbf{e}}_k})^*(\alpha_k) \right) = \int_{\mathcal{I}_x} (\tau^{\mathbf{e}_1})^*(\alpha_1) \dots (\tau^{\mathbf{e}_k})^*(\alpha_k), \end{aligned}$$

modulo terms in $\mathcal{W}_1^{-\infty}(U)$. This completes the proof of the lemma. \square

Remark 5.32. Geometrically, Lemma 5.31 means that terms of the form $\alpha_{\mathcal{T}} \tilde{\delta}_{n_{\mathcal{T}}} z^{m_{\mathcal{T}}} t^{j_{\mathcal{T}}}$ for $\mathcal{T} \in \text{LRT}_0^k$, which appear in the leading order contribution of the Maurer-Cartan solution Φ (introduced in Definition 5.24), can be expressed as integrals over the moduli space $\mathfrak{M}_{\mathcal{T}}(U)$ of tropical trees in U .

We will see in Section 5.3 that what we essentially care about is the path integral

$$(5.19) \quad \int_{\varrho} \alpha_{\mathcal{T}} = (-1)^{k-1} \int_{\mathcal{I}_{\varrho}} (\tau^{\mathbf{e}_1})^*(\alpha_1) \dots (\tau^{\mathbf{e}_k})^*(\alpha_k) + O(e^{-c_e/\hbar})$$

along an embedded affine path $\varrho : (a, b) \rightarrow U$ that crosses the wall $P_{\mathcal{T}}$ transversally and positively (meaning that $TP_{\mathcal{T}} \oplus \mathbb{R} \cdot \varrho'$ agrees with the orientation of B_0), where we let

$$(5.20) \quad \mathcal{I}_{\varrho} := \bigcup_{t \in (a, b)} \mathcal{I}_{\varrho}(t).$$

Here \mathcal{I}_{ϱ} is equipped with coordinates $(\{s_e\}_{e \in \mathcal{T}^{[1]}}, t)$, where $(s_e)_e \in \mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|}$ and $t \in (a, b)$. We are going to calculate the integral in (5.19) explicitly.

Recall that $\alpha_i = -\delta_{jk}^{(1)}$ or $-\delta_{jk}^{(2)}$, and each α_i has asymptotic support on P_{e_i} which is either P_1 or P_2 (supports of the two initial walls). For each $i \in \{1, \dots, k\}$, we take an affine coordinate η_i associated to the corresponding $\delta_{jk}^{(i)}$ as in Definition 5.16, namely, such that $\{\eta_i = 0\} = P_{e_i}$ and $\iota_{\nu_{P_{e_i}}} d\eta_i > 0$, so that we can write $\alpha_i = -(\pi\hbar)^{-1/2} e^{-\frac{\eta_i^2}{\hbar}} d\eta_i$ locally near P_{e_i} .

The flow τ^{e_o} corresponding to the outgoing edge e_o is an affine map $\tau^{e_o}|_{\varrho} : \bigcup_{t \in (a, b)} \mathcal{I}_{\varrho}(t) \rightarrow U$, where $\mathcal{I}_{\varrho}(t) \subset \mathbb{R}_{\leq 0}$ is the maximal domain of backward flow associated to $-m_{e_o}$ starting from the

point $\varrho(t)$. The property that $\varrho \pitchfork P_{\mathcal{T}}$ is equivalent to the condition that the image of $\tau^{e_o}|_{\varrho}$ intersects transversally with Q at a point q_1 . For each $i \in \{1, \dots, k\}$, we let $N_i \subset U$ be the affine line through the point $q_1 \in Q$ transversal to P_{e_i} . Then we consider the affine space $\prod_{i=1}^k N_i$ with local affine coordinates η_1, \dots, η_k . See Figure 13.




FIGURE 13. The lines N_i 's through $q_1 \in Q$

We define the affine map

$$(5.21) \quad \vec{\tau} : \mathcal{I}_{\varrho} \rightarrow \prod_{i=1}^k N_i$$

by requiring that $(\vec{\tau})^*(\eta_i) = \eta_i(\tau^{e_i}(\vec{s}, x))$, where τ^{e_i} is defined in Definition 5.28.

Lemma 5.33. *There exists some constant $c > 0$ such that $(\vec{\tau})^*(d\eta_1 \wedge \dots \wedge d\eta_k) = c(-1)^{\chi(\mathcal{T})} \nu_{\mathcal{T}} \wedge dt$, where we set $(-1)^{\chi(\mathcal{T})} := \prod_{v \in \mathcal{T}^{[0]}} (-1)^{\chi(\mathcal{T}, v)}$ (with the convention that $(-1)^{\chi(\mathcal{T})} = 1$ if $\mathcal{T}^{[0]} = \emptyset$) and $(-1)^{\chi(\mathcal{T}, v)}$ is defined for each trivalent vertex v (attached to two incoming edges e_1, e_2 and one outgoing edge e_3 so that e_1, e_2, e_3 are arranged in the clockwise orientation) by comparing the orientation of the ordered basis $\{-m_{e_1}, -m_{e_2}\}$ with that of $\{-m_1, -m_2\}$ of NQ (cf. Notations 5.11). In particular, $\vec{\tau}$ is an affine isomorphism onto its image $C(\vec{\tau}) \subset \prod_{i=1}^k N_i$.*

Proof. Once again, we will prove by induction on the number of vertices of the labeled ribbon tree \mathcal{T} . The initial step concerning labeled ribbon 1-trees is trivial because in this case $\mathcal{I}_{\varrho} = (a, b)$ and $\vec{\tau} = \varrho$.

As before, we will consider the next step, or the simplest nontrivial case, namely, when we are given a labeled ribbon 2-tree \mathcal{T} with only one trivalent vertex v , two incoming edges e_1, e_2 and one outgoing edge e_o meeting v , to illustrate the induction step.

We first assume that the orientation of $\{-m_{e_1}, -m_{e_2}\}$ agrees with that of $\{-m_1, -m_2\}$. We can treat η_1, η_2 as oriented affine linear coordinates on the fiber NQ_{q_1} of the normal bundle NQ . We will use (s, t) for the coordinates of $\mathcal{I}_{\varrho} = \mathcal{I}_{\varrho}^{e_o}$ defined in (5.20). Let $x_0 := \vec{\tau}(0, t_0) \in \varrho \cap P_{\mathcal{T}}$ be the unique intersection point between ϱ and $P_{\mathcal{T}}$. Then there exists a unique $s_0 \leq 0$ such that $q_1 = \vec{\tau}(s_0, t_0) \in P_{e_1} \cap P_{e_2}$. We see that $\{(d\vec{\tau})_{(s_0, t_0)}(\frac{\partial}{\partial s}) = -m_{\mathcal{T}} \neq 0, (d\vec{\tau})_{(s_0, t_0)}(\frac{\partial}{\partial t})\}$ is an oriented basis of NQ_{q_1} by the assumption that ϱ intersects positively with $P_{\mathcal{T}}$; in other words, $(d\vec{\tau})_{(s_0, t_0)}$ is a linear isomorphism from $T_{(s_0, t_0)}\mathcal{I}_{\varrho}$ onto NQ_{q_1} and we have $(\vec{\tau})^*(d\eta_1 \wedge d\eta_2) = cds \wedge dt$ for some $c > 0$.

In the opposite case when the orientation of $\{-m_{e_1}, -m_{e_2}\}$ disagrees with that of $\{-m_1, -m_2\}$, η_2, η_1 are oriented coordinates of NQ_{q_1} . So we get $(\vec{\tau})^*(d\eta_1 \wedge d\eta_2) = -c ds \wedge dt = c(-1)^{\chi(\mathcal{T})} ds \wedge dt$ for some $c > 0$, because $\chi(\mathcal{T}) = 1$ in this case.

For the induction step, we again split a general k -tree $\mathcal{T} \in \text{LRT}_0^k$ at the root vertex v_r to get two trees \mathcal{T}_1 and \mathcal{T}_2 , as in the proof of Lemma 5.27. Since $P_{\mathcal{T}} \neq \emptyset$, both $P_{\mathcal{T}_1}$ and $P_{\mathcal{T}_2}$ are non-empty and they intersect transversally. We take two embedded paths ϱ_1 and ϱ_2 intersecting positively with $P_{\mathcal{T}_1}$ and $P_{\mathcal{T}_2}$ at q_1 with coordinates $\eta_{\mathcal{T}_1}$ and $\eta_{\mathcal{T}_2}$ respectively. By the induction hypothesis, the forms $(\tau^{\tilde{e}_1})^*(d\eta_1) \wedge \cdots \wedge (\tau^{\tilde{e}_{k_1}})^*(d\eta_{k_1}) = c_1(-1)^{\chi(\mathcal{T}_1)}\nu_{\mathcal{T}_1} \wedge d\eta_{\mathcal{T}_1}$ and $(\tau^{\tilde{e}_{k_1+1}})^*(d\eta_{k_1+1}) \wedge \cdots \wedge (\tau^{\tilde{e}_k})^*(d\eta_k) = c_2(-1)^{\chi(\mathcal{T}_2)}\nu_{\mathcal{T}_2} \wedge d\eta_{\mathcal{T}_2}$ are non-degenerate on \mathcal{I}_{ϱ_1} and \mathcal{I}_{ϱ_2} respectively. Therefore we have a nontrivial product $(\tau^{\tilde{e}_1})^*(d\eta_1) \wedge \cdots \wedge (\tau^{\tilde{e}_k})^*(d\eta_k) = (-1)^{\chi(\mathcal{T}_1)+\chi(\mathcal{T}_2)+\bar{\nu}_{\mathcal{T}_2}}\nu_{\mathcal{T}_1} \wedge \nu_{\mathcal{T}_2} \wedge d\eta_{\mathcal{T}_1} \wedge d\eta_{\mathcal{T}_2}$ on $\mathcal{I}_{\varrho_1} \times \mathcal{I}_{\varrho_2}$, where $\bar{\nu}_{\mathcal{T}_2}$ denotes the degree of the differential form $\nu_{\mathcal{T}_2}$.

Assuming that the orientation of $\{-m_{P_{\mathcal{T}_1}}, -m_{P_{\mathcal{T}_2}}\}$ agrees with that of $\{-m_1, -m_2\}$ on NQ_{q_1} , we can treat $\{\eta_{\mathcal{T}_1}, \eta_{\mathcal{T}_2}\}$ as an oriented basis for NQ_{q_1} . Using the same argument as in the 2-tree case, we use (s, t) as coordinates for $\bigcup_{t \in (a,b)} \mathcal{I}_{\varrho(t)}^{e_o}$ and obtain the relation $(\tau^{e_o}|_{\varrho})^*(d\eta_{\mathcal{T}_1} \wedge d\eta_{\mathcal{T}_2}) = cds \wedge dt$ for some $c > 0$. Combining with the induction hypothesis, we get

$$\begin{aligned} & (\tau^{e_1})^*(d\eta_1) \wedge \cdots \wedge (\tau^{e_k})^*(d\eta_k) = (\tau^{e_o}|_{\varrho})^* \left((\tau^{\tilde{e}_1})^*(d\eta_1) \wedge \cdots \wedge (\tau^{\tilde{e}_k})^*(d\eta_k) \right) \\ & = (-1)^{\chi(\mathcal{T}_1)+\chi(\mathcal{T}_2)+\bar{\nu}_{\mathcal{T}_2}} (\tau^{e_o}|_{\varrho})^* (\nu_{\mathcal{T}_1} \wedge \nu_{\mathcal{T}_2} \wedge d\eta_{\mathcal{T}_1} \wedge d\eta_{\mathcal{T}_2}) \\ & = (-1)^{\chi(\mathcal{T}_1)+\chi(\mathcal{T}_2)+\bar{\nu}_{\mathcal{T}_2}} (\tau^{e_o}|_{\varrho})^* (\nu_{\mathcal{T}_1} \wedge \nu_{\mathcal{T}_2}) \wedge ds \wedge dt = (-1)^{\chi(\mathcal{T})} \nu_{\mathcal{T}} \wedge dt; \end{aligned}$$

here we have $\chi(\mathcal{T}, v_r) = 0$ because we assume that the orientation of $\{-m_{P_{\mathcal{T}_1}}, -m_{P_{\mathcal{T}_2}}\}$ agrees with that of $\{-m_{P_{\mathcal{T}_1}}, -m_{P_{\mathcal{T}_2}}\}$.

Reversing the orientation condition, we will have $\chi(\mathcal{T}, v_r) = 1$, while at the same time we get an extra (-1) in the above formula because $(\tau^{e_o}|_{\varrho})^*(d\eta_{\mathcal{T}_1} \wedge d\eta_{\mathcal{T}_2}) = -cds \wedge dt$, exactly as in the 2-tree case. This completes the proof. \square

Now \mathcal{I}_{ϱ} is an open neighborhood of $\vec{0} \times \text{Im}(\varrho)$ in the cone $\mathbb{R}_{\leq 0}^{|\mathcal{T}^{[1]}|} \times \text{Im}(\varrho)$, and we let $C(\vec{\tau}) \subset \prod_{i=1}^k N_i$ be its image under the map $\vec{\tau}$. The local diffeomorphism $\vec{\tau}$ allows us to transform the integral in (5.19) to an integral over $C(\vec{\tau})$, so we have the identity

$$\begin{aligned} & \int_{x \in \varrho} \alpha_{\mathcal{T}} = (-1)^{k-1} \int_{C(\vec{\tau})} (\alpha_1 \wedge \cdots \wedge \alpha_k) + O(e^{-c_{\varrho}/\hbar}) \\ & = (-1)^{\chi(\mathcal{T})+1} (\pi\hbar)^{-\frac{k}{2}} \int_{C(\vec{\tau})} e^{-\frac{\sum_{i=1}^k \eta_i^2}{\hbar}} d\eta_1 \cdots d\eta_k + O(e^{-c_{\varrho}/\hbar}) = (-1)^{\chi(\mathcal{T})+1} \lim_{\epsilon \rightarrow 0} \frac{\text{vol}(C(\vec{\tau}) \cap B_{\epsilon})}{\text{vol}(B_{\epsilon})} + O(\hbar^{1/2}), \end{aligned}$$

which computes the leading order contribution of $\int_{x \in \varrho} \alpha_{\mathcal{T}}$; here B_{ϵ} is the ϵ -ball in $\prod_{i=1}^k N_i$ and vol is the volume with respect to the standard metric $\sum_{i=1}^k d\eta_i^2$.


Remark 5.34. *The meaning of the above equation is that the leading order contribution of $\int_{x \in \varrho} \alpha_{\mathcal{T}}$ (corresponding to the effect of crossing the new wall $P_{\mathcal{T}}$) depends on how the image $C(\vec{\tau})$ of the locus \mathcal{I}_{ϱ} in the moduli space $\mathfrak{M}_{\mathcal{T}}(U)$ of tropical trees in U intersects with the normals of the initial walls P_1, P_2 . Figure 14 illustrates the situation for a tree with only two incoming edges e_1, e_2 and one outgoing edge, where the ansatz for the initial walls P_1, P_2 are drawn as in Figure 4.*

The following lemma summarizes the results of this subsection:

Lemma 5.35. *For a labeled ribbon k -tree $\mathcal{T} \in \text{LRT}_0^k$ with $P_{\mathcal{T}} \neq \emptyset$, we write $\alpha_{\mathcal{T}} \check{\partial}_{n_{\mathcal{T}}} = \iota_{k, \mathcal{T}}(\vec{\alpha}, \vec{n})$ as in (5.17). Then, for any embedded affine line $\varrho : (a, b) \rightarrow U$ intersecting transversally and positively with $P_{\mathcal{T}}$, we have*

$$\int_{\varrho} \alpha_{\mathcal{T}} = (-1)^{\chi(\mathcal{T})+1} \lim_{\epsilon \rightarrow 0} \frac{\text{vol}(C(\vec{\tau}) \cap B_{\epsilon})}{\text{vol}(B_{\epsilon})} + O(\hbar^{1/2}),$$

where vol is the volume with respect to the standard metric $\sum_{i=1}^k d\eta_i^2$ on $\prod_{i=1}^k N_i$, and $\chi(\mathcal{T})$ is defined as in Lemma 5.33. Moreover, we have $n_{\mathcal{T}} \in (TP_{\mathcal{T}})^{\perp}$ and $(-1)^{\chi(\mathcal{T})} (\nu_{P_{\mathcal{T}}}, n_{\mathcal{T}}) < 0$.

FIGURE 14. Intersection $C(\vec{\tau})$ with P_i 's giving tropical counting

Proof. It remains to prove the last statement, which is yet another induction on the number of vertices of the labeled ribbon tree \mathcal{T} . The initial step is trivially true.

For the simplest non-trivial case, we look at a labeled ribbon 2-tree \mathcal{T} with only one trivalent vertex v , two incoming edges e_1, e_2 and one outgoing edge e_o meeting v , as before. Since $n_i \in (TP_i)^\perp$ and $(\nu_{P_{e_i}}, n_i) < 0$ for $i = 1, 2$, we have, by the formula (2.8), $n_{\mathcal{T}} = (m_{e_2}, n_1)n_2 - (m_{e_1}, n_2)n_1 \in (TP_{\mathcal{T}})^\perp$ since $P_{\mathcal{T}} = Q - \mathbb{R}_{\geq} \cdot m_{\mathcal{T}}$.

When the orientation of $\{-m_{e_1}, -m_{e_2}\}$ agrees with that of $\{-m_1, -m_2\}$, we can choose $-m_{e_2}$ as an oriented normal to $P_{\mathcal{T}}$ (here we only care about orientation so there are many different choices), so we have $(-m_{e_2}, n_{\mathcal{T}}) = (-m_{e_1}, n_2)(-m_{e_2}, n_1) < 0$ which means that $(\nu_{P_{\mathcal{T}}}, n_{\mathcal{T}}) < 0$ in this case.

When the orientation of $\{-m_{e_1}, -m_{e_2}\}$ disagrees with that of $\{-m_1, -m_2\}$, we have $(\nu_{P_{\mathcal{T}}}, n_{\mathcal{T}}) > 0$ from the above argument as now $-m_{e_1}$ is chosen as an oriented normal to $P_{\mathcal{T}}$. This completes the proof of the 2-tree case.

By (once again) splitting a k -tree $\mathcal{T} \in \text{LRT}_0^k$ at the root vertex v_r into two trees \mathcal{T}_1 and \mathcal{T}_2 , we can prove the induction step using exactly the same argument as above with n_1, n_2 replaced by $n_{\mathcal{T}_1}, n_{\mathcal{T}_2}$ respectively. \square

Remark 5.36. *The integral in Lemma 5.31 depends on the ribbon structure on \mathcal{T} because the order in taking the wedge product and the orientation of \mathcal{I}_x given by Definition 5.29 depend on it. Nevertheless, as Lemmas 5.33 and 5.35 show, the whole expression $\alpha_{\mathcal{T}} \check{\partial}_{\mathcal{T}}$ is independent of the ribbon structure, as only the sign of $\check{\partial}_{\mathcal{T}}$ depends on the ribbon structure and so does $\alpha_{\mathcal{T}}$, and this dependence cancels out with each other. This matches our earlier observation in equation (5.16) that the term $\mathfrak{I}_{k, \mathcal{T}}(\check{\Pi}, \dots, \check{\Pi})$ is independent of the ribbon structure.*

5.3. Consistent scattering diagrams from Maurer-Cartan solutions. In this section, we apply the results we obtained in Sections 5.2.3 and 5.2.4 to prove Theorems 1.4 and 1.5 in the Introduction.

5.3.1. The scattering diagram associated to the MC solution Φ . Recall that the Maurer-Cartan solution Φ constructed in (5.14) is decomposed as a sum of Fourier modes $\Phi^{(a)} = \Psi^{(a)} + F^{(a)}$ (see Definition 5.24). The asymptotic behavior of each $\Phi^{(a)}$ is described by Theorem 5.25 and a precise expression for the leading order terms $\Psi^{(a)}$ is obtained in Lemma 5.35. Applying these results, we are going to associate a scattering diagram $\mathcal{D}(\Phi)$ to Φ .

We will first construct a finite diagram $\mathcal{D}(\Phi)_N$ for each fixed $N \in \mathbb{Z}_{>0}$, producing a sequence $\{\mathcal{D}(\Phi)_N\}_{N \in \mathbb{Z}_{>0}}$ such that $\mathcal{D}(\Phi)_{N+1}$ is extension of $\mathcal{D}(\Phi)_N$ in the sense that there is an inclusion $\mathcal{D}(\Phi)_N \subset \mathcal{D}(\Phi)_{N+1}$ identifying walls (mod \mathbf{m}^{N+1}) and each $\mathbf{w} \in \mathcal{D}(\Phi)_{N+1} \setminus \mathcal{D}(\Phi)_N$ has a trivial wall crossing factor (mod \mathbf{m}^{N+1}). Then we define $\mathcal{D}(\Phi)$ as the limit of this sequence.

The order N scattering diagram $\mathcal{D}(\Phi)_N$ will be constructed by adding to the initial diagram $\mathcal{D}(\Phi)_1 = \{\mathbf{w}_1, \mathbf{w}_2\}$ new walls \mathbf{w}_a parametrized by a finite set of $a \in (\mathbb{Z}_{>0})_{\text{prim}}^2$, where each \mathbf{w}_a is supported on the half-hyperplane $P_a = Q - \mathbb{R}_{\geq 0} \cdot m_a$ and equipped with a wall crossing factor Θ_a (which could be trivial) determined from the leading order term $\Psi^{(a)}$ in the asymptotic expansion of $\Phi^{(a)}$. In order to parametrize the old and new walls by the same parameter space, we introduce the following notations:

Notations 5.37. We set $a \in \widetilde{(\mathbb{Z}_{\geq 0})}_{\text{prim}}^2 := (\mathbb{Z}_{\geq 0})_{\text{prim}}^2 \cup \{(-1, 0), (0, -1)\}$, and use the (rather unusual) convention that $m_{(-1,0)} = m_{(1,0)} = m_1$ and $m_{(0,-1)} = m_{(0,1)} = m_2$ for the Fourier modes corresponding to the two initial walls \mathbf{w}_1 and \mathbf{w}_2 . We use $\widetilde{(\mathbb{Z}_{\geq 0})}_{\text{prim}}^2$ to parametrize the set of half-hyperplanes P_a emanating from Q with slope $-m_a = -(a_1 m_1 + a_2 m_2)$ for $a = (a_1, a_2)$, where we are regarding each initial wall \mathbf{w}_i as a union of two half-hyperplanes.

For a fixed $N \in \mathbb{Z}_{>0}$, there will only be finitely many Fourier modes involved in the expression for the MC solution Φ in Definition 5.24. For this purpose, we use

$$\mathbb{W}(N) := \{a \in \widetilde{(\mathbb{Z}_{\geq 0})}_{\text{prim}}^2 \mid l m_a = m_{\mathcal{T}} \text{ for some } l \geq 1 \text{ and } \mathcal{T} \in \text{LRT}^k \text{ with } 1 \leq j_{\mathcal{T}} \leq N\},$$

where $(m_{\mathcal{T}}, j_{\mathcal{T}})$ is the labeling of the unique outgoing edge e_o attached to the outgoing vertex v_o in \mathcal{T} (see Definition 5.19 and Notations 5.20), to parametrize the possible walls involved in Φ (mod \mathbf{m}^{N+1}).

It makes sense to regard each of the two initial walls $\mathbf{w}_1, \mathbf{w}_2$ as a union of two half-hyperplanes in Notations 5.37 because of the following construction:

Definition 5.38. Given an input term $\Pi^{(i)}$ in the form of (5.11) having asymptotic support on P_i for $i = 1, 2$, we take an affine coordinate function $u_{m_i,1}$ along $-m_i$ which assumes the value 0 along Q . Then the functions

$$\chi_{i,+}(u_{m_i,1}) := \left(\frac{1}{\hbar\pi}\right)^{\frac{1}{2}} \int_{-\infty}^{u_{m_i,1}} e^{-\frac{s^2}{\hbar}} ds; \quad \chi_{i,-}(u_{m_i,1}) := 1 - \chi_{1,+}(u_{m_i,1}) = \left(\frac{1}{\hbar\pi}\right)^{\frac{1}{2}} \int_{u_{m_i,1}}^{\infty} e^{-\frac{s^2}{\hbar}} ds$$

have asymptotic support on $\{u_{m_i,1} \geq 0\} \cap U$ and $\{u_{m_i,1} \leq 0\} \cap U$ respectively, which implies that the cut-offs $\Phi^{(1,0)} := \chi_{1,+}\Pi^{(1)}$, and $\Phi^{(-1,0)} := \chi_{1,-}\Pi^{(1)}$ have asymptotic support on $\{u_{m_1,1} \geq 0\} \cap P_1$ and $\{u_{m_1,1} \leq 0\} \cap P_1$ respectively as well; the cut-offs $\Phi^{(0,\pm 1)}$ can be defined similarly using $\chi_{2,\pm}$ and they have asymptotic support on $\{u_{m_2,1} \geq 0\} \cap P_2$ and $\{u_{m_2,1} \leq 0\} \cap P_2$ respectively.

From this construction, we see that both $\bar{\partial}\Phi^{(1,0)}, \bar{\partial}\Phi^{(-1,0)}$ (resp. $\bar{\partial}\Phi^{(0,1)}, \bar{\partial}\Phi^{(0,-1)}$) have asymptotic support on $\{u_{m_1,1} = 0\} \cap P_1 = Q$ (resp. $\{u_{m_2,1} = 0\} \cap P_2 = Q$).

To prove that $\mathcal{D}(\Phi)_N$ is consistent (mod \mathbf{m}^{N+1}), we will remove $Q = P_1 \cap P_2 = \text{Sing}(\mathcal{D})$ from the spherical neighborhood U and apply a monodromy argument on the annulus $A := U \setminus Q$ by considering the universal cover $\mathbf{p} : \tilde{A} \rightarrow A$, which is endowed with the pullback affine structure from A . We use polar coordinates (r, θ) on a fiber of the normal bundle NQ (identified with a slice of a tubular neighborhood around Q) together with a set of affine coordinates $b := (b_3, \dots, b_n)$ on Q to get the coordinates $\hat{b} := (b_1 = r, b_2 = \theta, b_3, \dots, b_n)$ on \tilde{A} .¹⁴

We fix, once and for all, an angle θ_0 (chosen up to multiples of 2π) such that the half-hyperplane \mathcal{R}_{θ_0} with slope θ_0 through Q contains the center x^0 of the spherical neighborhood U (recall that

¹⁴Note that the polar coordinates (r, θ) are *not* affine coordinates.

the initial walls $\mathbf{w}_1, \mathbf{w}_2$ are dividing $U \cap NQ$ into 4 quadrants and x^0 lies in the 3rd quadrant; see Figures 10 and 11) and also a base point $\hat{b}^0 = (r_0, \theta_0, b^0) \in \mathcal{R}_{\theta_0}$ such that $\mathbf{p}(\hat{b}_0) = x^0$.

Notations 5.39. For each $a \in \mathbb{W}(N)$, we associate to the wall P_a an angle θ_a in the branch $\{(r, \theta, \hat{b}) \mid \theta_0 < \theta < \theta_0 + 2\pi\}$ to parametrize the lifting of $P_a \cap A$ in \tilde{A} . We identify $P_a \cap A$ with its lift in \tilde{A} , and will denote it again by P_a by abusing notations.

We choose a sufficiently small ϵ_0 and set $\mathbb{V} := \{(r, \theta, b) \mid \theta_0 - \epsilon_0 + 2\pi < \theta < \theta_0 + 2\pi\}$ so that the open subset $\mathbb{V} - 2\pi = \{(r, \theta, b) \mid \theta_0 - \epsilon_0 < \theta < \theta_0\}$ stays away from all the possible walls $\{\mathbf{w}_a\}$, as shown in Figure 15.




FIGURE 15.

For computation of some monodromy around Q , we consider the open subset $\tilde{A}_0 := \{(r, \theta, b) \mid \theta_0 - \epsilon_0 < \theta < \theta_0 + 2\pi\} \subset \tilde{A}$. Through the covering map $\mathbf{p} : \tilde{A}_0 \rightarrow A$, we pull back the dgLa's $\mathfrak{g}_N^*(A)$, $\mathcal{E}_N^*(A)$ and $\mathfrak{g}_N^*(A)/\mathcal{E}_N^*(A)$ to \tilde{A}_0 , and consider $\mathfrak{g}_N^*(\tilde{A}_0)$, $\mathcal{E}_N^*(\tilde{A}_0)$ and $\mathfrak{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$.

We write $\tilde{\Phi}^{(a)} := \mathbf{p}^*(\Phi^{(a)})$, $\tilde{\Psi}^{(a)} := \mathbf{p}^*(\Psi^{(a)})$, $\tilde{F}^{(a)} := \mathbf{p}^*(F^{(a)})$, and $\mathbf{p}^*(\Phi) := \sum_{a \in \mathbb{W}(N)} \mathbf{p}^*(\Phi^{(a)}) = \sum_{a \in \mathbb{W}(N)} \tilde{\Psi}^{(a)} + \tilde{F}^{(a)} \pmod{\mathfrak{m}^{N+1}}$, for the pullbacks to \tilde{A}_0 . We then have the following lemma.

Lemma 5.40. For each $a \in \widetilde{(\mathbb{Z}_{\geq 0})}_{\text{prim}}^2$, the Fourier mode $\tilde{\Phi}^{(a)}$ is itself a solution of the Maurer-Cartan equation (5.1) of the dgLa $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$ which further satisfies $\bar{\partial}\tilde{\Phi}^{(a)} = [\tilde{\Phi}^{(a)}, \tilde{\Phi}^{(a)}] = 0$.

Proof. For any fixed $N \in \mathbb{Z}_{>0}$, $\tilde{\Phi} = \sum_{a \in \mathbb{W}(N)} \tilde{\Phi}^{(a)} \pmod{\mathfrak{m}^{N+1}}$ by Definition 5.24 and $\mathbb{W}(N)$ is a finite set. Now for two different $a, a' \in \mathbb{W}(N)$, we have $(P_a \cap P_{a'}) \cap A = \emptyset$, and hence $[\tilde{\Phi}^{(a)}, \tilde{\Phi}^{(a')}] \in \mathcal{E}_N^2(\tilde{A}_0)$ which means that $[\tilde{\Phi}^{(a)}, \tilde{\Phi}^{(a')}] = 0$ in $\mathfrak{g}_N^2(\tilde{A}_0)/\mathcal{E}_N^2(\tilde{A}_0)$. Therefore each $\tilde{\Phi}^{(a)}$ is itself a MC solution in $\mathfrak{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$. Taking inverse limit shows that $\tilde{\Phi}^{(a)}$ is a MC solution in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$. Furthermore, as $\tilde{\Phi}^{(a)}$ is having asymptotic support on P_a , we have $[\tilde{\Phi}^{(a)}, \tilde{\Phi}^{(a)}] = 0$ in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$ by Lemma 4.25 (because P_a intersects itself non-transversally), so $\bar{\partial}\tilde{\Phi}^{(a)} = \bar{\partial}\tilde{\Phi}^{(a)} + \frac{1}{2}[\tilde{\Phi}^{(a)}, \tilde{\Phi}^{(a)}] = 0$. \square

Lemma 5.40 says that $\tilde{\Phi}^{(a)} = \tilde{\Psi}^{(a)} + \tilde{F}^{(a)}$ is a Maurer-Cartan solution in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$ with support concentrated along the wall P_a . Using a similar argument as the proof of Lemma 5.10, we note that the higher cohomologies of the complex $\mathfrak{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$ and $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$ are all trivial. So there are no non-trivial deformations of the dgLa $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$. In particular, the MC solution $\tilde{\Phi}^{(a)}$ is gauge equivalent to 0, i.e. there exists $\varphi_a \in \widehat{\mathfrak{g}^1/\mathcal{E}^1}(\tilde{A}_0)$ such that

$$(5.22) \quad e^{\varphi_a} * 0 = \tilde{\Psi}^{(a)} + \tilde{F}^{(a)}$$

on \tilde{A}_0 . As in the single wall case (Section 4.2), we need to define a homotopy operator $\hat{\mathcal{H}}$ on $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$ in order to fix the choice of φ_a .

We take a smooth homotopy $h : [0, 1] \times \tilde{A}_0 \rightarrow \tilde{A}_0$ contracting \tilde{A}_0 to the fixed point \hat{b}^0 with the property that $h(1, \hat{b}) = \hat{b}$ and $h(0, \hat{b}) = \hat{b}^0$. We define the homotopy operator $\hat{\mathcal{H}} : F_*^\infty(\tilde{A}_0) \rightarrow F_{*-1}^\infty(\tilde{A}_0)$ by

$$(5.23) \quad \hat{\mathcal{H}}(\alpha) := \int_0^1 h^*(\alpha)$$

for $\alpha \in F_*^\infty(\tilde{A}_0)$, and we also define $\hat{\mathcal{P}}$ by the evaluation at the base point \hat{b}^0 and \hat{i} by the embedding of constant functions on \tilde{A}_0 , as before (cf. Section 4.2); one can see that these operators descend to the quotient $F_*^\infty(\tilde{A}_0)/F_*^{-\infty}(\tilde{A}_0)$ using the same argument as in the explanation for Definition 5.14. We extend the above operators to the complex $\mathfrak{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$ as follows.

Definition 5.41. *We define the operators $\hat{\mathcal{H}}$, $\hat{\mathcal{P}}$ and \hat{i} by extending linearly the formulas $\hat{\mathcal{H}}(\alpha z^m \check{\partial}_n t^j) := \hat{\mathcal{H}}(\alpha) z^m \check{\partial}_n t^j$, $\hat{\mathcal{P}}(\alpha z^m \check{\partial}_n t^j) := \hat{\mathcal{P}}(\alpha) z^m \check{\partial}_n t^j$ and $\hat{i}(\alpha z^m \check{\partial}_n t^j) := \hat{i}(\alpha) z^m \check{\partial}_n t^j$, descending to the quotient and taking inverse limit.*

Definition 5.42. *Similar to the deduction of (4.6) from (4.5) [43], we solve the equation (5.22) in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$ iteratively to obtain the gauge: $\varphi_a := -\hat{\mathcal{H}}(ad_{\varphi_a}/(e^{ad_{\varphi_a}} - Id))(\tilde{\Psi}^{(a)} + \tilde{F}^{(a)})$ associated to $\tilde{\Phi}^{(a)} = \tilde{\Psi}^{(a)} + \tilde{F}^{(a)}$ which satisfies the gauge fixing condition $\hat{\mathcal{P}}(\varphi_a) = 0$; this gauge is unique by Lemma 4.7.*

We now apply asymptotic analysis to the gauge φ_a , similar to what we have done in the single wall case. First of all, as in Section 4.2.2 (see Remark 4.8 and the setup before Lemma 4.23), we shall replace $\hat{\mathcal{H}}$ by another operator $\hat{\mathcal{L}}$, which is defined using an integral over affine lines transversal to the wall P_a . For this purpose, we consider the half-space $\hat{\mathbb{H}}(P_a) := \{(r, \theta, b) \in \tilde{A}_0 \mid \theta \geq \theta_a\}$ in \tilde{A}_0 , on which φ_a is possibly having asymptotic support. Note that θ is not an affine coordinate but we can always express $\hat{\mathbb{H}}(P_a)$ as a tropical half-space in \tilde{A}_0 (by pulling back an affine linear function defining P_a to \tilde{A}_0 and parallel transporting to hyperplanes parallel to P_a).

We write $\tilde{\Psi}^{(a)} = \sum_{s=1}^\infty \tilde{\Psi}_s^{(a)}$, $\tilde{F}^{(a)} = \sum_{s=1}^\infty \tilde{F}_s^{(a)}$ and $\varphi_a = \sum_{s=1}^\infty \varphi_{a,s}$ according to powers of the formal variable t . We also set $\varphi_a^s := \varphi_{a,1} + \dots + \varphi_{a,s}$, $\tilde{\Psi}^{(a),s} := \tilde{\Psi}_1^{(a)} + \dots + \tilde{\Psi}_s^{(a)}$ and $\tilde{F}^{(a),s} := \tilde{F}_1^{(a)} + \dots + \tilde{F}_s^{(a)}$. Then we have the following lemma, which is parallel to Lemma 4.27 in Section 4.

Lemma 5.43. *The gauge φ_a has asymptotic support on the (codimension 0) tropical half-space $\hat{\mathbb{H}}(P_a) \subset \tilde{A}_0$, and we have*

$$\begin{aligned} \varphi_{a,s} &\in \bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{\hat{\mathbb{H}}(P_a)}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^s, \\ \varphi_{a,s} + \hat{\mathcal{H}}(\tilde{\Psi}_s^{(a)}) &\in \bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{\hat{\mathbb{H}}(P_a)}^{-1}(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^s, \\ ad_{\varphi_a^l}^l(\check{\partial} \varphi_a^s) &\in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq s(l+1)}} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{P_a}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^j \end{aligned}$$

for all $s \geq 1$ and $l \geq 1$.

Proof. We prove by induction on s (the power of the formal variable t). In the initial case, the equation defining $\varphi_{a,1}$ is $\varphi_{a,1} = -\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)} + \tilde{F}_1^{(a)})$. We want define an integral operator to replace $\hat{\mathcal{H}}$ in order to apply Lemma 4.23. Since all the assertions that we need to prove are local properties, we will work locally around any given point in \tilde{A}_0 .

So we fix a point $\hat{b}^1 \in \tilde{A}_0$ and choose a sufficiently small pre-compact open neighborhood $K \subset \tilde{A}_0$ of \hat{b}^1 , and then try to prove the initial case in K . We will also need to choose a family of piecewise affine lines, as in the single wall case. There are two scenarios:

- (1) If $\hat{b}^1 \in \hat{\mathbb{H}}(P_a)$, we choose a sufficiently small pre-compact open neighborhood K of \hat{b}^1 and a family of paths $\varrho_K : [0, 1] \times K \rightarrow \tilde{A}_0$ such that $\varrho_K(0, \hat{b}) = \hat{b}^0$, $\varrho_K(1, \hat{b}) = \hat{b}$ and there exists a partition $0 = t_0 < \dots < t_{l-1} < t_l = 1$ so that only one of the intervals $[t_{i_0-1}, t_{i_0}]$ has its image possibly intersecting with P_a and that $\varrho_K|_{[t_{i_0-1}, t_{i_0}]}$ is a flow line of the affine vector field v_K pointing into $\hat{\mathbb{H}}(P_a)$.
- (2) If $\hat{b}^1 \notin \hat{\mathbb{H}}(P_a)$, we choose a sufficiently small pre-compact open neighborhood K of \hat{b}^1 with $K \cap \hat{\mathbb{H}}(P_a) = \emptyset$ and a family of paths $\varrho_K : [0, 1] \times K \rightarrow \tilde{A}_0$ such that $\varrho_K(0, \hat{b}) = \hat{b}^0$, $\varrho_K(1, \hat{b}) = \hat{b}$ and $\text{Im}(\varrho_K) \cap P_a = \emptyset$.

Such a family always exists when K is sufficiently small. Figure 16 illustrates the difference between the integral operators $\hat{\mathcal{H}}$ and $\hat{\mathcal{I}}_{a,K}$. Then we set $\hat{\mathcal{I}}_{a,K}(\alpha)(\hat{b}) := \int_0^1 \varrho^*(\alpha)(s, \hat{b})$. Applying Lemma 4.23




FIGURE 16.

to the piece $\varrho_K|_{[t_{i_0-1}, t_{i_0}]}$ gives

$$-\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)}) \in F_{\hat{\mathbb{H}}(P_a)}^0(K) \cdot (z^{km_a} \check{\partial}_{n_a}) t^1; \quad -\hat{\mathcal{H}}(\tilde{F}_1^{(a)}) \in \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{\hat{\mathbb{H}}(P_a)}^{-1}(K) \cdot (z^{km_a} \check{\partial}_n) t^1,$$

which proves the first two assertions in the initial case.

For the third assertion, we have

$$\bar{\partial} \varphi_{a,1} = -\bar{\partial} \hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)} + \tilde{F}_1^{(a)}) = -\tilde{\Psi}_1^{(a)} - \tilde{F}_1^{(a)} \in \bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{P_a}^1(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^1$$

from the gauge fixing condition in Definition 5.42. Upon repeated applications of Lemma 4.25, we have $ad_{\varphi_a^1}^l(\tilde{F}_1^{(a)}) \in \bigoplus_{1 \leq j \leq l+1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{P_a}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^j$, so we only need to take care of the term $ad_{\varphi_a^1}^l(\tilde{\Psi}_1^{(a)})$. Writing $\varphi_{a,1} = -\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)} + \tilde{F}_1^{(a)})$ and applying Lemma 4.25 again, we see that the only term we have to consider is $ad_{-\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)})}^l(\tilde{\Psi}_1^{(a)})$ in the expression $ad_{-\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)} + \tilde{F}_1^{(a)})}^l(\tilde{\Psi}_1^{(a)})$, because the appearance of any of $ad_{-\hat{\mathcal{H}}(\tilde{F}_1^{(a)})}$ in the above expression will result in a term in

$\bigoplus_{j,k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{P_a}^0(\tilde{A}_0) \cdot (z^{km_a} \check{\partial}_n) t^j$. Concerning the term $ad_{-\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)})}^l(\tilde{\Psi}_1^{(a)})$, Theorem 5.25 says that that $n_a \perp P_a$ in the expressions

$$\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)}) \in \bigoplus_{k \geq 1} F_{\hat{\mathbb{H}}(P_a)}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_{n_a} t^1; \quad \tilde{\Psi}_1^{(a)} \in \bigoplus_{k \geq 1} F_{P_a}^1(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_{n_a} t^1,$$

meaning that the leading order term of $ad_{-\hat{\mathcal{H}}(\tilde{\Psi}_1^{(a)})}(\tilde{\Psi}_1^{(a)})$ given by Lemma 4.25 vanishes. Hence the third assertion follows.

Now we assume that the assertions hold for $s' \leq s$. We consider the equation $\varphi_{a,s+1} = -\hat{\mathcal{H}}(\tilde{\Psi}_{s+1}^{(a)}) + \tilde{F}_{s+1}^{(a)} + \sum_{k \geq 0} \frac{ad_{\varphi_a^s}^k}{(k+1)!} \bar{\partial} \varphi_a^s$ which determines $\varphi_{a,s+1}$ iteratively. From the induction hypothesis, we have

$$\tilde{F}_{s+1}^{(a)} + \left(\sum_{k \geq 0} \frac{ad_{\varphi_a^s}^k}{(k+1)!} \bar{\partial} \varphi_a^s \right)_{s+1} \in \bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{P_a}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^{s+1}.$$

Applying $\hat{\mathcal{H}}$ (replacing $\hat{\mathcal{H}}$ by $\hat{\mathcal{L}}_{a,K}$ again) to this expression give the first two assertions of the induction step by Lemma 4.23.

For the third assertion, we have $\bar{\partial} \varphi_{a,s+1} = -(\tilde{\Psi}_{s+1}^{(a)} + \tilde{F}_{s+1}^{(a)} + \sum_{k \geq 0} \frac{ad_{\varphi_a^s}^k}{(k+1)!} \bar{\partial} \varphi_a^s)_{s+1}$ in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\tilde{A}_0)$, again from the gauge fixing condition in Definition 5.42. Applying Lemma 4.25 as in the proof of the initial step, we see that the essential term to be considered is $ad_{-\hat{\mathcal{H}}(\tilde{\Psi}^{(a),s+1})}^l(\tilde{\Psi}^{(a),s+1})$. By Theorem 5.25 again, we have $n \perp P_a$ in the following expressions

$$\hat{\mathcal{H}}(\tilde{\Psi}^{(a),s+1}) \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq s+1}} F_{\hat{\mathbb{H}}(P_a)}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_{n_a} t^j; \quad \tilde{\Psi}^{(a),s+1} \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq s+1}} F_{P_a}^1(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_{n_a} t^j.$$

Then we can conclude that $ad_{-\hat{\mathcal{H}}(\tilde{\Psi}^{(a),s+1})}^l(\tilde{\Psi}^{(a),s+1}) \in \bigoplus_{\substack{k \geq 1 \\ 1 \leq j \leq (s+1)(l+1)}} F_{P_a}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_{n_a} t^j$ because the leading order term given by Lemma 4.25 vanishes, as in the proof of the initial step. This finishes the proof of the induction step. \square

The following lemma is parallel to Proposition 4.28 in Section 4.

Lemma 5.44. *Over the half-space $\hat{\mathbb{H}}(P_a) \setminus P_a$, we have*

$$\varphi_a \in \psi_a + \left(\bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{\hat{\mathbb{H}}(P_a) \setminus P_a}^{-1}(\hat{\mathbb{H}}(P_a) \setminus P_a) \cdot z^{km_a} \check{\partial}_n \right) [[t]] \cdot t,$$

where $\psi_a = \text{Log}(\Theta_a)$ for some element Θ_a of the tropical vertex group of the form $\psi_a = \sum_{j,k \geq 1} b_{jk}^{(a)} \cdot z^{km_a} \check{\partial}_{n_a} t^j$, where $b_{jk}^{(a)}$'s are constants independent of \hbar with $b_{jk}^{(a)} \neq 0$ only for finitely many k 's for each fixed j and n_a is the unique primitive normal to P_a satisfying $(\nu_{P_a}, n_a) < 0$; while over the other half-space $\tilde{A}_0 \setminus \hat{\mathbb{H}}(P_a)$, we have $\varphi_a = 0$.

Proof. We first consider φ_a over $\hat{\mathbb{H}}(P_a) \setminus P_a$. From the proof of Lemma 5.43, we see that

$$\bar{\partial} \varphi_{a,s} = - \left(\tilde{\Psi}_s^{(a)} + \tilde{F}_s^{(a)} + \sum_{k \geq 0} \frac{ad_{\varphi_a^{s-1}}^k}{(k+1)!} \bar{\partial} \varphi_a^{s-1} \right)_s \in \bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^\vee(U)} F_{P_a}^0(\tilde{A}_0) \cdot z^{km_a} \check{\partial}_n t^s$$

for every $s \geq 1$. In particular, we have $\bar{\partial} \varphi_{a,s} = 0$ in $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\hat{\mathbb{H}}(P_a) \setminus P_a)$.

Applying Lemma 5.10 to $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(\hat{\mathbb{H}}(P_a) \setminus P_a)$, we can write $\varphi_{a,s} = (\hat{l}_1 \circ \hat{P}_1)(\varphi_{a,s})$ where \hat{P}_1 is the projection operator defined by evaluating at a point $\hat{b}^1 \in \hat{\mathbb{H}}(P_a) \setminus P_a$ and \hat{l}_1 is the corresponding

embedding operator, constructed similarly as $\hat{\mathcal{P}}$ in Definition 5.41.¹⁵ By Lemma 5.43 and the above discussion, it remains to show that the leading order term of the asymptotic expansion of $-\hat{\mathcal{P}}_1(\hat{\mathcal{H}}(\tilde{\Psi}_s^{(a)} + \tilde{F}_s^{(a)} + \sum_{k \geq 0} \frac{ad^k}{\varphi_a^{s-1}} \bar{\partial} \varphi_a^{s-1})_s$ is exactly of the form ψ_a over $\hat{\mathbb{H}}(P_a) \setminus P_a$.

Choose a neighborhood $K \subset \tilde{A}_0$ of \hat{b}_1 and a family of paths ϱ_K , and using the operator $\tilde{\mathcal{L}}_{a,K}$ as defined in the proof of Lemma 5.43, we see that

$$\hat{\mathcal{P}}_1 \left(\hat{\mathcal{H}} \left(\tilde{\Psi}_s^{(a)} + \tilde{F}_s^{(a)} + \sum_{k \geq 0} \frac{ad^k}{\varphi_a^{s-1}} \bar{\partial} \varphi_a^{s-1} \right)_s \right) = \int_{\varrho_K(\cdot, \hat{b}_1)} \left(\tilde{\Psi}_s^{(a)} + \tilde{F}_s^{(a)} + \sum_{k \geq 0} \frac{ad^k}{\varphi_a^{s-1}} \bar{\partial} \varphi_a^{s-1} \right)_s.$$

Also we have

$$\hat{1}_1 \int_{\varrho_K(\cdot, \hat{b}_1)} \left(\tilde{F}_s^{(a)} + \sum_{k \geq 0} \frac{ad^k}{\varphi_a^{s-1}} \bar{\partial} \varphi_a^{s-1} \right)_s \in \bigoplus_{k \geq 1} \sum_{n \in \Lambda_{B_0}^{\vee}(U)} F_{\hat{\mathbb{H}}(P_a) \setminus P_a}^{-1}(\hat{\mathbb{H}}(P_a) \setminus P_a) \cdot z^{km_a} \check{\partial}_n t^s.$$

So it remains to compute $\int_{\varrho(\cdot, \hat{b}_1)} (\tilde{\Psi}_s^{(a)})_s$.

Lemma 5.27 together with Lemma 5.35 allow us to compute the leading order term of the integral $-\int_{\varrho(\cdot, \hat{b}_1)} \tilde{\Psi}^{(a)}$ explicitly as $-\int_{\varrho_K(\cdot, \hat{b}_1)} \tilde{\Psi}^{(a)} = \sum_k \frac{1}{2^{k-1}} \sum_{\mathcal{T} \in \text{LRT}_0^k: P_{\mathcal{T}} \neq \emptyset, m_{\mathcal{T}} \| m_a} \int_{\varrho_K(\cdot, \hat{b}_1)} \mathfrak{l}_{k, \mathcal{T}}(\check{\Pi}, \dots, \check{\Pi})$. From the discussion in Section 5.2.4, we learn that $\mathfrak{l}_{k, \mathcal{T}}(\check{\Pi}, \dots, \check{\Pi}) = \alpha_{\mathcal{T}} \check{\partial}_{n_{\mathcal{T}}} z^{m_{\mathcal{T}}} t^{j_{\mathcal{T}}}$ for each $\mathcal{T} \in \text{LRT}_0^k$. Therefore, restricting to the interval $[t_{i_0-1}, t_{i_0}]$ of $\varrho_K(\cdot, \hat{b}_1)$ and applying Lemma 5.35, we find that the \hbar order expansion of $\int_{\varrho(\cdot, \hat{b}_1)} \mathfrak{l}_{k, \mathcal{T}}(\check{\Pi}, \dots, \check{\Pi})$ is of the form $\int_{\varrho_K(\cdot, \hat{b}_1)} \mathfrak{l}_{k, \mathcal{T}}(\check{\Pi}, \dots, \check{\Pi}) \in (b_{j_{\mathcal{T}}, k_{\mathcal{T}}}^{(a)} + O(\hbar^{1/2})) z^{k_{\mathcal{T}} m_a} \check{\partial}_{n_{\mathcal{T}}} t^{j_{\mathcal{T}}}$, where $k_{\mathcal{T}} m_a = m_{\mathcal{T}}$ (here $m_{\mathcal{T}}$, $j_{\mathcal{T}}$ and $n_{\mathcal{T}}$ are introduced in Definition 5.19 and the equation (5.17)). This proves the desired result over $\hat{\mathbb{H}}(P_a) \setminus P_a$.

Over the other half-space $\tilde{A}_0 \setminus \hat{\mathbb{H}}(P_a)$, the same reason yields $\bar{\partial} \varphi_{a,s} = 0$ in $\widehat{\mathfrak{g}^* / \mathcal{E}^*}(\tilde{A}_0 \setminus \hat{\mathbb{H}}(P_a))$. Therefore we have $\varphi_{a,s} = (\hat{1} \circ \hat{\mathcal{P}})(\varphi_{a,s}) = 0$ from the gauge fixing condition in Definition 5.42, where $\hat{\mathcal{P}}$ is treated as an operator acting on $\widehat{\mathfrak{g}^* / \mathcal{E}^*}(\tilde{A}_0 \setminus \hat{\mathbb{H}}(P_a))$. \square

Now we are ready to construct the order N scattering diagram $\mathcal{D}(\Phi)_N$ for any fixed $N \in \mathbb{Z}_{>0}$. Given $a \in \mathbb{W}(N)$, Lemma 5.44 says that the leading order term in the asymptotic expansion of the gauge φ_a produces the element $\psi_a = \sum_{\substack{k \geq 1 \\ 1 \leq j \leq N}} b_{jk}^{(a)} \cdot z^{km_a} \check{\partial}_{n_a} t^j \pmod{\mathfrak{m}^{N+1}}$ over $\hat{\mathbb{H}}(P_a) \setminus P_a$.

Definition 5.45. We define the order N scattering diagram as $\mathcal{D}(\Phi)_N := \{\mathbf{w}_a \mid a \in \mathbb{W}(N)\}$, where each newly added wall \mathbf{w}_a is supported on the tropical half-hyperplane $P_a = Q - \mathbb{R}_{\geq 0} m_a \subset U$ and equipped with the wall crossing factor Θ_a defined by $\text{Log}(\Theta_a) := \sum_{\substack{k \geq 1 \\ 1 \leq j \leq N}} b_{jk}^{(a)} \cdot z^{km_a} \check{\partial}_{n_a} t^j \pmod{\mathfrak{m}^{N+1}}$.

The order $N+1$ diagram $\mathcal{D}(\Phi)_{N+1}$ is naturally an extension of the order N diagram $\mathcal{D}(\Phi)_N$ because $\Phi^{(a)}$, $\Psi^{(a)}$ and hence φ_a are defined for all orders of t . Hence this defines a scattering diagram $\mathcal{D}(\Phi)$ associated to Φ .

5.3.2. *Consistency of $\mathcal{D}(\Phi)$.* We are now ready to prove Theorem 1.5:

Theorem 5.46 (=Theorem 1.5). For the Maurer-Cartan solution Φ constructed in (5.14), the associated scattering diagram $\mathcal{D}(\Phi)$ defined in Definition 5.45 is consistent, i.e. we have the identity $\Theta_{\gamma, \mathcal{D}(\Phi)} = \prod_{\mathbf{w}_a \in \mathcal{D}(\Phi)}^{\gamma} \Theta_a = \text{Id}$ for any embedded loop γ in $U \setminus \text{Sing}(\mathcal{D}(\Phi))$ intersecting $\mathcal{D}(\Phi)$ generically.

Proof. Let us first recall that we are working over the open subset $\tilde{A}_0 = \{(r, \theta, b) \mid \theta_0 - \epsilon_0 < \theta < \theta_0 + 2\pi\}$, in the universal cover \tilde{A} of $A = U \setminus Q$, where $Q = P_1 \cap P_2 = \text{Sing}(\mathcal{D}(\Phi))$. We have also fixed a strip

¹⁵Note that $\hat{\mathcal{P}}$ and $\hat{\mathcal{P}}_1$ are defined by evaluation at two different points \hat{b}_0 and \hat{b}_1 respectively.

$\mathbb{V} = \{(r, \theta, b) \mid \theta_0 - \epsilon_0 + 2\pi < \theta < \theta_0 + 2\pi\}$ so that the strip $\mathbb{V} - 2\pi = \{(r, \theta, b) \mid \theta_0 - \epsilon_0 < \theta < \theta_0\}$ stays away from all the possible walls in $\mathcal{D}(\Phi)$; see Figure 17.




FIGURE 17.

It is enough to show that $\mathcal{D}(\Phi)_N$ is a consistent scattering diagram for each fixed $N \in \mathbb{Z}_{>0}$. Recall from Definition 5.42 that the gauge φ_a is written as $\varphi_a = -\hat{\mathcal{H}}(ad_{\varphi_a}/(e^{ad_{\varphi_a}} - \text{Id})) \left(\tilde{\Psi}^{(a)} + \tilde{F}^{(a)} \right) \pmod{\mathfrak{m}^{N+1}}$, and it satisfies the gauge fixing condition $\hat{\mathcal{P}}(\varphi_a) = 0$ and solves the equation $e^{\varphi_a} * 0 = \tilde{\Psi}^{(a)} + \tilde{F}^{(a)}$ in $\mathfrak{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$. We first show that, given any embedded loop γ in $U \setminus Q$ intersecting $\mathcal{D}(\Phi)$ generically (see Figure 17), we have

$$(5.24) \quad \prod_{a \in \mathbb{W}(N)}^{\gamma} e^{\varphi_a} = \text{Id} \pmod{\mathfrak{m}^{N+1}},$$

over \mathbb{V} .

Lemma 5.47. *Over \tilde{A}_0 , we have $\left(\prod_{a \in \mathbb{W}(N)}^{\gamma} e^{\varphi_a} \right) * 0 = \sum_{a \in \mathbb{W}(N)} \left(\tilde{\Psi}^{(a)} + \tilde{F}^{(a)} \right) \pmod{\mathfrak{m}^{N+1}}$, where the (finite) product on the left hand side is taken according to the orientation of γ .*

Proof of Lemma 5.47. By Lemma 5.44, we have $\varphi_a \equiv 0$ over the half-space $\tilde{A}_0 \setminus \hat{\mathbb{H}}(P_a) = \{(r, \theta, b) \in \tilde{A}_0 \mid \theta < \theta_a\}$ for any $a \in \mathbb{W}(N)$. So $\text{supp}(\varphi_{a'}) \cap P_a = \emptyset$ for any $a, a' \in \mathbb{W}(N)$ with $\theta_a < \theta_{a'}$ (see Figure 17). As a result we have $[\varphi_{a'}, \tilde{\Psi}^{(a)} + \tilde{F}^{(a)}] \equiv 0 \pmod{\mathfrak{m}^{N+1}}$, and we get

$$\begin{aligned} e^{\varphi_{a'}} * (\tilde{\Psi}^{(a)} + \tilde{F}^{(a)}) &= (\tilde{\Psi}^{(a)} + \tilde{F}^{(a)}) - \left(\frac{e^{ad_{\varphi_{a'}}} - \text{Id}}{ad_{\varphi_{a'}}} \right) (d\varphi_{a'} + \{\tilde{\Psi}^{(a)} + \tilde{F}^{(a)}, \varphi_{a'}\}) \\ &= \tilde{\Psi}^{(a')} + \tilde{F}^{(a')} + \tilde{\Psi}^{(a)} + \tilde{F}^{(a)} \end{aligned}$$

in $\mathfrak{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$. The lemma follows by applying this argument repeatedly according to the anti-clockwise ordering (i.e. increasing values of θ_a). \square

On the other hand, since Φ is a Maurer-Cartan solution of $\mathfrak{g}_N^*(U)/\mathcal{E}_N^*(U)$, whose deformations are all trivial in view of Lemma 5.10, we can find a gauge φ solving the equation $e^{\varphi} * 0 = \Phi$ and satisfying the condition $\mathcal{P}(\varphi) = 0$ over U . Pulling back via $\mathfrak{p}: A_0 \rightarrow A$, we get $\mathfrak{p}^*(\varphi)$ solving $e^{\mathfrak{p}^*(\varphi)} * 0 = \mathfrak{p}^*(\Phi)$

and satisfying $\hat{\mathcal{P}}(\mathbf{p}^*(\varphi)) = 0$ (the latter using the fact that $\mathbf{p}(\hat{b}_0) = x^0$ over \tilde{A}_0). Then uniqueness in Lemma 4.7 and Lemma 5.24 imply that $e^{\mathbf{p}^*(\varphi)} = \prod_{a \in \mathbb{W}(N)}^\gamma e^{\varphi_a}$ in $\mathbf{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$.

But φ is defined over the whole spherical neighborhood U , instead of just over the annulus $A = U \setminus Q$, so in fact $\prod_{a \in \mathbb{W}(N)}^\gamma e^{\varphi_a} \in \mathbf{g}_N^*(\tilde{A}_0)/\mathcal{E}_N^*(\tilde{A}_0)$ is monodromy free. In particular, this tells us that

$$\left(\prod_{a \in \mathbb{W}(N)}^\gamma e^{\varphi_a} \right) |_{\mathbb{V}-2\pi} = \left(\prod_{a \in \mathbb{W}(N)}^\gamma e^{\varphi_a} \right) |_{\mathbb{V}}$$

modulo \mathbf{m}^{N+1} . Note that \mathbb{V} is chosen so that $\mathbb{V} - 2\pi$ stays away from $\bigcup_{a \in \mathbb{W}(N)} \hat{\mathbb{H}}(P_a)$. Thus we have $\left(\prod_{a \in \mathbb{W}(N)}^\gamma e^{\varphi_a} \right) |_{\mathbb{V}-2\pi} = \text{Id} \pmod{\mathbf{m}^{N+1}}$ by Lemma 5.44, so we obtain the identity

$$(5.25) \quad \prod_{a \in \mathbb{W}(N)}^\gamma e^{\varphi_a} = \text{Id} \pmod{\mathbf{m}^{N+1}}$$

over the strip \mathbb{V} .

Equation (5.25) is an identity in the Lie algebra $\bigoplus_{m \in \Lambda_{B_0}(U)} \sum_{\substack{n \in \Lambda_{B_0}(U) \\ 1 \leq j \leq N}} F_{\mathbb{V}}^0(\mathbb{V}) \cdot z^m \check{\partial}_n t^j$. Passing to the quotient by the ideal $\bigoplus_{m \in \Lambda_{B_0}(U)} \sum_{\substack{n \in \Lambda_{B_0}(U) \\ 1 \leq j \leq N}} F_{\mathbb{V}}^{-1}(\mathbb{V}) \cdot z^m \check{\partial}_n t^j$ gives the identity

$$(5.26) \quad \prod_{a \in \mathbb{W}(N)}^\gamma e^{\psi_a} = \text{Id} \pmod{\mathbf{m}^{N+1}}$$

in $\bigoplus_{m \in \Lambda_{B_0}(U)} \sum_{\substack{n \in \Lambda_{B_0}(U) \\ 1 \leq j \leq N}} (F_{\mathbb{V}}^0(\mathbb{V})/F_{\mathbb{V}}^{-1}(\mathbb{V})) \cdot z^m \check{\partial}_n t^j$ over \mathbb{V} .

The embedding $\mathfrak{h}(\mathbb{V}) \otimes_R (\mathbf{m}/\mathbf{m}^{N+1}) \hookrightarrow \bigoplus_{m \in \Lambda_{B_0}(U)} \sum_{\substack{n \in \Lambda_{B_0}(U) \\ 1 \leq j \leq N}} F_{\mathbb{V}}^0(\mathbb{V}) \cdot z^m \check{\partial}_n t^j$, whose image has trivial intersection with $\bigoplus_{m \in \Lambda_{B_0}(U)} \sum_{\substack{n \in \Lambda_{B_0}(U) \\ 1 \leq j \leq N}} F_{\mathbb{V}}^{-1}(\mathbb{V}) \cdot z^m \check{\partial}_n t^j$ because coefficients of an element in the image are all constants independent of \hbar , so it descends to the quotient to give an embedding $\mathfrak{h}(\mathbb{V}) \otimes_R (\mathbf{m}/\mathbf{m}^{N+1}) \hookrightarrow \bigoplus_{m \in \Lambda_{B_0}(U)} \sum_{\substack{n \in \Lambda_{B_0}(U) \\ 1 \leq j \leq N}} (F_{\mathbb{V}}^0(\mathbb{V})/F_{\mathbb{V}}^{-1}(\mathbb{V})) \cdot z^m \check{\partial}_n t^j$. Therefore we obtain

$$\prod_{a \in \mathbb{W}(N)}^\gamma \Theta_a = \text{Id} \pmod{\mathbf{m}^{N+1}}$$

from (5.26) and completes the proof of the theorem. \square

5.3.3. Consistent scattering diagrams from more general Maurer-Cartan solutions. From the proof of Theorem 5.46, we observe a general relation between Maurer-Cartan solutions of the dgLa $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$ with suitable asymptotic behavior and consistent scattering diagrams in U .

We work with a contractible open coordinate chart $U \subset B_0$ and the dgLa's $\mathbf{g}_N^*(U)/\mathcal{E}_N^*(U)$ as well as $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$. We also fix a codimension 2 tropical subspace $Q \subset U$, which plays the role of the common boundary of the walls.¹⁶ In order to obtain a consistent scattering diagram from a Maurer-Cartan solution Φ of $\widehat{\mathbf{g}^*/\mathcal{E}^*}(U)$, we put two Assumptions 5.48 and 5.49 on the asymptotic behavior of Φ , the first of which is the following.

¹⁶One can regard Q as a joint in the Gross-Siebert program [28] and we are indeed considering MC solutions near a joint Q in B_0 .

Assumption 5.48. We assume that Φ admits a (Fourier) decomposition of the form

$$(5.27) \quad \Phi = \sum_{a \in \mathbb{W}} \Phi^{(a)},$$

where we have a partition of the index set \mathbb{W} into three subsets $\mathbb{W} = \mathbb{W}_{in} \sqcup \mathbb{W}_{out} \sqcup \mathbb{W}_{un}$; here the subscripts stand for incoming walls, outgoing walls and unidirectional walls respectively, following the notations in [28]. We further assume that there is an association $a \in \mathbb{W} \mapsto m_a \in M \cong \Lambda_{B_0}(U)$ satisfying m_a is not parallel to Q if $a \in \mathbb{W}_{in} \sqcup \mathbb{W}_{out}$, and m_a is parallel to Q if $a \in \mathbb{W}_{un}$ and an association

$$a \in \mathbb{W} \mapsto \text{a tropical half-hyperplane } P_a \text{ containing } Q \text{ in } U$$

satisfying $P_a = Q - \mathbb{R}_{\leq 0} \cdot m_a$ if $a \in \mathbb{W}_{in}$, $P_a = Q - \mathbb{R}_{\geq 0} \cdot m_a$ if $a \in \mathbb{W}_{out}$, and $P_a \neq P_{a'}$ if $a \neq a'$ in \mathbb{W} ,¹⁷ such that the summand $\Phi^{(a)}$ has asymptotic support on P_a and admits a decomposition $\Phi^{(a)} = \Psi^{(a)} + F^{(a)}$, where

$$\Psi^{(a)} \in \left(\bigoplus_{k \geq 1} F_{P_a}^1(U) z^{km_a} \check{\partial}_{n_a} \right) [[t]]; \quad F^{(a)} \in \left(\bigoplus_{k \geq 1} \sum_n F_{P_a}^0(U) z^{km_a} \check{\partial}_n \right) [[t]];$$

here n_a is a primitive normal to P_a .¹⁸

Under Assumption 5.48, we can solve for the gauge φ_a by the same process as in Definition 5.42 and prove the same statement as in Lemma 5.43 for each φ_a (because we have $(m_a, n_a) = 0$ even for unidirectional walls).

Next, we consider the annulus $A := U \setminus Q$ and the universal cover $\mathbf{p} : \tilde{A} \rightarrow A$, as before. We choose a reference half-hyperplane \mathcal{R}_{θ_0} of the form $\mathcal{R}_{\theta_0} = Q - \mathbb{R}_{\geq 0} m_{\theta_0}$ with $m_{\theta_0} \in M_{\mathbb{R}} \setminus M$, so that \mathcal{R}_{θ_0} cannot overlap with any of the possible walls. Again we combine polar coordinates (r, θ) on a fiber of NQ with affine coordinates $b := (b_3, \dots, b_n)$ on Q to obtain coordinates $\hat{b} = (b_1 = r, b_2 = \theta, b_3, \dots, b_n)$ on \tilde{A} . We consider the branch $\{(r, \theta, b) \mid \theta_0 < \theta < \theta_0 + 2\pi\}$, where θ_0 is a fixed angular coordinate for the half-hyperplane \mathcal{R}_{θ_0} . For each $a \in \mathbb{W}$, we let $\theta_0 < \theta_a < \theta_0 + 2\pi$ be the angular coordinate of the half-hyperplane P_a and set $\hat{\mathbb{H}}(P_a) := \{(r, \theta, b) \mid \theta_a \leq \theta < \theta_0 + 2\pi\}$.

Assumption 5.49. We assume that there exists an element $\psi_a = \sum_{j, k \geq 1} b_{jk}^{(a)} z^{km_a} \check{\partial}_{n_a} t^j$, where $b_{jk}^{(a)}$'s are constants independent of \hbar with $b_{jk}^{(a)} \neq 0$ only for finitely many k 's for each fixed j and n_a is a primitive normal to P_a , such that

$$\hat{\mathcal{H}}(\Psi^{(a)})|_{\hat{\mathbb{H}}(P_a) \setminus P_a} \in \psi_a + \left(\bigoplus_{k \geq 1} F_{\hat{\mathbb{H}}(P_a) \setminus P_a}^{-1}(\hat{\mathbb{H}}(P_a) \setminus P_a) \cdot z^{km_a} \check{\partial}_{n_a} \right) [[t]] \cdot t.$$

For each fixed $N \in \mathbb{Z}_{>0}$, we choose a sufficiently small $\epsilon_N > 0$ such that the subset $\mathbb{V}_N := \{(r, \theta, b) \mid \theta_0 - \epsilon_N + 2\pi < \theta < \theta_0 + 2\pi\}$ is disjoint from all the P_a 's. We then restrict our attention to $\tilde{A}_0 := \{(r, \theta, b) \mid \theta_0 - \epsilon_N < \theta < \theta_0 + 2\pi\}$ in order to apply a monodromy argument as in Section 5.3.1. We also fix the homotopy operator $\hat{\mathcal{H}}$ as in Definition 5.41, together with $\hat{\mathcal{P}}$ and \hat{i} . Then we can prove the same statement as in Lemma 5.44 under Assumption 5.49.

So altogether, assuming both Assumptions 5.48 and 5.49, we have Lemmas 5.43 and 5.44, and a scattering diagram $\mathcal{D}(\Phi)$ can be associated to the given Maurer-Cartan solution Φ in exactly the same way as in Definition 5.45. Finally, the same proof as in Theorem 5.46 gives the following:

Theorem 5.50 (= Theorem 1.4). Suppose that we have a Maurer-Cartan solution Φ of $\widehat{\mathfrak{g}^*/\mathcal{E}^*}(U)$ satisfying both Assumptions 5.48 and 5.49. Then the scattering diagram $\mathcal{D}(\Phi)$ associated to Φ is

¹⁷Note that there is no restriction on P_a if $a \in \mathbb{W}_{un}$, hence the name *unidirectional walls*.

¹⁸Note that we do not need to specify the sign of (ν_{P_a}, n_a) in this assumption.

consistent, i.e. we have the following identity $\Theta_{\gamma, \mathcal{D}(\Phi)} = \prod_{\mathbf{w}_a \in \mathcal{D}(\Phi)}^{\gamma} \Theta_a = \text{Id}$ along any embedded loop γ in $U \setminus \text{Sing}(\mathcal{D}(\Phi))$ intersecting $\mathcal{D}(\Phi)$ generically.

REFERENCES

1. M. Abouzaid, *Homogeneous coordinate rings and mirror symmetry for toric varieties*, *Geom. Topol.* **10** (2006), 1097–1157 (electronic). MR 2240909 (2007h:14052)
2. ———, *Morse homology, tropical geometry, and homological mirror symmetry for toric varieties*, *Selecta Math. (N.S.)* **15** (2009), no. 2, 189–270. MR 2529936 (2011h:53123)
3. M. Abouzaid, D. Auroux, and L. Katzarkov, *Lagrangian fibrations on blowups of toric varieties and mirror symmetry for hypersurfaces*, *Publ. Math. Inst. Hautes Études Sci.* **123** (2016), 199–282. MR 3502098
4. P. Aspinwall, T. Bridgeland, A. Craw, M. R. Douglas, M. Gross, A. Kapustin, G. W. Moore, G. Segal, B. Szendrői, and P. M. H. Wilson, *Dirichlet branes and mirror symmetry*, *Clay Mathematics Monographs*, vol. 4, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2009. MR 2567952 (2011e:53148)
5. D. Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, *J. Gökova Geom. Topol. GGT* **1** (2007), 51–91. MR 2386535 (2009f:53141)
6. ———, *Special Lagrangian fibrations, wall-crossing, and mirror symmetry*, *Surveys in differential geometry. Vol. XIII. Geometry, analysis, and algebraic geometry: forty years of the Journal of Differential Geometry*, *Surv. Differ. Geom.*, vol. 13, Int. Press, Somerville, MA, 2009, pp. 1–47. MR 2537081 (2010j:53181)
7. T. Bridgeland and V. Toledano Laredo, *Stability conditions and Stokes factors*, *Invent. Math.* **187** (2012), no. 1, 61–98. MR 2874935
8. K. Chan, C.-H. Cho, S.-C. Lau, and H.-H. Tseng, *Gross fibrations, SYZ mirror symmetry, and open Gromov-Witten invariants for toric Calabi-Yau orbifolds*, *Journal of Differential Geometry* **103** (2016), no. 2, 207–288.
9. K. Chan, S.-C. Lau, and N. C. Leung, *SYZ mirror symmetry for toric Calabi-Yau manifolds*, *J. Differential Geom.* **90** (2012), no. 2, 177–250. MR 2899874
10. K. Chan and N. C. Leung, *Mirror symmetry for toric Fano manifolds via SYZ transformations*, *Adv. Math.* **223** (2010), no. 3, 797–839. MR 2565550 (2011k:14047)
11. ———, *On SYZ mirror transformations*, *New developments in algebraic geometry, integrable systems and mirror symmetry (RIMS, Kyoto, 2008)*, *Adv. Stud. Pure Math.*, vol. 59, Math. Soc. Japan, Tokyo, 2010, pp. 1–30. MR 2683205 (2011g:53186)
12. K.-L. Chan, N. C. Leung, and Z. N. Ma, *Fukaya’s conjecture on Witten’s twisted A_{∞} structure*, *J. Differential Geom.*, to appear, arXiv:1401.5867.
13. C.-H. Cho, *Products of Floer cohomology of torus fibers in toric Fano manifolds*, *Comm. Math. Phys.* **260** (2005), no. 3, 613–640. MR 2183959 (2006h:53094)
14. C.-H. Cho and Y.-G. Oh, *Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds*, *Asian J. Math.* **10** (2006), no. 4, 773–814. MR 2282365 (2007k:53150)
15. M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, *London Mathematical Society Lecture Note Series*, vol. 268, Cambridge University Press, Cambridge, 1999. MR 1735654
16. J. J. Duistermaat, *On global action-angle coordinates*, *Comm. Pure Appl. Math.* **33** (1980), no. 6, 687–706. MR 596430 (82d:58029)
17. B. Fang, *Homological mirror symmetry is T-duality for \mathbb{P}^n* , *Commun. Number Theory Phys.* **2** (2008), no. 4, 719–742. MR 2492197 (2010f:53154)
18. B. Fang, C.-C. M. Liu, D. Treumann, and E. Zaslow, *T-duality and homological mirror symmetry for toric varieties*, *Adv. Math.* **229** (2012), no. 3, 1875–1911. MR 2871160
19. S. A. Filippini, M. Garcia-Fernandez, and J. Stoppa, *Stability data, irregular connections and tropical curves*, *Selecta Math. (N.S.)* **23** (2017), no. 2, 1355–1418. MR 3624914
20. K. Fukaya, *Deformation theory, homological algebra and Mirror Symmetry*, *Geometry and physics of branes (Como, 2001)*, *Ser. High Energy Phys. Cosmol. Gravit.*, IOP Bristol (2003), 121–209.
21. ———, *Multivalued Morse theory, asymptotic analysis and mirror symmetry*, *Graphs and patterns in mathematics and theoretical physics*, *Proc. Sympos. Pure Math.*, vol. 73, Amer. Math. Soc., Providence, RI, 2005, pp. 205–278. MR 2131017 (2006a:53100)
22. K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian Floer theory on compact toric manifolds. I*, *Duke Math. J.* **151** (2010), no. 1, 23–174. MR 2573826 (2011d:53220)
23. ———, *Lagrangian Floer theory on compact toric manifolds II: bulk deformations*, *Selecta Math. (N.S.)* **17** (2011), no. 3, 609–711. MR 2827178
24. ———, *Lagrangian Floer theory and mirror symmetry on compact toric manifolds*, *Astérisque* (2016), no. 376, vi+340. MR 3460884
25. M. Gross, *Examples of special Lagrangian fibrations*, *Symplectic geometry and mirror symmetry (Seoul, 2000)*, *World Sci. Publ.*, River Edge, NJ, 2001, pp. 81–109. MR 1882328 (2003f:53085)

26. ———, *Topological mirror symmetry*, Invent. Math. **144** (2001), no. 1, 75–137. MR 1821145 (2002c:14062)
27. M. Gross, R. Pandharipande, and B. Siebert, *The tropical vertex*, Duke Math. J. **153** (2010), no. 2, 297–362. MR 2667135 (2011f:14093)
28. M. Gross and B. Siebert, *From real affine geometry to complex geometry*, Ann. of Math. (2) **174** (2011), no. 3, 1301–1428. MR 2846484
29. ———, *Local mirror symmetry in the tropics*, Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II, Kyung Moon Sa, Seoul, 2014, pp. 723–744. MR 3728635
30. B. Helffer and J. Sjöstrand, *Multiple wells in the semi-classical limit I*, Comm. in PDE **9** (1984), no. 4, 337–408.
31. ———, *Puits multiples en limite semi-classique IV - Etude du complexe de Witten*, Comm. in PDE **10** (1985), no. 3, 245–340.
32. N. J. Hitchin, *The moduli space of special Lagrangian submanifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **25** (1997), no. 3-4, 503–515 (1998), Dedicated to Ennio De Giorgi. MR 1655530
33. K. Hori, A. Iqbal, and C. Vafa, *D-branes and mirror symmetry*, preprint (2000), arXiv:hep-th/0005247.
34. K. Hori and C. Vafa, *Mirror symmetry*, preprint (2000), arXiv:hep-th/0002222.
35. K. Kodaira and J. Morrow, *Complex manifolds*, AMS Chelsea Publishing, Providence, RI, 2006, Reprint of the 1971 edition with errata. MR 2214741 (2006j:32001)
36. M. Kontsevich and Y. Soibelman, *Homological mirror symmetry and torus fibrations*, Symplectic geometry and mirror symmetry (Seoul, 2000), World Sci. Publ., River Edge, NJ, 2001, pp. 203–263. MR 1882331 (2003c:32025)
37. ———, *Affine structures and non-Archimedean analytic spaces*, The unity of mathematics, Progr. Math., vol. 244, Birkhäuser Boston, Boston, MA, 2006, pp. 321–385. MR 2181810 (2006j:14054)
38. M. Kuranishi, *New proof for the existence of locally complete families of complex structures*, Proc. Conf. Complex Analysis (Minneapolis, 1964), Springer, Berlin, 1965, pp. 142–154. MR 0176496
39. S.-C. Lau, *Gross-Siebert's slab functions and open GW invariants for toric Calabi-Yau manifolds*, Math. Res. Lett. **22** (2015), no. 3, 881–898. MR 3350109
40. N. C. Leung, *Mirror symmetry without corrections*, Comm. Anal. Geom. **13** (2005), no. 2, 287–331. MR 2154821 (2006c:32028)
41. N. C. Leung and C. Vafa, *Branes and toric geometry*, Adv. Theor. Math. Phys. **2** (1998), no. 1, 91–118. MR 1635926 (99f:81170)
42. Z. N. Ma, *SYZ mirror symmetry from Witten-Morse theory*, Gromov-Witten theory, gauge theory and dualities, Proc. Centre Math. Appl. Austral. Nat. Univ., vol. 48, Austral. Nat. Univ., Canberra, 2019, arXiv:1811.09042, p. 36.
43. M. Manetti, *Differential graded Lie algebras and formal deformation theory*, Algebraic geometry—Seattle 2005. Part 2, Proc. Sympos. Pure Math., vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 785–810. MR 2483955 (2009m:17015)
44. M. Reineke, *Poisson automorphisms and quiver moduli*, J. Inst. Math. Jussieu **9** (2010), no. 3, 653–667. MR 2650811
45. A. Strominger, S.-T. Yau, and E. Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B **479** (1996), no. 1-2, 243–259. MR 1429831 (97j:32022)
46. E. Witten, *Supersymmetry and Morse theory*, J. Differential Geom. **17** (1982), no. 4, 661–692 (1983). MR 683171 (84b:58111)

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

Email address: kwchan@math.cuhk.edu.hk

THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

Email address: leung@math.cuhk.edu.hk

THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

Email address: zmma@math.cuhk.edu.hk